



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



## RIVINGTON'S MATHEMATICAL SERIES

---

By J. HAMBLIN SMITH, M.A.,

GONVILLE AND CAIUS COLLEGE, AND LATE LECTURER AT ST. PETER'S  
COLLEGE, CAMBRIDGE.

*Arithmetic.* 3s. 6d. A KEY, 9s.

*Elementary Algebra.* 3s. Without Answers, 2s. 6d.  
A KEY, 9s.

*Exercises on Algebra.* 2s. 6d.  
[Copies may be had without the Answers.]

*Elementary Trigonometry.* 4s. 6d. A KEY, 7s. 6d.

*Elements of Geometry.*

Containing Books 1 to 6, and portions of Books 11 and 12 of  
EUCLID, with Exercises and Notes. 3s. 6d. A KEY, 8s. 6d.  
Books 1 and 2, 1s. 6d., may be had separately.

*Elementary Statics.* 3s. }  
*Elementary Hydrostatics.* 3s. } A KEY, 6s.

*Book of Enunciations.*

FOR HAMBLIN SMITH'S GEOMETRY, ALGEBRA, TRIGONOMETRY,  
STATICS, AND HYDROSTATICS. 1s.

*An Introduction to the Study of Heat.* 3s.

---

By E. J. GROSS, M.A.,

FELLOW OF GONVILLE AND CAIUS COLLEGE, CAMBRIDGE.

*Algebra.* - Part II. 8s. 6d.

*Kinematics and Kinetics.* 5s. 6d.

---

By G. RICHARDSON, M.A.,

ASSISTANT MASTER AT WINCHESTER COLLEGE.

*Geometrical Conic Sections.* 4s. 6d.

---

Waterloo Place, Pall Mall, London.

## ENGLISH SCHOOL CLASSICS

EDITED BY FRANCIS STORR, B.A.,

CHIEF MASTER OF MODERN SUBJECTS IN MERCHANT TAYLORS' SCHOOL.

- THOMSON'S SEASONS: Winter.**  
With an Introduction to the Series. By the Rev. J. F. RUGHT, M.A. 12.
- COWPER'S TASK.**  
By FRANCIS STORR, B.A. 2s.; or in Three Parts, 9d. each.
- COWPER'S SIMPLE POEMS.**  
By FRANCIS STORR, B.A. 1s.
- SCOTT'S LAY OF THE LAST MINSTREL.**  
By J. SURTEES PHILLIPOTS, M.A., Head Master of Bedford School. 2s. 6d.;  
or in Four Parts, 9d. each.
- SCOTT'S LADY OF THE LAKE.**  
By R. W. TAYLOR, M.A., Head Master of Kelly College, Tavistock. 2s.  
or in Three Parts, 9d. each.
- SCOTT'S MARMION.**  
By F. S. ARNOLD, M.A., Assistant Master at Bedford School.
- NOTES TO SCOTT'S WAVERLEY.**  
By H. W. EVE, M.A., Head Master of University College School, London.  
1s.; WAVERLEY AND NOTES, 2s. 6d.
- BACON'S ESSAYS. Complete Edition.**  
By FRANCIS STORR, B.A.
- TWENTY-OF BACON'S ESSAYS.**  
By FRANCIS STORR, B.A. 1s.
- SIMPLE POEMS.**  
By W. M. MULLINS, M.A., Assistant Master at Marlborough College. 2d.
- SELECTIONS FROM WORDSWORTH'S POEMS.**  
By H. M. TURNER, B.A., late Scholar of Trinity College, Cambridge. 1s.
- WORDSWORTH'S EXCURSION: The Wanderer.**  
By H. M. TURNER, B.A. 1s.
- MILTON'S PARADISE LOST.**  
By FRANCIS STORR, B.A. Book I., 5d. Book II., 9d.
- MILTON'S L'ALLEGRO, IL PENSEROSO, AND LYCIDAS.**  
By FRANCIS STORR, M.A., late Scholar of New College, Oxford. 1s.
- SELECTIONS FROM THE SPECTATOR.**  
By OSMAIR ALBY, M.A., late Assistant Master at Wellington College. 1s.
- BROWNE'S RELIGIO MEDICI.**  
By W. P. SMITH, M.A., Assistant Master at Winchester College. 1s.
- GOLDSMITH'S TRAVELLER AND DESERTED VILLAGE.**  
By C. SANKEY, M.A., Head Master of Bury St. Edmund's Grammar  
School. 1s.
- EXTRACTS from GOLDSMITH'S VICAR OF WAKEFIELD.**  
By C. SANKEY, M.A. 1s.
- POEMS SELECTED from the WORKS OF ROBERT BURNS.**  
By A. M. BELL, M.A., Balliol College, Oxford. 2s.
- MACAULAY'S ESSAYS:**  
**MOORE'S LIFE OF BYRON.** By FRANCIS STORR, B.A. 9d.  
**BOSWELL'S LIFE OF JOHNSON.** By FRANCIS STORR, B.A. 9d.  
**MALLAM'S CONSTITUTIONAL HISTORY.** By H. F. BOND, late  
Scholar of Brasenose College, Oxford. 1s.
- SOUTHEW'S LIFE OF NELSON.**  
By W. E. MULLINS, M.A. 2s. 6d.
- GRAY'S POEMS. SELECTION FROM LETTERS, with LIFE**  
by JOHNSON. By FRANCIS STORR, B.A. 1s.

---

Waterloo Place, Pall Mall, London.



600047144Q

A KEY  
TO  
ELEMENTARY TRIGONOMETRY


BY  
J. HAMBLIN SMITH, M.A.  
OF GONVILLE AND CAIUS COLLEGE  
AND LATE LECTURER AT ST. PETER'S COLLEGE, CAMBRIDGE

*THIRD EDITION*

RIVINGTONS  
WATERLOO PLACE, LONDON

MDCCCLXXXV

18381. E. 2.



## PREFACE.

I HAVE to acknowledge most gratefully the assistance rendered me in the preparation of this book by Mr. T. H. Gascoigne, son of the Rev. T. Gascoigne, of Spondon House School, Derby. For the solutions of a few of the Problems I am indebted to Mr. Gaskin's *Trigonometrical Examples*, and to Mr. Hymers' *Trigonometry*. I shall be glad to receive corrections of errors that may be discovered in my work.

CAMBRIDGE, *October 1876.*





# ELEMENTARY TRIGONOMETRY.

## KEY.

### EXAMPLES—I. (pp. 1, 2).

- (1) 4 feet 6 inches = 54 inches ;  $\therefore$  number is 54.
- (2) 15 feet 2 inches = 182 inches ;  $\therefore$  number is  $182 \div 7$ , or, 26.
- (3) Unit of square measurement is  $(192 \div 12)$  square inches, or, 16 square inches ;  $\therefore$  unit of linear measurement is  $\sqrt{16}$  inches, or, 4 inches.
- (4) Unit of square measurement is  $(1000 \div 40)$  square inches, or, 25 square inches ;  $\therefore$  unit of linear measurement is  $\sqrt{25}$  inches, or, 5 inches.
- (5) Unit of cubic measurement is  $(216 \div 8)$  cubic inches, or, 27 cubic inches ;  $\therefore$  unit of linear measurement is  $\sqrt[3]{27}$  inches, or, 3 inches.
- (6) Unit of cubic measurement is  $(2000 \div 16)$  cubic inches, or, 125 cubic inches ;  $\therefore$  unit of linear measurement is  $\sqrt[3]{125}$  inches, or, 5 inches.
- (7) Measure of 1 yard is  $\frac{1}{a}$  ;  
 $\therefore$  measure of 1 foot is  $\frac{1}{3a}$  ;  
 $\therefore$  measure of  $b$  feet is  $\frac{b}{3a}$ .

#### 4 KEY TO ELEMENTARY TRIGONOMETRY.

(13) Taking the diagram in Example 12, let measure of  $AB$  be  $x$ .

$$\text{Then } x^2 = \frac{x^2}{4} + (15)^2;$$

$$\therefore 3x^2 = 4 \times (15)^2, \text{ or, } x^2 = \frac{4 \times (15)^2}{3} = \frac{4 \times (15)^2 \times 3}{9};$$

$$\therefore x = \frac{2 \times 15 \sqrt{3}}{3} = 10 \sqrt{3}.$$




FIG. 3.

(14)  $OD$ , a perpendicular from the centre on the chord  $AB$ , bisects  $AB$ .

Let  $x$  = measure of  $OD$  in inches.

$$\begin{aligned} \text{Then } x^2 &= (OA)^2 - (AD)^2 \\ &= (37)^2 - (35)^2 = 144; \end{aligned}$$

$$\therefore \text{distance} = \sqrt{144} \text{ inches} = 12 \text{ inches.}$$

(15) Taking the diagram of Example (14).

Let measure of  $AD$  in inches be  $x$ .

$$\text{Then } x^2 = (181)^2 - (180)^2 = 361;$$

$$\therefore x = 19, \text{ and } \therefore AB = (2 \times 19) \text{ inches} = 38 \text{ inches.}$$

(16) Taking the diagram of Example (14).

Let measure of  $AO$  in feet be  $x$ .

$$\text{Then } x^2 = (308)^2 + (75)^2 = 100489;$$

$$\therefore x = 317, \text{ and } \therefore \text{diameter} = (2 \times 317) \text{ feet} = 634 \text{ feet.}$$




FIG. 4.

$$(17) (AC)^2 + (BC)^2 = (AB)^2.$$

$$\therefore 2(AC)^2 = (AB)^2;$$

$$\therefore \frac{(AC)^2}{(AB)^2} = \frac{1}{2};$$

$$\therefore \frac{AC}{AB} = \frac{1}{\sqrt{2}}.$$




FIG. 5.

(18) Let  $x$  be the measure of  $EG$ .

Then  $2x$  is the measure of  $ED$ ;

and measure of  $DG = \sqrt{(4x^2 - x^2)} = \sqrt{3} \cdot x$ ;

$$\therefore EG : ED : DG = x : 2x : \sqrt{3} \cdot x$$

$$= 1 : 2 : \sqrt{3}.$$

EXAMPLES—III. (p. 9).

$$(1) \text{ Circumference} = \frac{22 \times 5}{7} \text{ feet} = \frac{110}{7} \text{ feet} = 15\frac{5}{7} \text{ feet.}$$

$$(2) \text{ Radius} = \frac{7 \times 542\frac{5}{44}}{44} \text{ feet} = \frac{3797\frac{5}{44}}{44} \text{ feet} = 86\cdot306\frac{1}{4} \text{ feet.}$$

$$(3) \text{ Train goes in a second } \frac{22 \times 12}{7} \text{ feet.}$$

$$\text{Rate in miles per hour} = \frac{22 \times 12 \times 60 \times 60}{7 \times 3 \times 1760} = \frac{180}{7} = 25\cdot71428\frac{5}{7}.$$

$$(4) \text{ Diameter in miles} = \frac{7 \times 25000}{22} = 7954\frac{6}{11}.$$

$$(5) \text{ Circumference in miles} = \frac{22 \times 883220}{7} = 2775834\frac{2}{7}.$$

$$(6) \text{ Radius in miles} = \frac{7 \times 6850}{44} = \frac{23975}{22} = 1089\frac{17}{22}.$$

$$(7) \text{ Circumference in feet} = \frac{22 \times 12\frac{1}{2} \times 2}{7} = \frac{22 \times 25}{7};$$

$$\therefore \frac{1}{12} \text{ of circumference} = \frac{22 \times 25}{12 \times 7} \text{ feet} = 6 \text{ feet } 6\frac{1}{4} \text{ inches.}$$

$$(8) \text{ Circumference in feet} = \frac{22 \times 21}{7};$$

$$\therefore \frac{5}{7} \text{ of circumference} = \frac{22 \times 21 \times 5}{7 \times 7} \text{ feet} = 47\frac{1}{7} \text{ feet.}$$

$$(9) \text{ If } x \text{ be the side of the square,}$$

$$(\text{diameter})^2 = 2x^2;$$

$$\therefore x = \frac{\text{diameter}}{\sqrt{2}} = \frac{7 \times 150}{22 \times \sqrt{2}} \text{ feet} = \frac{7 \times 150 \times \sqrt{2}}{22 \times \sqrt{2} \times \sqrt{2}} \text{ feet} = \frac{525\sqrt{2}}{22} \text{ feet.}$$

$$(10) x = \frac{\text{diameter}}{\sqrt{2}} = \frac{7 \times 200}{22 \times \sqrt{2}} \text{ feet} = \frac{7 \times 200 \times \sqrt{2}}{22 \times 2} \text{ feet} = \frac{350\sqrt{2}}{11} \text{ feet.}$$

$$(11) \text{ Point goes in a minute } \frac{22 \times 12 \times 30}{7} \text{ feet.}$$

$$\text{Rate in miles per hour} = \frac{22 \times 12 \times 30 \times 60}{7 \times 1760 \times 3} = \frac{90}{7} = 12\frac{6}{7}.$$

## 6 KEY TO ELEMENTARY TRIGONOMETRY.

(12) End goes in a minute  $\frac{22 \times 2 \times 15 \times 21}{7}$  feet.

$$\text{Rate in miles per hour} = \frac{22 \times 2 \times 15 \times 21 \times 60}{7 \times 3 \times 1760} = \frac{45}{2} = 22\frac{1}{2}.$$

### EXAMPLES—IV. (p. 12).

$$\begin{array}{r} (1) \quad 60 \quad \overline{) 5} \\ 60 \quad \overline{) 16'08\ddot{3}} \\ \hline \quad \quad \quad \cdot 2680\ddot{5} \end{array}$$

$$\therefore 24^\circ. 16'. 5'' = 24^\circ. 2680\ddot{5}$$

$$\begin{array}{r} (2) \quad 60 \quad \overline{) 43} \\ 60 \quad \overline{) 2'71\ddot{6}} \\ \hline \quad \quad \quad \cdot 0452\ddot{7} \end{array}$$

$$\therefore 37^\circ. 2'. 43'' = 37^\circ. 0452\ddot{7}$$

$$\begin{array}{r} (3) \quad 60 \quad \overline{) 14} \\ 60 \quad \overline{) 0'2\ddot{3}} \\ \hline \quad \quad \quad \cdot 003\ddot{8} \end{array}$$

$$\therefore 175^\circ. 0'. 14'' = 175^\circ. 003\ddot{8}.$$

$$\begin{array}{r} (4) \quad 60 \quad \overline{) 28} \\ 60 \quad \overline{) 5'4\ddot{6}} \\ \hline \quad \quad \quad \cdot 09\ddot{1} \end{array}$$

$$\therefore 5'. 28'' = 09\ddot{1}.$$

$$\begin{array}{r} (5) \quad 60 \quad \overline{) 4} \\ \hline \quad \quad \quad \cdot 0\ddot{6} \end{array}$$

$$\therefore 375^\circ. 4' = 375^\circ. 0\ddot{6}.$$

$$\begin{array}{r} (6) \quad 60 \quad \overline{) 4} \\ 60 \quad \overline{) 12'0\ddot{6}} \\ \hline \quad \quad \quad \cdot 20\ddot{1} \end{array}$$

$$\therefore 78^\circ. 12'. 4'' = 78^\circ. 20\ddot{1}.$$

### EXAMPLES—V. (p. 13).

$$(1) \quad 25^\circ. 14'. 25'' = 25^\circ. 1425.$$

$$(4) \quad 15'. 7''. 45 = 150745.$$

$$(2) \quad 38^\circ. 4'. 15'' = 38^\circ. 0415.$$

$$(5) \quad 425^\circ. 13'. 5''. 54 = 425^\circ. 130554.$$

$$(3) \quad 214^\circ. 3'. 7'' = 214^\circ. 0307.$$

$$(6) \quad 2^\circ. 2'. 2''. 22 = 2^\circ. 020222.$$

### EXAMPLES—VI. (p. 19).

$$(1) \quad 27^\circ. 15'. 46'' = 27^\circ. 262\ddot{7}$$

$$\begin{array}{r} 10 \\ \overline{) 272'62\ddot{7}} \end{array}$$

$$30'291975 \dots$$

$$\therefore 27^\circ. 15'. 46'' = 30^\circ. 29'. 19''. 75 \dots$$

$$\begin{array}{r}
 (2) \ 157^{\circ}.4'.9'' = 157^{\circ}.0691\bar{6} \\
 \qquad \qquad \qquad 10 \\
 9 \overline{) 1570.691\bar{6}} \\
 \qquad \qquad \qquad 1745212\bar{9}6\bar{2} \\
 \therefore 157^{\circ}.4'.9'' = 174^{\circ}.52'.12''.\bar{9}6\bar{2}.
 \end{array}$$

$$\begin{array}{r}
 (3) \ 24'.18'' = 0^{\circ}.405 \\
 \qquad \qquad \qquad 10 \\
 9 \overline{) 4.05} \\
 \qquad \qquad \qquad 45 \\
 \therefore 24'.18'' = 45'.
 \end{array}$$

$$\begin{array}{r}
 (4) \ 19^{\circ}.0'.18'' = 19^{\circ}.005 \\
 \qquad \qquad \qquad 10 \\
 9 \overline{) 190.05} \\
 \qquad \qquad \qquad 21.116\bar{6}\bar{8} \\
 \therefore 19^{\circ}.0'.18'' = 21^{\circ}.11'.66''.\bar{6}.
 \end{array}$$

$$\begin{array}{r}
 (5) \ 143^{\circ}.9' = 143^{\circ}.15 \\
 \qquad \qquad \qquad 10 \\
 9 \overline{) 1431.5} \\
 \qquad \qquad \qquad 159.055\bar{5}\bar{5} \\
 \therefore 143^{\circ}.9' = 159^{\circ}.5'.55''.\bar{5}.
 \end{array}$$

$$\begin{array}{r}
 (6) \ 28^{\circ} = 28^{\circ} \\
 \qquad \qquad \qquad 10 \\
 9 \overline{) 280} \\
 \qquad \qquad \qquad 31.111\bar{1}\bar{1} \\
 \therefore 28^{\circ} = 31^{\circ}.11'.11''.\bar{1}.
 \end{array}$$

$$\begin{array}{r}
 (7) \ 10^{\circ}.25'.48'' = 10^{\circ}.43 \\
 \qquad \qquad \qquad 10 \\
 9 \overline{) 104.3} \\
 \qquad \qquad \qquad 11.588\bar{8}\bar{8} \\
 \therefore 10^{\circ}.25'.48'' = 11^{\circ}.58'.88''.\bar{8}.
 \end{array}$$

# 8 KEY TO ELEMENTARY TRIGONOMETRY.

$$(8) 27^{\circ}.38'.12'' = 27^{\circ}.63\dot{6}$$

$$\begin{array}{r} 10 \\ 9 \overline{) 276.3\dot{6}} \\ 30.7\dot{0}7\dot{1} \end{array}$$

$$\therefore 27^{\circ}.38'.12'' = 30^{\circ}.70'.74''.\dot{0}7\dot{1}.$$

$$(9) 300^{\circ}.15'.58'' = 300^{\circ}.266\dot{1}$$

$$\begin{array}{r} 10 \\ 9 \overline{) 3002.66\dot{1}} \\ 333.629012345679\dot{0} \end{array}$$

$$\therefore 300^{\circ}.15'.58'' = 333^{\circ}.62'.90''.\dot{1}2345679\dot{0}$$

$$(10) 422^{\circ}.7'.22'' = 422^{\circ}.122\dot{7}$$

$$\begin{array}{r} 10 \\ 9 \overline{) 4221.22\dot{7}} \\ 469.0253\dot{0}8641975\dot{3} \end{array}$$

$$\therefore 422^{\circ}.7'.22'' = 469^{\circ}.2'.53''.\dot{0}8641975\dot{3}.$$

## EXAMPLES—VII (p. 20).

$$(1) 19^{\circ}.45'.95'' = 19^{\circ}.4595$$

$$\begin{array}{r} 9 \\ 10 \overline{) 175.135\dot{5}} \\ \text{degrees} \quad 17.51355 \\ \quad \quad \quad 60 \\ \text{minutes} \quad 30.81300 \\ \quad \quad \quad 60 \\ \text{seconds} \quad 48.78000 \end{array}$$

$$\therefore 19^{\circ}.45'.95'' = 17^{\circ}.30'.48''.78.$$

$$(2) 124^{\circ}.5'.8'' = 124^{\circ}.0508$$

$$\begin{array}{r} 9 \\ 10 \overline{) 1116.457\dot{2}} \\ \text{degrees} \quad 111.64572 \\ \quad \quad \quad 60 \\ \text{minutes} \quad 38.4320 \\ \quad \quad \quad 60 \\ \text{seconds} \quad 44.59200 \end{array}$$

$$\therefore 124^{\circ}.5'.8'' = 111^{\circ}.38'.44''.592.$$

$$(3) 29^{\circ}.75' = 29^{\circ}.75$$

$$\begin{array}{r} 9 \\ 10 \overline{) 267.75} \\ \text{degrees} \quad 26.775 \\ \quad 60 \\ \text{minutes} \quad 46.500 \\ \quad 60 \\ \text{seconds} \quad 30.000 \\ \therefore 29^{\circ}.75' = 26^{\circ}.46'.30''. \end{array}$$

$$(4) 15^{\circ}.0'.15'' = 15^{\circ}.0015$$

$$\begin{array}{r} 9 \\ 10 \overline{) 135.0135} \\ \text{degrees} \quad 13.50135 \\ \quad 60 \\ \text{minutes} \quad 30.08100 \\ \quad 60 \\ \text{seconds} \quad 4.86000 \\ \therefore 15^{\circ}.0'.15'' = 13^{\circ}.30'.4''.86. \end{array}$$

$$(5) 154^{\circ}.7'.24'' = 154^{\circ}.0724$$

$$\begin{array}{r} 9 \\ 10 \overline{) 1386.6516} \\ \text{degrees} \quad 138.66516 \\ \quad 60 \\ \text{minutes} \quad 39.90960 \\ \quad 60 \\ \text{seconds} \quad 54.57600 \\ \therefore 154^{\circ}.7'.24'' = 138^{\circ}.39'.54''.576. \end{array}$$

$$(6) 43^{\circ} = 43^{\circ}$$

$$\begin{array}{r} 9 \\ 10 \overline{) 387} \\ \text{degrees} \quad 38.7 \\ \quad 60 \\ \text{minutes} \quad 42.0 \\ \therefore 43^{\circ} = 38^{\circ}.42'. \end{array}$$



$$(7) 38^{\circ}.71'.20'' \cdot 3 = 38^{\circ}.71203$$

	9
	10   348·40827
degrees	34·840827
	60
minutes	50·449620
	60
seconds	26·977200

$$\therefore 38^{\circ}.71'.20'' \cdot 3 = 34^{\circ}.50'.26'' \cdot 9772.$$

$$(8) 50^{\circ}.76'.94'' \cdot 3 = 50^{\circ}.76943$$

	9
	10   456·92487
degrees	45·692487
	60
minutes	41·549220
	60
seconds	32·953200

$$\therefore 50^{\circ}.76'.94'' \cdot 3 = 45^{\circ}.41'.32'' \cdot 9532.$$

$$(9) 170^{\circ}.63'.27'' = 170^{\circ}.6327$$

	9
	10   1535·6943
degrees	153·56943
	60
minutes	34·16580
	60
seconds	9·94800

$$\therefore 170^{\circ}.63'.27'' = 153^{\circ}.34'.9'' \cdot 948.$$

$$(10) 324^{\circ}.13'.88'' \cdot 7 = 324^{\circ}.13887$$

	9
	10   2917·24983
degrees	291·724983
	60
minutes	43·498980
	60
seconds	29·938800

$$\therefore 324^{\circ}.13'.88'' \cdot 7 = 291^{\circ}.43'.29'' \cdot 9388$$

## EXAMPLES—VIII. (p. 21).

(1) Circular measure is  $\frac{60 \times \pi}{180} = \frac{\pi}{3}$ .

(2) Circular measure is  $\frac{22.5 \times \pi}{180} = \frac{\pi}{8}$ .

(3) Circular measure is  $\frac{11.25 \times \pi}{180} = \frac{\pi}{16}$ .

(4) Circular measure is  $\frac{270 \times \pi}{180} = \frac{3\pi}{2}$ .

(5) Circular measure is  $\frac{315 \times \pi}{180} = \frac{7\pi}{4}$ .

(6) Circular measure is  $\frac{241\frac{3}{4} \times \pi}{180} = \frac{1453\pi}{60 \times 180} = \frac{1453\pi}{10800}$ .

(7) Circular measure is  $\frac{95\frac{1}{3} \times \pi}{180} = \frac{286 \times \pi}{180 \times 3} = \frac{143\pi}{270}$ .

(8) Circular measure is  $\frac{12\frac{304}{5600} \times \pi}{180} = \frac{43504 \times \pi}{180 \times 3600} = \frac{2719\pi}{40500}$ .

(9) Circular measure of each angle is  $\frac{60 \times \pi}{180} = \frac{\pi}{3}$ .

(10) The angles are  $90^\circ$ ,  $45^\circ$ ,  $45^\circ$ , and of these the circular measures are  $\frac{\pi}{2}$ ,  $\frac{\pi}{4}$ ,  $\frac{\pi}{4}$ .

## EXAMPLES—IX. (p. 22).

(1) Measure in degrees is  $\frac{\pi \times 180}{2 \times \pi} = 90$ .

(2) Measure in degrees is  $\frac{\pi \times 180}{3 \times \pi} = 60$ .

(3) Measure in degrees is  $\frac{\pi \times 180}{4 \times \pi} = 45$ .

(4) Measure in degrees is  $\frac{\pi \times 180}{6 \times \pi} = 30.$

(5) Measure in degrees is  $\frac{2\pi \times 180}{3 \times \pi} = 120.$

(6) Measure in degrees is  $\frac{1 \times 180}{2 \times \pi} = \frac{90}{\pi}.$

(7) Measure in degrees is  $\frac{1 \times 180}{3 \times \pi} = \frac{60}{\pi}.$

(8) Measure in degrees is  $\frac{1 \times 180}{4 \times \pi} = \frac{45}{\pi}.$

(9) Measure in degrees is  $\frac{1 \times 180}{6 \times \pi} = \frac{30}{\pi}.$

(10) Measure in degrees is  $\frac{2 \times 180}{3 \times \pi} = \frac{120}{\pi}.$

EXAMPLES—X. (p. 22).

(1) Circular measure is  $\frac{50 \times \pi}{200} = \frac{\pi}{4}.$

(2) Circular measure is  $\frac{25 \times \pi}{200} = \frac{\pi}{8}.$

(3) Circular measure is  $\frac{6 \cdot 25 \times \pi}{200} = \frac{\pi}{32}.$

(4) Circular measure is  $\frac{250 \times \pi}{200} = \frac{5\pi}{4}.$

(5) Circular measure is  $\frac{500 \times \pi}{200} = \frac{5\pi}{2}.$

(6) Circular measure is  $\frac{13 \cdot 0505 \times \pi}{200} = \cdot 0652525\pi.$

(7) Circular measure is  $\frac{24 \cdot 150215 \times \pi}{200} = \cdot 120751075\pi.$

(8) Circular measure is  $\frac{125 \cdot 0013 \times \pi}{200} = \cdot 6250065\pi.$

$$(9) \text{ Circular measure is } \frac{.03 \times \pi}{200} = .00015\pi.$$

$$(10) \text{ Circular measure is } \frac{.0005 \times \pi}{200} = .0000025\pi.$$

EXAMPLES—XI. (p. 22).

$$(1) \text{ Measure in grades is } \frac{\pi \times 200}{3 \times \pi} = 66\frac{2}{3}.$$

$$(2) \text{ Measure in grades is } \frac{\pi \times 200}{5 \times \pi} = 40.$$

$$(3) \text{ Measure in grades is } \frac{\pi \times 200}{6 \times \pi} = 33\frac{1}{3}.$$

$$(4) \text{ Measure in grades is } \frac{2\pi \times 200}{3 \times \pi} = 133\frac{1}{3}.$$

$$(5) \text{ Measure in grades is } \frac{3\pi \times 200}{5 \times \pi} = 120.$$

$$(6) \text{ Measure in grades is } \frac{1 \times 200}{3 \times \pi} = \frac{200}{3\pi}.$$

$$(7) \text{ Measure in grades is } \frac{1 \times 200}{5 \times \pi} = \frac{40}{\pi}.$$

$$(8) \text{ Measure in grades is } \frac{1 \times 200}{8 \times \pi} = \frac{25}{\pi}.$$

$$(9) \text{ Measure in grades is } \frac{3 \times 200}{5 \times \pi} = \frac{120}{\pi}.$$

$$(10) \text{ Measure in grades is } \frac{23 \times 200}{10 \times \pi} = \frac{460}{\pi}.$$

EXAMPLES—XII. (p. 23).

$$(1) \text{ Measure} = 22\frac{1}{2} \div 5 = 22.5 \div 5 = 4.5.$$

$$(2) \text{ Unit} = 42.5^\circ \div 10 = 4.25^\circ.$$

$$(3) \text{ Angle} = 8 \times 2^\circ, \text{ or, } 16^\circ;$$

$$\therefore \text{ larger unit} = 16^\circ \div 5 = 3\frac{1}{5}^\circ.$$

Then, smaller unit in terms of larger is  $2 \div 3\frac{1}{5}$ , or,  $\frac{5}{8}$ ,

and larger unit in terms of smaller is  $3\frac{1}{5} \div 2$ , or,  $\frac{8}{5}$ .

(4) Angle =  $7 \times 3^\circ$ , or,  $21^\circ$ ;

$\therefore$  larger unit =  $21^\circ \div 6 = 3\frac{1}{2}^\circ$ .

Then, smaller unit in terms of larger is  $3 \div 3\frac{1}{2}$ , or,  $\frac{2}{5}$ ,  
and larger unit in terms of smaller is  $3\frac{1}{2} \div 3$ , or,  $\frac{7}{6}$ .

(5) Measure =  $42 \div 45 = 1\frac{4}{5}$ .

(6)  $13^\circ.13'.48'' = 47628''$ ,

$14^\circ.7' = \frac{227934''}{5}$ ;

$\therefore$  ratio =  $47628 \times 5 : 227934 = 70 : 67$ .

(7)  $G : D = 10 : 9$ ,  $\therefore$ ,  $9G = 10D$ ,  $\therefore$ ,  $G = D + \frac{1}{9}D$ .

(8) The angles of each triangle are  $90^\circ$ ,  $60^\circ$ ,  $30^\circ$ , because the line, drawn from any angle of an equilateral triangle to bisect the base, cuts the base at right angles, and bisects the vertical angle.

Expressed in grades the angles are  $100^\circ$ ,  $66\frac{2}{3}^\circ$ ,  $33\frac{1}{3}^\circ$ .

(9) Let  $x + y$ ,  $x$ ,  $x - y$  be the angles.

Then  $x + y + x + x - y = 180^\circ$ , or,  $3x = 180^\circ$ , or,  $x = 60^\circ$ .

(10)	$39^\circ.012$
	$\phantom{39^\circ.}9$
	$10 \overline{) 351.108}$
degrees	$35.1108$
	$\phantom{35.}60$
minutes	$6.6480$
	$\phantom{6.}60$
seconds	$38.8800$

(11) Number of degrees in the angle =  $\frac{m}{60}$ .

Number of grades                   ,,   =  $\frac{m \times 10}{60 \times 9}$

Number of French seconds,,   =  $\frac{m \times 10 \times 100 \times 100}{60 \times 9} = \frac{5000m}{27}$ .

(12)  $5^\circ.33'.20'' = 20000''$ ; and  $90^\circ = 324000''$ ;

$\therefore$  fraction =  $\frac{20000}{5} \div 324000 = \frac{4000}{324000} = \frac{1}{81}$ .



- (13) Let
- $x$
- be the measure of the angle in degrees.

Then  $\frac{10x}{9}$  is the measure of the angle in grades,

$$\text{and } \frac{1}{x} + \frac{9}{10x} = 1, \text{ or, } 10x = 19, \text{ or, } x = 1.9;$$

 $\therefore$  unit angle is  $1.9^\circ$ .

- (14) Let
- $x+y$
- ,
- $x$
- ,
- $x-y$
- be the angles expressed in degrees.

$$\text{Then } \frac{10(x+y)}{9} = x + (x-y);$$

$$\text{or, } 10x + 10y = 18x - 9y, \text{ and } \therefore x = \frac{19y}{8};$$

$$\therefore \text{ the angles are } \frac{27y}{8}, \frac{19y}{8}, \frac{11y}{8},$$

and these are in the ratio  $27 : 19 : 11$ .

- (15)
- $\frac{180^\circ}{\sqrt{3}} = \frac{10 \times 180^\circ}{9 \times \sqrt{3}} = \frac{200^\circ}{\sqrt{3}} = \frac{200\sqrt{3}^\circ}{3} = 115.47^\circ$
- nearly.

- (16) Let
- $x+y$
- ,
- $x$
- ,
- $x-y$
- be the angles expressed in degrees.
- 
- Then
- $x+y+x+x-y=180^\circ$
- , or,
- $3x=180^\circ$
- , or,
- $x=60^\circ$
- .

$$\text{Also } \frac{10}{9}(60-y) : 60+y = 2 : 9;$$

$$\text{or, } 600 - 10y = 120 + 2y, \text{ and } \therefore y = 40^\circ.$$

Hence the angles are  $100^\circ$ ,  $60^\circ$ ,  $20^\circ$ .

- (17) Circumference : diameter
- $= 360 : 2 \times 57.29577$
- 
- $= 180 : 57.29577$
- 
- $= 3.14159 \dots : 1.$

- (18) The sum of the two angles is
- $90^\circ$
- , because the third angle is
- $90^\circ$
- .
- 
- Hence, dividing
- $90^\circ$
- into two parts proportional to 2 and 3,
- 
- we have
- $36^\circ$
- and
- $54^\circ$
- for the angles.

$$\therefore \text{ angles expressed in degrees are } 90^\circ, 54^\circ, 36^\circ.$$

$$\text{,, ,, grades are } 100^\circ, 60^\circ, 40^\circ.$$

$$\text{,, ,, circular measure are } \frac{\pi}{2}, \frac{3\pi}{10}, \frac{\pi}{5}.$$

- (19) Angle :
- $360^\circ = 13 : 27$
- ;

$$\therefore \text{ angle} = \frac{360 \times 13}{27} \text{ degrees} = \frac{40 \times 13}{3} \text{ degrees} = 173\frac{1}{3}^\circ.$$

(20) Angle :  $400^s = 17 : 54$  ;

$$\therefore \text{angle} = \frac{400 \times 17}{54} \text{ grades} = 125.925 \text{ grades.}$$

(21) Angle subtended by an arc 18 inches long = unit of circular

$$\text{measure} = \frac{200}{\pi} \text{ grades ;}$$

$$\therefore \text{angle subtended by an arc 24 inches long} = \frac{24 \times 200}{18 \times \pi} \text{ gr.} = \frac{800}{3\pi} \text{ gr.}$$

(22) 1st angle contains  $\frac{2 \times 200}{\pi}$  grades, or,  $\frac{400}{\pi}$  grades.

2d angle contains  $\frac{10 \times 20}{9}$  grades, or,  $\frac{200}{9}$  grades.

3d angle contains  $\left(200 - \frac{400}{\pi} - \frac{200}{9}\right) \text{ gr.}$ , or,  $\frac{1600\pi - 3600}{9\pi} \text{ gr.}$

(23) Angle required =  $\frac{7}{2}$  of  $15^\circ.39'.7'' = 54^\circ.46'.54''.5$ .

(24) Circular measure =  $\frac{11.3 \times \pi}{200} = \frac{113 \times 355}{2000 \times 113} = \frac{71}{400} = .1775$ .

(25) Measure in degrees =  $\frac{180 \times \pi^2}{\pi \times 9} = 20\pi$ .

(26) Larger circumference = 400 times smaller circumference.

Then, since  $\frac{1}{360}$ th part of smaller circumference subtends an angle of  $1^\circ$  at the centre, it follows that  $\frac{1}{400}$  of  $\frac{1}{360}$ th part of the larger circumference will subtend the same angle.

$$\therefore \text{part required} = \frac{1}{400 \times 360} = \frac{1}{144000}.$$

(27) 4 right angles =  $360^\circ = 400^s = 2\pi^c$  ;

$$\therefore \text{the measure of } 1^\circ \text{ will be } \frac{1}{360},$$

$$\text{the measure of } 1^s \text{ will be } \frac{1}{400},$$

$$\text{the measure of } 1^c \text{ will be } \frac{1}{2\pi}.$$

(28) Length of whole circumference of earth =  $7980\pi$  miles ;

$$\therefore \text{length of 1 degree of meridian} = \frac{7980\pi}{360} \text{ miles} = \frac{133\pi}{6} \text{ miles.}$$

(29) (1)  $\frac{3}{2} \times 45^\circ = 67\frac{1}{2}^\circ$  ;  $4 \times 45^\circ = 180^\circ$  ;  $\pi \times 45^\circ = 45\pi^\circ$  ;

$$\left(4n + \frac{1}{3}\right) \times 45^\circ = (n \cdot 180 + 15)^\circ.$$

$$(2) \frac{3}{2} \times \frac{\pi}{4} = \frac{3\pi}{8} ; 4 \times \frac{\pi}{4} = \pi ; \pi \times \frac{\pi}{4} = \left(\frac{\pi}{2}\right)^2 ;$$

$$\left(4n + \frac{1}{3}\right) \times \frac{\pi}{4} = n\pi + \frac{\pi}{12}.$$

(30) Number of degrees in the unit angle =  $\frac{3 \times 180}{\pi}$  ;

$$\therefore \text{measure of an angle of } 45^\circ = 45 \div \frac{3 \times 180}{\pi} = \frac{45 \times \pi}{3 \times 180} = \frac{\pi}{12}.$$

(31) (1) Sum of angles =  $(12 - 4)$  right angles = 8 right angles.

(EUCLID, I. XXXII., COR. 1.)

$$\therefore \text{each angle} = \frac{8 \times 90}{6} \text{ degrees} = 120^\circ.$$

(2) Sum of angles =  $(10 - 4)$  right angles = 6 right angles ;

$$\therefore \text{each angle} = \frac{6 \times 90}{5} \text{ degrees} = 108^\circ.$$

(32) (1) Each angle =  $\frac{6 \times 100}{5}$  grades =  $120^\circ$ .

(2) Each angle =  $\frac{12 \times 100}{8}$  grades =  $150^\circ$ .

(33) (1) Circular measure of each angle =  $\frac{\pi}{3}$ .

$$(2) \text{Circular measure of each angle} = \frac{8 \times \pi}{6 \times 2} = \frac{2\pi}{3}.$$

(34) Sum of all the angles =  $(2n - 4)$  right angles ;

$$\therefore \text{circular measure of each angle} = \frac{2n - 4}{n} \cdot \frac{\pi}{2} = \pi - \frac{2\pi}{n}.$$

(35) Arc subtending an angle of  $180^\circ = 18\pi$  feet.

$$\therefore \text{arc subtending an angle of } 10^\circ = \frac{18\pi}{18} \text{ feet} = \pi \text{ feet.}$$



(36) Let  $2n$  and  $n$  be the number of sides in the polygons, respectively.

Each angle in first polygon contains  $\frac{4n-4}{2n}$  right angles.

Each angle in second polygon contains  $\frac{2n-4}{n}$  right angles.

$$\therefore \frac{4n-4}{2n} : \frac{2n-4}{n} = 3 : 2 ;$$

$$\therefore 4n-4 = 6n-12, \text{ or, } 2n=8, \text{ or, } n=4.$$

Hence the number of sides will be 8 and 4 respectively.

#### EXAMPLES—XIII. (p. 36).

$$\begin{aligned} (1) \quad \sin BAD &= \frac{BD}{AB}; \quad \cos BAD = \frac{AD}{AB}; \quad \tan BAD = \frac{BD}{AD}; \\ \sin ABD &= \frac{AD}{AB}; \quad \cot ABD = \frac{BD}{AD}; \quad \operatorname{cosec} ABD = \frac{AB}{AD}; \\ \sin BCD &= \frac{BD}{BC}; \quad \sin CBD = \frac{CD}{BC}; \quad \tan BCD = \frac{BD}{CD}. \end{aligned}$$




FIG. 6.

$$(2) \quad \frac{a}{b} = \sin A, \therefore a = b \cdot \sin A,$$

$$\frac{a}{b} = \cos C, \therefore a = b \cdot \cos C,$$

$$\frac{a}{c} = \tan A, \therefore a = c \cdot \tan A,$$

$$\frac{a}{c} = \cot C, \therefore a = c \cdot \cot C;$$

and similarly for the rest of the Examples.

EXAMPLES—XIV. (p. 49).

$$(1) \cos \alpha \cdot \sin \gamma \cdot \cos \delta = \cos 0^\circ \cdot \sin 45^\circ \cdot \cos 60^\circ = 1 \times \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{1}{2\sqrt{2}}.$$

$$(2) \sin \theta \cdot \cos \frac{\pi}{4} \cdot \operatorname{cosec} \delta = \sin 90^\circ \cdot \cos 45^\circ \cdot \operatorname{cosec} 60^\circ \\ = 1 \times \frac{1}{\sqrt{2}} \times \frac{2}{\sqrt{3}} = \sqrt{\frac{2}{3}}.$$

$$(3) \sin \frac{\pi}{2} + \cos \frac{\pi}{6} \cdot \sec \alpha = \sin 90^\circ + \cos 30^\circ - \sec 0^\circ = 1 + \frac{\sqrt{3}}{2} - 1 = \frac{\sqrt{3}}{2}.$$

$$(4) \sin \frac{\pi}{3} \cdot \operatorname{cosec} \frac{\pi}{2} \cdot \sec \delta = \sin 60^\circ \cdot \operatorname{cosec} 90^\circ \cdot \sec 60^\circ = \frac{\sqrt{3}}{2} \times 1 \times 2 = \sqrt{3}.$$

$$(5) (\sin \theta - \cos \theta + \operatorname{cosec} \beta) \left( \cos \theta + \sec \frac{\pi}{4} + \cot \delta \right) \\ = (\sin 90^\circ - \cos 90^\circ + \operatorname{cosec} 30^\circ) \cdot (\cos 90^\circ + \sec 45^\circ + \cot 60^\circ) \\ = (1 - 0 + 2) \cdot \left( 0 + \sqrt{2} + \frac{1}{\sqrt{3}} \right) = 3 \times \left( \sqrt{2} + \frac{1}{\sqrt{3}} \right) = 3\sqrt{2} + \sqrt{3}.$$

$$(6) (\sin \delta - \sin \gamma) (\cos \beta + \cos \gamma) = (\sin 60^\circ - \sin 45^\circ) (\cos 30^\circ + \cos 45^\circ) \\ = \left( \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \right) \left( \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \right) = \frac{3}{4} - \frac{1}{2} = \frac{1}{4}.$$

$$\sin^2 \beta = \sin^2 30^\circ = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}.$$

$$(7) \cot^2 \frac{\pi}{4} - \cot^2 \frac{\pi}{6} = \cot^2 45^\circ - \cot^2 30^\circ = 1 - 3 = -2.$$

$$\frac{\sin^2 \frac{\pi}{6} - \sin^2 \frac{\pi}{4}}{\sin^2 \frac{\pi}{4} \cdot \sin^2 \frac{\pi}{6}} = \frac{\frac{1}{4} - \frac{1}{2}}{\frac{1}{4} \cdot \frac{1}{2}} = \frac{-2}{1} = -2.$$

$$(8) \left( \sin \frac{\pi}{6} + \cos \frac{\pi}{6} \right) \left( \sin \frac{\pi}{3} - \cos \frac{\pi}{3} \right) = \left( \frac{1}{2} + \frac{\sqrt{3}}{2} \right) \left( \frac{\sqrt{3}}{2} - \frac{1}{2} \right) \\ = \frac{3}{4} - \frac{1}{4} = \frac{1}{2} = \cos \frac{\pi}{3}.$$

$$(9) \cos \frac{\pi}{3} \cdot \cos \frac{\pi}{6} = \frac{1}{2} \cdot \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{4}.$$

$$\frac{1}{2} \cos \left( \frac{\pi}{3} + \frac{\pi}{6} \right) + \frac{1}{2} \cos \left( \frac{\pi}{3} - \frac{\pi}{6} \right) = \frac{1}{2} \cos \frac{\pi}{2} + \frac{1}{2} \cos \frac{\pi}{6} \\ = \frac{1}{2} \times 0 + \frac{1}{2} \times \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{4}.$$

$$(10) \tan^2 \frac{\pi}{3} - \tan^2 \frac{\pi}{6} = 3 - \frac{1}{3} = \frac{8}{3}.$$

$$\frac{\sin^2 \frac{\pi}{3} - \sin^2 \frac{\pi}{6}}{\cos^2 \frac{\pi}{3} \cdot \cos^2 \frac{\pi}{6}} = \frac{\frac{3}{4} - \frac{1}{4}}{\frac{1}{4} \cdot \frac{3}{4}} = \frac{8}{3}.$$

EXAMPLES—XV. (p. 52).




FIG. 7.

(1.) Let  $PM$  be the tower;  $A$  the place of observation.

Then  $AM = 200$  feet, and  $\angle PAM = 60^\circ$ .

Now  $PM = AM \cdot \tan PAM$

$= (200 \times \sqrt{3})$  feet  $= 346.4101 \dots$  feet.




FIG. 8.

(2) Let  $RO$  be the tower;  $P$  the point of observation.

Then  $AP = 140$  feet, and  $\angle RPA = 30^\circ$ .

Now  $RA = PA \cdot \tan 30^\circ$ .

$= \frac{140}{\sqrt{3}}$  feet  $= \frac{140\sqrt{3}}{3}$  feet  $= 80.829037 \dots$  feet.

$\therefore RO = 80.829037 \dots$  feet  $+ 5$  feet  $= 85.829037 \dots$  feet.

(3) Taking the diagram in Art. 87,

$$AB:BQ=\sqrt{3}:1;$$

$$\therefore \tan SQR=\sqrt{3}, \text{ and } \therefore \angle SQR=60^\circ.$$

(4) Let  $AB$  be the steeple;  $P$  the point of observation.

$$\text{Then } PB=300 \text{ feet, and } \angle APB=30^\circ.$$

$$\text{Then } AB=PB \cdot \tan \angle APB$$

$$\begin{aligned} &= 300 \cdot \frac{1}{\sqrt{3}} \text{ feet} = 100\sqrt{3} \cdot \text{feet} \\ &= 173 \cdot 205 \dots \text{feet.} \end{aligned}$$




FIG. 9.

(5) Let  $AP$  be the rock;  $O$  the position of the ship.

$$\text{Then } AP=245 \text{ feet; and } \angle AOP=30^\circ.$$

$$\text{Now } PO=AP \cdot \cot \angle AOP$$

$$= 245 \cdot \sqrt{3} \text{ ft.} = 424 \cdot 352 \dots \text{ft.}$$




FIG. 10.

(6) Let  $AB$  be the hill;  $C$  and  $D$  the positions of the milestones.

$$\text{Then } DC=1 \text{ mile; } \angle ACB=45^\circ; \quad \cdot$$

$$\angle ADB=30^\circ.$$

$$\text{Hence } \angle CAB=45^\circ, \text{ and } AB=BC.$$

$$\text{Let } x=\text{height of hill in miles.}$$

$$\text{Then } AB=BD \cdot \tan \angle ADB$$

$$= (BC + CD) \cdot \tan 30^\circ;$$

$$\therefore x = (x+1) \cdot \frac{1}{\sqrt{3}};$$

$$\therefore \sqrt{3} \cdot x = x+1, \text{ or, } x = \frac{1}{\sqrt{3}-1} = \frac{\sqrt{3}+1}{3-1} = \frac{\sqrt{3}+1}{2};$$

$$\therefore x = \frac{2 \cdot 732}{2} \text{ miles} = 1 \cdot 366 \dots \text{miles.}$$




FIG. 11.

(7) Let  $AO$  be the flag-staff;  $PO$  the tower;  $M$  the point of observation.




FIG. 12.

Then  $PM = 100$  feet;  $\angle AMP = 45^\circ$ ;  
 $\angle OMP = 30^\circ$ .

$$\begin{aligned}\text{Then } AO &= AP - OP \\ &= PM - PM \cdot \tan OMP \\ &= \left(100 - 100 \cdot \frac{1}{\sqrt{3}}\right) \text{ feet.} \\ &= \frac{300 - 100\sqrt{3}}{3} \text{ feet} \\ &= \frac{300 - 173.205 \dots}{3} \text{ feet} = 42.265 \dots \text{ feet.}\end{aligned}$$

(8) Let  $AP$  be the tower;  $MO$  the column;  $MD$  parallel to  $OP$ .




FIG. 13.

Then  $\angle AMD = 30^\circ$ , and  $\angle AOP = 60^\circ$ .

$$\text{Then } MD = OP = AP \cdot \cot AOP = \frac{108}{\sqrt{3}} \text{ feet.}$$

$$\begin{aligned}\text{And } AD &= MD \cdot \tan AMD = \frac{108}{\sqrt{3}} \cdot \frac{1}{\sqrt{3}} \text{ feet} \\ &= \frac{108}{3} \text{ feet} = 36 \text{ feet.}\end{aligned}$$

$$\text{Hence } OM = AP - AD = (108 - 36) \text{ feet} = 72 \text{ feet.}$$

(9.) Let  $AP$  be the tower;  $M$  and  $O$  the points of observation.




FIG. 14.

Then  $\angle AMP = 60^\circ$ ; and  $\angle AOP = 30^\circ$ .

Let  $x$  = height of tower in yards.

Now  $\angle AMP = \angle AOP + \angle OAM$ ,

$$\text{or, } 60^\circ = 30^\circ + \angle OAM;$$

$$\therefore \angle OAM = 30^\circ = \angle AOM,$$

$$\text{and } \therefore MA = OM = 100 \text{ yards.}$$

$$\text{Then } AP = MA \cdot \sin AMP = 100 \cdot \frac{\sqrt{3}}{2} \text{ yards} = 50\sqrt{3} \text{ yards.}$$

(10) Taking the diagram in Art. 87.

$$\tan AQB = \frac{AB}{BQ} = \frac{10}{25} = .4;$$

$\therefore$  altitude of the sun is  $25^\circ$ .

(11) The diagram represents a vertical section of the spire and tower.

Let  $x$  represent the height of the spire in feet.

$$\text{Then } AM = x + 35 - 23 = x + 12,$$

$$BM = 60 + 17\frac{1}{2} = 77.5,$$

$$\text{and } \frac{x+12}{77.5} = \tan ABM = 1.5;$$

$$\therefore x + 12 = 116.25, \text{ or, } x = 104.25 \text{ feet.}$$




FIG. 15.

(12) Let  $OB$  be the height of the kite in yards.

Let  $AO$  be the string.

$$\text{Then } OB = AO \cdot \sin OAB$$

$$= \left(250 \times \frac{1}{2}\right) \text{ yards} = 125 \text{ yards.}$$




FIG. 16.

(13) Let  $AC$  be the rope;  $AB$  the height of the house.

$$\text{Then } \angle ACB = 40^\circ. 30'.$$

$$\text{And } AC = \frac{AB}{\sin ACB} = \frac{60}{.65} \text{ feet} = 92\frac{4}{13} \text{ feet.}$$




FIG. 17.

(14) Let  $AC$  be the tower;  $BC$  the breadth of the river.




FIG. 18.

Then  $\angle ABC = 20^\circ$ .

$$\begin{aligned} \text{And } BC &= \frac{AC}{\tan \angle ABC} \\ &= \frac{120}{.35} \text{ feet} = 342\frac{2}{5} \text{ feet.} \end{aligned}$$

(15) Taking the diagram of Art. 87.

$$\text{Length of shadow} = QB = \frac{AB}{\tan \angle QB} = \frac{6}{.745} \text{ feet} = 8.053 \dots \text{ feet.}$$

#### EXAMPLES—XVI. (p. 57).

$$(1) \cos \theta \cdot \tan \theta = \cos \theta \cdot \frac{\sin \theta}{\cos \theta} = \sin \theta.$$

$$(2) \sin \theta \cdot \cot \theta = \sin \theta \cdot \frac{\cos \theta}{\sin \theta} = \cos \theta.$$

$$(3) \sin a \cdot \sec a = \sin a \cdot \frac{1}{\cos a} = \frac{\sin a}{\cos a} = \tan a.$$

$$(4) \cos a \cdot \operatorname{cosec} a = \cos a \cdot \frac{1}{\sin a} = \frac{\cos a}{\sin a} = \cot a.$$

$$(5) (1 + \tan^2 \theta) \cdot \cos^2 \theta = \sec^2 \theta \cdot \cos^2 \theta = \frac{\cos^2 \theta}{\cos^2 \theta} = 1.$$

$$(6) (1 + \cot^2 \theta) \cdot \sin^2 \theta = \operatorname{cosec}^2 \theta \cdot \sin^2 \theta = \frac{\sin^2 \theta}{\sin^2 \theta} = 1.$$

$$(7) \frac{\tan^2 a}{1 + \tan^2 a} = \frac{\tan^2 a}{\sec^2 a} = \frac{\sin^2 a}{\cos^2 a} \cdot \cos^2 a = \sin^2 a.$$

$$(8) \frac{\operatorname{cosec}^2 a - 1}{\operatorname{cosec}^2 a} = 1 - \frac{1}{\operatorname{cosec}^2 a} = 1 - \sin^2 a = \cos^2 a.$$

$$(9) \tan x + \cot x = \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{\sin^2 x + \cos^2 x}{\cos x \cdot \sin x} = \frac{1}{\cos x \cdot \sin x} = \sec x \cdot \operatorname{cosec} x.$$

- $$(10) \frac{\cos x \cdot \operatorname{cosec} x \cdot \tan x}{\sin x \cdot \sec x \cdot \cot x} = \frac{\cos x \cdot \frac{1}{\sin x} \cdot \frac{\sin x}{\cos x}}{\sin x \cdot \frac{1}{\cos x} \cdot \frac{\cos x}{\sin x}} = \frac{\cos x \cdot \sin x \cdot \cos x \cdot \sin x}{\sin x \cdot \sin x \cdot \cos x \cdot \cos x} = 1.$$
- $$(11) \cos x + \sin x \cdot \tan x = \cos x + \frac{\sin^2 x}{\cos x} = \frac{\cos^2 x + \sin^2 x}{\cos x} = \frac{1}{\cos x} = \sec x.$$
- $$(12) \frac{\cos \theta}{\tan \theta \cdot \cot^2 \theta} = \frac{\cos \theta}{\cot \theta} = \frac{\cos \theta \cdot \sin \theta}{\cos \theta} = \sin \theta.$$
- $$(13) (\cos^2 \theta - 1)(\cot^2 \theta + 1) = (\cos^2 \theta - 1) \cdot \operatorname{cosec}^2 \theta = -\sin^2 \theta \times \frac{1}{\sin^2 \theta} = -1.$$
- $$(14) \cot^2 a - \cos^2 a = \frac{\cos^2 a}{\sin^2 a} - \cos^2 a = \cos^2 a \left( \frac{1}{\sin^2 a} - 1 \right) = \cos^2 a \cdot \frac{1 - \sin^2 a}{\sin^2 a} \\ = \cos^2 a \cdot \frac{\cos^2 a}{\sin^2 a} = \cot^2 a \cdot \cos^2 a.$$
- $$(15) \sec^2 a \cdot \operatorname{cosec}^2 a = \sec^2 a (1 + \cot^2 a) = \sec^2 a + \sec^2 a \cdot \frac{\cos^2 a}{\sin^2 a} \\ = \sec^2 a + \operatorname{cosec}^2 a.$$
- $$(16) \sin^2 \phi + \sin^2 \phi \cdot \tan^2 \phi = \sin^2 \phi (1 + \tan^2 \phi) = \sin^2 \phi \cdot \sec^2 \phi = \tan^2 \phi.$$
- $$(17) \cot^2 \phi \cdot \sin^2 \phi + \sin^2 \phi = \sin^2 \phi (\cot^2 \phi + 1) = \sin^2 \phi \cdot \operatorname{cosec}^2 \phi = 1.$$
- $$(18) \sec^2 \phi - 1 = \tan^2 \phi = \frac{\sin^2 \phi}{\cos^2 \phi} = \sin^2 \phi \cdot \sec^2 \phi.$$
- $$(19) 2 \operatorname{versin} \phi - \operatorname{versin}^2 \phi = 2(1 - \cos \phi) - (1 - \cos \phi)^2 \\ = 2 - 2 \cos \phi - 1 + 2 \cos \phi - \cos^2 \phi = 1 - \cos^2 \phi = \sin^2 \phi.$$
- $$(20) \frac{\sec \theta - 1}{\sec \theta} = 1 - \frac{1}{\sec \theta} = 1 - \cos \theta = \operatorname{versin} \theta.$$

## EXAMPLES—XVII. (p. 60).

- (1) Let  $PAM$  be an angle whose cosine is  $c$ .

Draw  $PM$  perpendicular to  $AM$ .

Then if  $AP$  be represented by 1,

$AM$  will be represented by  $c$ , and

$PM$  will be represented by  $\sqrt{1-c^2}$ .

Then, denoting  $\angle PAM$  by  $A$ ,

$$\sin A = \frac{PM}{AP} = \frac{\sqrt{1-c^2}}{1} = \sqrt{1-\cos^2 A}$$




FIG. 19.



$$\begin{aligned}\tan A &= \frac{PM}{AM} = \frac{\sqrt{1-c^2}}{c} = \frac{\sqrt{1-\cos^2 A}}{\cos A} \\ \sec A &= \frac{AP}{AM} = \frac{1}{c} = \frac{1}{\cos A} \\ \operatorname{cosec} A &= \frac{AP}{PM} = \frac{1}{\sqrt{1-c^2}} = \frac{1}{\sqrt{1-\cos^2 A}} \\ \cot A &= \frac{AM}{PM} = \frac{c}{\sqrt{1-c^2}} = \frac{\cos A}{\sqrt{1-\cos^2 A}}\end{aligned}$$

(2) Let  $PAM$  be an angle whose cosecant is  $c$ .

Constructing a diagram as in Example (1), the measures of  $AP$ ,  $PM$ ,  $AM$  may be taken as  $c$ ,  $1$ ,  $\sqrt{c^2-1}$  respectively.

$$\begin{aligned}\text{Then } \sin A &= \frac{PM}{AP} = \frac{1}{c} = \frac{1}{\operatorname{cosec} A} \\ \cos A &= \frac{AM}{AP} = \frac{\sqrt{c^2-1}}{c} = \frac{\sqrt{\operatorname{cosec}^2 A - 1}}{\operatorname{cosec} A} \\ \tan A &= \frac{PM}{AM} = \frac{1}{\sqrt{c^2-1}} = \frac{1}{\sqrt{\operatorname{cosec}^2 A - 1}} \\ \sec A &= \frac{AP}{AM} = \frac{c}{\sqrt{c^2-1}} = \frac{\operatorname{cosec} A}{\sqrt{\operatorname{cosec}^2 A - 1}} \\ \cot A &= \frac{AM}{MP} = \frac{\sqrt{c^2-1}}{1} = \sqrt{\operatorname{cosec}^2 A - 1}.\end{aligned}$$

(3) Let  $PAM$  be an angle whose secant is  $s$ .

Constructing a diagram as in Example (1), the measures of  $AP$ ,  $AM$ ,  $PM$ , may be taken as  $s$ ,  $1$ ,  $\sqrt{s^2-1}$  respectively.

$$\begin{aligned}\text{Then } \sin A &= \frac{PM}{AP} = \frac{\sqrt{s^2-1}}{s} = \frac{\sqrt{\sec^2 A - 1}}{\sec A} \\ \cos A &= \frac{AM}{AP} = \frac{1}{s} = \frac{1}{\sec A} \\ \tan A &= \frac{PM}{AM} = \frac{\sqrt{s^2-1}}{1} = \sqrt{\sec^2 A - 1} \\ \operatorname{cosec} A &= \frac{AP}{PM} = \frac{s}{\sqrt{s^2-1}} = \frac{\sec A}{\sqrt{\sec^2 A - 1}} \\ \cot A &= \frac{AM}{PM} = \frac{1}{\sqrt{s^2-1}} = \frac{1}{\sqrt{\sec^2 A - 1}}.\end{aligned}$$

(4) Let  $PAM$  be an angle whose cotangent is  $c$ .

Constructing a diagram as in Example (1), the measures of  $AM$ ,  $PM$ ,  $AP$  may be taken as  $c$ ,  $1$ ,  $\sqrt{1+c^2}$  respectively.

$$\text{Then } \sin A = \frac{PM}{AP} = \frac{1}{\sqrt{1+c^2}} = \frac{1}{\sqrt{1+\cot^2 A}}$$

$$\cos A = \frac{AM}{AP} = \frac{c}{\sqrt{1+c^2}} = \frac{\cot A}{\sqrt{1+\cot^2 A}}$$

$$\tan A = \frac{PM}{AM} = \frac{1}{c} = \frac{1}{\cot A}$$

$$\operatorname{cosec} A = \frac{AP}{PM} = \frac{\sqrt{1+c^2}}{1} = \sqrt{1+\cot^2 A}$$

$$\sec A = \frac{AP}{AM} = \frac{\sqrt{1+c^2}}{c} = \frac{\sqrt{1+\cot^2 A}}{\cot A}.$$

#### EXAMPLES—XVIII. (p. 61).

(1) Take the diagram as before; then if  $\angle PAM$  be denoted by  $a$ , the measure of  $PM$  may be denoted by 2, the measure of  $AP$  by 3, and therefore the measure of  $AM$  by  $\sqrt{9-4} = \sqrt{5}$ .

$$\text{Then } \cos a = \frac{\sqrt{5}}{3} \text{ and } \tan a = \frac{2}{\sqrt{5}}.$$

(2) Let the measure of  $AM$  be 4, and that of  $AP$  be 5; then that of  $PM$  will be  $\sqrt{25-16}$ , or, 3.

$$\text{Then } \sin a = \frac{3}{5}, \text{ and } \tan a = \frac{3}{4}.$$

(3) Let the measure of  $AP$  be 4, and that of  $PM$  be 3; then that of  $AM$  will be  $\sqrt{16-9}$ , or,  $\sqrt{7}$ .

$$\text{Then } \cos \theta = \frac{\sqrt{7}}{4}, \text{ and } \tan \theta = \frac{3}{\sqrt{7}}.$$

(4) Let the measure of  $PM$  be 1, and that of  $AP$  be  $\sqrt{3}$ ; then that of  $AM$  will be  $\sqrt{3-1}$ , or,  $\sqrt{2}$ .

$$\text{Then } \cos \theta = \frac{\sqrt{2}}{\sqrt{3}}, \text{ and } \tan \theta = \frac{1}{\sqrt{2}}.$$

(5) Let the measure of  $PM$  be  $a^2$ , and that of  $AM$  be  $b^2$ ; then that of  $AP$  will be  $\sqrt{a^4 + b^4}$ .

$$\text{Then cosec} \theta = \frac{\sqrt{a^4 + b^4}}{a^2}, \text{ and sec} \theta = \frac{\sqrt{a^4 + b^4}}{b^2}.$$

(6) Let the measure of  $AM$  be  $a$ , and that of  $AP$  be  $b$ ; then that of  $PM$  will be  $\sqrt{b^2 - a^2}$ .

$$\text{Then tan} \theta = \frac{\sqrt{b^2 - a^2}}{a}, \text{ and cosec} \theta = \frac{b}{\sqrt{b^2 - a^2}}.$$

(7) Let the measure of  $PM$  be  $a$ , and that of  $AP$  be 1; then that of  $AM$  will be  $\sqrt{1 - a^2}$ .

$$\text{Then tan} \theta = \frac{a}{\sqrt{1 - a^2}}, \text{ and sec} \theta = \frac{1}{\sqrt{1 - a^2}}.$$

(8) Let the measure of  $AM$  be  $b$ , and that of  $AP$  be 1; then that of  $PM$  will be  $\sqrt{1 - b^2}$ .

$$\text{Then tan} \theta = \frac{\sqrt{1 - b^2}}{b}, \text{ and cosec} \theta = \frac{1}{\sqrt{1 - b^2}}.$$

(9) Let the measure of  $PM$  be 6, and that of  $AP$  be 10; then that of  $AM$  will be  $\sqrt{100 - 36}$ , or, 8.

$$\text{Then cos} \theta = \frac{8}{10} = \frac{4}{5}, \text{ and cot} \theta = \frac{8}{6} = \frac{4}{3}.$$

(10) Let the measure of  $AM$  be 5, and that of  $AP$  be 9; then that of  $PM$  will be  $\sqrt{81 - 25} = \sqrt{56} = 2\sqrt{14}$ .

$$\text{Then cot} \theta = \frac{5}{2\sqrt{14}}, \text{ and cosec} \theta = \frac{9}{2\sqrt{14}}.$$

(11) Let the measure of  $AP$  be 22, and that of  $PM$  be 9; then that of  $AM$  will be  $\sqrt{484 - 81} = \sqrt{403}$ .

$$\text{Then cos} \theta = \frac{\sqrt{403}}{22}, \text{ and cot} \theta = \frac{\sqrt{403}}{9}.$$

$$(12) \ 1.0\dot{8} = \frac{103-10}{90} = \frac{93}{90} = \frac{31}{30}.$$

Let the measure of  $AP$  be 31, and that of  $AM$  be 30; then that of  $PM$  will be  $\sqrt{961-900}$ , or,  $\sqrt{61}$ .

$$\text{Then } \sin\theta = \frac{\sqrt{61}}{31}, \text{ and } \tan\theta = \frac{\sqrt{61}}{30}.$$

(13) Let the measure of  $PM$  be 99, and that of  $AP$  be 101; then that of  $AM$  will be  $\sqrt{10201-9801}$ , or, 20.

$$\text{Then } \cos\phi = \frac{20}{101}, \text{ and } \cot\phi = \frac{20}{99}.$$

(14) Let the measure of  $AM$  be 20, and that of  $AP$  be 101; then that of  $PM$  will be  $\sqrt{10201-400}$ , or, 99.

$$\text{Then } \sin\phi = \frac{99}{101}, \text{ and } \tan\phi = \frac{99}{20}.$$

$$(15) \ \cos\theta = 1 - \text{versin}\theta = 1 - \frac{1}{13} = \frac{12}{13}.$$

Let the measure of  $AM$  be 12, and that of  $AP$  be 13; then that of  $PM$  will be  $\sqrt{169-144}$ , or, 5.

$$\text{Then } \sin\theta = \frac{5}{13}, \text{ and } \sec\theta = \frac{13}{12}.$$

#### EXAMPLES—XIX. (p. 63).

$$(1) \ \sin A = \frac{1}{\text{cosec} A} = \frac{1}{\sqrt{\text{cosec}^2 A}} = \frac{1}{\sqrt{(1 + \cot^2 A)}}.$$

$$(2) \ \cos A = \frac{1}{\sec A} = \frac{1}{\sqrt{\sec^2 A}} = \frac{1}{\sqrt{(1 + \tan^2 A)}}.$$

$$(3) \ \cos x = \frac{\cot x}{\text{cosec} x} = \frac{\cot x}{\sqrt{(\text{cosec}^2 x)}} = \frac{\cot x}{\sqrt{(1 + \cot^2 x)}}.$$

$$(4) \ \tan x \cdot \cos x = \sin x = \sqrt{(1 - \cos^2 x)}.$$

$$(5) \ \cos\phi = \frac{\cot\phi}{\text{cosec}\phi} = \frac{\sqrt{(\text{cosec}^2\phi - 1)}}{\text{cosec}\phi}.$$

$$(6) \tan \phi = \frac{\sin \phi}{\cos \phi} = \frac{\sqrt{1 - \cos^2 \phi}}{\cos \phi} = \sqrt{\left( \frac{1 - \cos^2 \phi}{\cos^2 \phi} \right)}.$$

$$(7) \sin^2 a = 1 - \cos^2 a = (1 + \cos a)(1 - \cos a) = (1 + \cos a) \cdot \text{versina}.$$

$$(8) \tan^2 a - \tan^2 \beta = \frac{\sin^2 a}{\cos^2 a} - \frac{\sin^2 \beta}{\cos^2 \beta} = \frac{\sin^2 a \cdot \cos^2 \beta - \cos^2 a \cdot \sin^2 \beta}{\cos^2 a \cdot \cos^2 \beta} \\ = \frac{(1 - \cos^2 a) \cos^2 \beta - (1 - \cos^2 \beta) \cos^2 a}{\cos^2 a \cdot \cos^2 \beta} = \frac{\cos^2 \beta - \cos^2 a}{\cos^2 a \cdot \cos^2 \beta}.$$

$$(9) \cot^2 a - \cot^2 \beta = \frac{\cos^2 a}{\sin^2 a} - \frac{\cos^2 \beta}{\sin^2 \beta} = \frac{\cos^2 a \cdot \sin^2 \beta - \cos^2 \beta \cdot \sin^2 a}{\sin^2 a \cdot \sin^2 \beta} \\ = \frac{(1 - \sin^2 a) \sin^2 \beta - (1 - \sin^2 \beta) \sin^2 a}{\sin^2 a \cdot \sin^2 \beta} = \frac{\sin^2 \beta - \sin^2 a}{\sin^2 a \cdot \sin^2 \beta}.$$

$$(10) \sin^2 \theta \cdot \tan^2 \theta + \cos^2 \theta \cdot \cot^2 \theta = (1 - \cos^2 \theta) \cdot \tan^2 \theta + (1 - \sin^2 \theta) \cdot \cot^2 \theta \\ = \tan^2 \theta - \sin^2 \theta + \cot^2 \theta - \cos^2 \theta = \tan^2 \theta + \cot^2 \theta - (\sin^2 \theta + \cos^2 \theta) \\ = \tan^2 \theta + \cot^2 \theta - 1.$$

$$(11) \sec^4 \theta + \tan^4 \theta = (1 + \tan^2 \theta)^2 + \tan^4 \theta = 1 + 2 \tan^2 \theta + \tan^4 \theta + \tan^4 \theta \\ = 1 + 2 \tan^2 \theta (1 + \tan^2 \theta) = 1 + 2 \tan^2 \theta \cdot \sec^2 \theta.$$

$$(12) \operatorname{cosec} \theta (\sec \theta - 1) - \cot \theta (1 - \cos \theta) = \frac{1}{\sin \theta \cdot \cos \theta} - \frac{1}{\sin \theta} - \frac{\cos \theta}{\sin \theta} + \frac{\cos^2 \theta}{\sin \theta} \\ = \frac{1 - \cos^2 \theta}{\sin \theta \cdot \cos \theta} - \frac{1 - \cos^2 \theta}{\sin \theta} = \frac{\sin^2 \theta}{\sin \theta \cdot \cos \theta} - \frac{\sin^2 \theta}{\sin \theta} = \tan \theta - \sin \theta.$$

$$(13) \cot^2 b + \tan^2 b = (\operatorname{cosec}^2 b - 1) + (\sec^2 b - 1) = \operatorname{cosec}^2 b + \sec^2 b - 2 \\ = \frac{1}{\sin^2 b} + \frac{1}{\cos^2 b} - 2 = \frac{\cos^2 b + \sin^2 b}{\sin^2 b \cdot \cos^2 b} - 2 \\ = \frac{1}{\sin^2 b \cdot \cos^2 b} - 2 = \operatorname{cosec}^2 b \cdot \sec^2 b - 2.$$

$$(14) \cot^2 A - \cos^2 A = \frac{\cos^2 A}{\sin^2 A} - \cos^2 A = \cos^2 A \left( \frac{1}{\sin^2 A} - 1 \right) \\ = \cos^2 A \left( \frac{1 - \sin^2 A}{\sin^2 A} \right) = \cos^2 A \cdot \frac{\cos^2 A}{\sin^2 A} = \cos^4 A \cdot \operatorname{cosec}^2 A.$$

$$(15) \tan^2 \theta - \sin^2 \theta = \frac{\sin^2 \theta}{\cos^2 \theta} - \sin^2 \theta = \sin^2 \theta \left( \frac{1}{\cos^2 \theta} - 1 \right) \\ = \sin^2 \theta \cdot \frac{1 - \cos^2 \theta}{\cos^2 \theta} = \sin^2 \theta \cdot \frac{\sin^2 \theta}{\cos^2 \theta} = \sin^4 \theta \cdot \sec^2 \theta.$$



$$\begin{aligned}
 (16) \quad & (\sec\theta - \operatorname{cosec}\theta)(1 + \cot\theta + \tan\theta) = \left(\frac{1}{\cos\theta} - \frac{1}{\sin\theta}\right)\left(1 + \frac{\cos\theta}{\sin\theta} + \frac{\sin\theta}{\cos\theta}\right) \\
 & = \frac{\sin\theta - \cos\theta}{\sin\theta \cdot \cos\theta} \cdot \frac{\sin\theta \cdot \cos\theta + 1}{\sin\theta \cdot \cos\theta} = \frac{\sin^2\theta \cdot \cos\theta + \sin\theta - \sin\theta \cdot \cos^2\theta - \cos\theta}{\sin^2\theta \cdot \cos^2\theta} \\
 & = \frac{(1 - \cos^2\theta)\cos\theta + \sin\theta - \sin\theta(1 - \sin^2\theta) - \cos\theta}{\sin^2\theta \cdot \cos^2\theta} \\
 & = \frac{\cos\theta - \cos^3\theta + \sin\theta - \sin\theta + \sin^3\theta - \cos\theta}{\sin^2\theta \cdot \cos^2\theta} \\
 & = \frac{\sin^3\theta - \cos^3\theta}{\sin^2\theta \cdot \cos^2\theta} = \frac{\sin\theta}{\cos^2\theta} - \frac{\cos\theta}{\sin^2\theta} = \operatorname{cosec}\theta - \sec\theta.
 \end{aligned}$$

$$\begin{aligned}
 (17) \quad & \frac{\operatorname{cosec}\theta}{\sec\theta} + \frac{\sec\theta}{\operatorname{cosec}\theta} = \frac{\cos\theta}{\sin\theta} + \frac{\sin\theta}{\cos\theta} = \frac{\cos^2\theta + \sin^2\theta}{\sin\theta \cdot \cos\theta} \\
 & = \frac{1}{\sin\theta \cdot \cos\theta} = \sec\theta \cdot \operatorname{cosec}\theta.
 \end{aligned}$$

$$\begin{aligned}
 (18) \quad & \cos\theta(\tan\theta + 2)(2\tan\theta + 1) = \cos\theta(2\tan^2\theta + 5\tan\theta + 2) \\
 & = 2\cos\theta(\tan^2\theta + 1) + 5\cos\theta \cdot \tan\theta \\
 & = 2\cos\theta \cdot \sec^2\theta + 5 \cdot \sin\theta = 2\sec\theta + 5\sin\theta.
 \end{aligned}$$

$$\begin{aligned}
 (19) \quad & \cos x(2\sec x + \tan x)(\sec x - 2\tan x) \\
 & = \cos x(2\sec^2 x - 3\sec x \cdot \tan x - 2\tan^2 x) \\
 & = 2\cos x(\sec^2 x - \tan^2 x) - 3\cos x \cdot \sec x \cdot \tan x \\
 & = 2\cos x - 3\tan x.
 \end{aligned}$$

$$\begin{aligned}
 (20) \quad & (\operatorname{cosec}\theta - \cot\theta)^2 = \operatorname{cosec}^2\theta - 2\operatorname{cosec}\theta \cdot \cot\theta + \cot^2\theta \\
 & = \frac{1}{\sin^2\theta} - \frac{2\cos\theta}{\sin^2\theta} + \frac{\cos^2\theta}{\sin^2\theta} \\
 & = \frac{1 - 2\cos\theta + \cos^2\theta}{\sin^2\theta} = \frac{(1 - \cos\theta)^2}{1 - \cos^2\theta} \\
 & = \frac{(1 - \cos\theta)(1 - \cos\theta)}{(1 + \cos\theta)(1 - \cos\theta)} \\
 & = \frac{1 - \cos\theta}{1 + \cos\theta}.
 \end{aligned}$$

$$\begin{aligned}
 (21) \quad & \frac{\sec\theta \cdot \cot\theta - \operatorname{cosec}\theta \cdot \tan\theta}{\cos\theta - \sin\theta} = \frac{\frac{1}{\sin\theta} - \frac{1}{\cos\theta}}{\cos\theta - \sin\theta} = \frac{\frac{\cos\theta - \sin\theta}{\sin\theta \cdot \cos\theta}}{\cos\theta - \sin\theta} \\
 & = \frac{1}{\sin\theta \cdot \cos\theta} = \operatorname{cosec}\theta \cdot \sec\theta.
 \end{aligned}$$

$$\begin{aligned}
 (22) \quad \sec\theta + \operatorname{cosec}\theta \cdot \tan^2\theta(1 + \operatorname{cosec}^2\theta) &= \frac{1}{\cos\theta} + \frac{\sin^2\theta}{\cos^3\theta} + \frac{1}{\cos^3\theta} \\
 &= \frac{\cos^2\theta + \sin^2\theta + 1}{\cos^3\theta} = \frac{2}{\cos^3\theta} = 2 \sec^3\theta.
 \end{aligned}$$

$$\begin{aligned}
 (23) \quad &(\sin\theta + \sec\theta)^2 + (\cos\theta + \operatorname{cosec}\theta)^2 \\
 &= \sin^2\theta + \frac{2\sin\theta}{\cos\theta} + \frac{1}{\cos^2\theta} + \cos^2\theta + \frac{2\cos\theta}{\sin\theta} + \frac{1}{\sin^2\theta} \\
 &= (\sin^2\theta + \cos^2\theta) + \left(\frac{1}{\cos^2\theta} + \frac{1}{\sin^2\theta}\right) + \left(\frac{2\sin\theta}{\cos\theta} + \frac{2\cos\theta}{\sin\theta}\right) \\
 &= 1 + \frac{1}{\sin^2\theta \cdot \cos^2\theta} + \frac{2}{\sin\theta \cdot \cos\theta} = \left(1 + \frac{1}{\sin\theta \cdot \cos\theta}\right)^2 = (1 + \sec\theta \cdot \operatorname{cosec}\theta)^2.
 \end{aligned}$$

$$\begin{aligned}
 (24) \quad &\frac{1 + (\operatorname{cosec}\theta \cdot \tan\phi)^2}{1 + (\operatorname{cosec}a \cdot \tan\phi)^2} = \frac{1 + \frac{\sin^2\phi}{\sin^2\theta \cdot \cos^2\theta}}{1 + \frac{\sin^2\phi}{\sin^2a \cdot \cos^2a}} = \frac{\sin^2\theta \cdot \cos^2\phi + \sin^2\phi \cdot \frac{\sin^2a}{\sin^2\theta}}{\sin^2a \cdot \cos^2\phi + \sin^2\phi \cdot \frac{\sin^2a}{\sin^2\theta}} \\
 &= \frac{\sin^2\theta(1 - \sin^2\phi) + \sin^2\phi \cdot \frac{\sin^2a}{\sin^2\theta}}{\sin^2a(1 - \sin^2\phi) + \sin^2\phi \cdot \frac{\sin^2a}{\sin^2\theta}} = \frac{\sin^2\theta - \sin^2\theta \cdot \sin^2\phi + \sin^2\phi \cdot \frac{\sin^2a}{\sin^2\theta}}{\sin^2a - \sin^2a \cdot \sin^2\phi + \sin^2\phi \cdot \frac{\sin^2a}{\sin^2\theta}} \\
 &= \frac{\sin^2\theta + \sin^2\phi \cdot \frac{\cos^2\theta}{\cos^2a} \cdot \frac{\sin^2a}{\sin^2\theta}}{\sin^2a + \sin^2\phi \cdot \frac{\cos^2a}{\cos^2\theta} \cdot \frac{\sin^2\theta}{\sin^2a}} \\
 &= \frac{1 + \sin^2\phi \cdot \cot^2\theta}{1 + \sin^2\phi \cdot \cot^2a} = \frac{1 + (\cot\theta \cdot \sin\phi)^2}{1 + (\cot a \cdot \sin\phi)^2}.
 \end{aligned}$$

$$\begin{aligned}
 (25) \quad &(3 - 4 \sin^2 A)(1 - 3 \tan^2 A) = (3 - 4 \sin^2 A) \left(1 - \frac{3 \sin^2 A}{\cos^2 A}\right) \\
 &= (3 - 4 \sin^2 A) \left(\frac{\cos^2 A - 3 \sin^2 A}{\cos^2 A}\right) \\
 &= (3 - 4 \sin^2 A) \left(\frac{\cos^2 A - 3(1 - \cos^2 A)}{\cos^2 A}\right) \\
 &= \frac{3 - 4 \sin^2 A}{\cos^2 A} \cdot (4 \cos^2 A - 3) \\
 &= \frac{3 \cos^2 A + 3 \sin^2 A - 4 \sin^2 A}{\cos^2 A} (4 \cos^2 A - 3) \\
 &= \frac{3 \cos^2 A - \sin^2 A}{\cos^2 A} (4 \cos^2 A - 3) \\
 &= (3 - \tan^2 A) (4 \cos^2 A - 3).
 \end{aligned}$$

EXAMPLES—XX. (p. 65).

1. (1)  $90^\circ - (24^\circ. 14'. 42'') = 65^\circ. 45'. 18''.$   
 (2)  $90^\circ - (43^\circ. 2'. 57'') = 46^\circ. 57'. 3''.$   
 (3)  $90^\circ - (64^\circ. 0'. 14'') = 25^\circ. 59'. 46''.$   
 (4)  $90^\circ - (82^\circ. 4'. 15'') = 7^\circ. 55'. 45''.$   
 (5)  $90^\circ - (125^\circ. 15'. 42'') = - (35^\circ. 15'. 42'').$   
 (6)  $90^\circ - (178^\circ. 27'. 34'') = - (88^\circ. 27'. 34'').$   
 (7)  $90^\circ - 195^\circ = - 105^\circ.$   
 (8)  $90^\circ - 254^\circ = - 164^\circ.$   
 (9)  $90^\circ - (- 25^\circ) = 90^\circ + 25^\circ = 115^\circ.$   
 (10)  $90^\circ - (- 245^\circ) = 90^\circ + 245^\circ = 335^\circ.$

2. (1)  $100^\circ - (32^\circ. 23'. 24'') = 67^\circ. 76'. 76''.$   
 (2)  $100^\circ - (95^\circ. 3'. 75'') = 4^\circ. 96'. 25''.$   
 (3)  $100^\circ - (46^\circ. 0'. 84'') = 53^\circ. 99'. 16''.$   
 (4)  $100^\circ - (2^\circ. 5'. 4'') = 97^\circ. 94'. 96''.$   
 (5)  $100^\circ - (135^\circ. 2'. 5'') = - (35^\circ. 2'. 5'').$   
 (6)  $100^\circ - (169^\circ. 0'. 3'') = - (69^\circ. 0'. 3'').$   
 (7)  $100^\circ - 243^\circ = - 143^\circ.$   
 (8)  $100^\circ - 357^\circ = - 257^\circ.$   
 (9)  $100^\circ - (- 35^\circ) = 100^\circ + 35^\circ = 135^\circ.$   
 (10)  $100^\circ - (- 245^\circ) = 100^\circ + 245^\circ = 345^\circ.$

3. (1)  $\frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}.$       (2)  $\frac{\pi}{2} - \frac{\pi}{3} = \frac{\pi}{6}.$       (3)  $\frac{\pi}{2} - \frac{3\pi}{5} = - \frac{\pi}{10}.$   
 (4)  $\frac{\pi}{2} - \left(- \frac{\pi}{4}\right) = \frac{\pi}{2} + \frac{\pi}{4} = \frac{3\pi}{4}.$       (5)  $\frac{\pi}{2} - \left(- \frac{3\pi}{4}\right) = \frac{\pi}{2} + \frac{3\pi}{4} = \frac{5\pi}{4}.$



## EXAMPLES—XXI. (p. 68).

1. (1)  $180^\circ - (34^\circ. 12'. 49'') = 145^\circ. 47'. 11''$ .  
 (2)  $180^\circ - (132^\circ. 24'. 47'') = 47^\circ. 35'. 13''$ .  
 (3)  $180^\circ - (146^\circ. 0'. 41'') = 33^\circ. 59'. 19''$ .  
 (4)  $180^\circ - (28^\circ. 15'. 4'') = 151^\circ. 44'. 56''$ .  
 (5)  $180^\circ - (179^\circ. 59'. 59'') = 1''$ .  
 (6)  $180^\circ - (100^\circ. 49'. 53'') = 79^\circ. 10'. 7''$ .  
 (7)  $180^\circ - 245^\circ = -65^\circ$ .  
 (8)  $180^\circ - (437^\circ. 3'. 4'') = -(257^\circ. 3'. 4'')$ .  
 (9)  $180^\circ - (-49^\circ) = 180^\circ + 49^\circ = 229^\circ$ .  
 (10)  $180^\circ - (-355^\circ) = 180^\circ + 355^\circ = 535^\circ$ .
  
2. (1)  $200^\circ - (132^\circ. 32'. 42'') = 67^\circ. 67'. 58''$ .  
 (2)  $200^\circ - (195^\circ. 2'. 57'') = 4^\circ. 97'. 43''$ .  
 (3)  $200^\circ - (3^\circ. 97'. 98'') = 196^\circ. 2'. 2''$ .  
 (4)  $200^\circ - (65^\circ. 12'. 8'') = 134^\circ. 87'. 92''$ .  
 (5)  $200^\circ - (154^\circ. 3'. 6'') = 45^\circ. 96'. 94''$ .  
 (6)  $200^\circ - (174^\circ. 0'. 4'') = 25^\circ. 99'. 96''$ .  
 (7)  $200^\circ - 275^\circ = -75^\circ$ .  
 (8)  $200^\circ - (527^\circ. 2'. 14'') = (327^\circ. 2'. 14'')$ .  
 (9)  $200^\circ - (-35^\circ) = 200^\circ + 35^\circ = 235^\circ$ .  
 (10)  $200^\circ - (-325^\circ) = 200^\circ + 325^\circ = 525^\circ$ .
  
3. (1)  $\pi - \frac{\pi}{2} = \frac{\pi}{2}$ .      (2)  $\pi - \frac{\pi}{3} = \frac{2\pi}{3}$ .      (3)  $\pi - \frac{4\pi}{5} = \frac{\pi}{5}$ .  
 (4)  $\pi - \left(-\frac{\pi}{4}\right) = \pi + \frac{\pi}{4} = \frac{5\pi}{4}$ .      (5)  $\pi - \left(-\frac{3\pi}{4}\right) = \pi + \frac{3\pi}{4} = \frac{7\pi}{4}$ .

4. Let  $\theta$  be the circular measure of the angle.

Then  $\frac{\pi}{2} - \theta$  is the complement of  $\theta$ ;

and  $\pi - \left(\frac{\pi}{2} - \theta\right)$ , or,  $\frac{\pi}{2} + \theta$  is the supplement of the complement of  $\theta$ .

Again  $\pi - \theta$  is the supplement of  $\theta$ ,

and  $\frac{\pi}{2} - (\pi - \theta)$ , or,  $\theta - \frac{\pi}{2}$  is the complement of the supplement of  $\theta$ ;

$$\therefore \text{difference} = \frac{\pi}{2} + \theta - \left(\theta - \frac{\pi}{2}\right) = \frac{\pi}{2} + \frac{\pi}{2} = \pi.$$

# EXAMPLES—XXII. (p. 72).

1. (1) Take the construction and notation of Art. 101.

$$\text{Then } \sec(180^\circ - A) = \sec EOP' = \frac{OP'}{OM'} = \frac{OP}{-OM} = -\sec A.$$

- (2) Take the construction of Art. 102, and let  $\angle EOP = \theta$ .

$$\text{Then } \operatorname{cosec}\left(\frac{\pi}{2} + \theta\right) = \frac{OP'}{P'M'} = \frac{OP}{OM} = \sec \theta.$$

- (3) Take the construction and notation of Art. 103.

$$\text{Then } \tan(180^\circ + A) = \frac{P'M'}{OM'} = \frac{-PM}{-OM} = \frac{PM}{OM} = \tan A.$$

- (4) Take the construction of Art. 103, and let  $\angle EOP = \theta$ .

$$\text{Then } \sec(\pi + \theta) = \frac{OP'}{OM'} = \frac{OP}{-OM} = -\sec \theta.$$

- (5) Take the construction of Art. 104, and let  $\angle EOP = \theta$ .

$$\text{Then } \tan(-\theta) = \frac{MP'}{MO} = \frac{-MP}{MO} = -\tan \theta.$$

- (6) Take the construction of Art. 104, and let  $\angle EOP = \theta$ .

$$\text{Then } \cot(2\pi - \theta) = \cot EOP' = \frac{OM}{MP'} = \frac{OM}{-MP} = -\cot \theta.$$

2. (1) Take the construction of Art. 102, and let  $\angle EOP = B$ .

$$\text{Then } \operatorname{cosec}(90^\circ + B) = \operatorname{cosec} EOP' = \frac{OP'}{P'M'} = \frac{OP}{OM} = \sec B = \frac{\operatorname{cosec} B}{\sqrt{\operatorname{cosec}^2 B - 1}}.$$

(Ex. XVII. 2.)

(2) Take the construction of Art. 103, and let  $\angle EOP = \phi$ .

$$\text{Then } \operatorname{cosec}(\pi + \phi) = \operatorname{cosec} EOP' = \frac{OP'}{P'M'} = \frac{OP}{-PM} = -\operatorname{cosec} \phi.$$

3. (1) Take the construction of Art. 102, and let  $\angle EOP = A$ .

$$\text{Then } \sec(90^\circ + A) = \sec EOP' = \frac{OP'}{OM'} = \frac{OP}{-PM} = -\operatorname{cosec} A = -\frac{\sec A}{\sqrt{\sec^2 A - 1}}.$$

(Ex. XVII. 3.)

(2) Take the construction of Art. 99, and let  $\angle EOP = \theta$ .

$$\text{Then } \sec\left(\frac{\pi}{2} - \theta\right) = \sec EOP' = \frac{OP'}{OM'} = \frac{OP}{PM} = \operatorname{cosec} \theta = \frac{\sec \theta}{\sqrt{\sec^2 \theta - 1}}.$$

(Ex. XVII. 3.)

#### EXAMPLES—XXIII. (p. 72).

$$(1) \sin 120^\circ = \sin(180^\circ - 120^\circ) = \sin 60^\circ = \frac{\sqrt{3}}{2}.$$

$$(2) \cos 120^\circ = -\cos(180^\circ - 120^\circ) = -\cos 60^\circ = -\frac{1}{2}.$$

$$(3) \sin 135^\circ = \sin(180^\circ - 135^\circ) = \sin 45^\circ = \frac{1}{\sqrt{2}}.$$

$$(4) \cos 135^\circ = -\cos(180^\circ - 135^\circ) = -\cos 45^\circ = -\frac{1}{\sqrt{2}}.$$

$$(5) \sin 150^\circ = \sin(180^\circ - 150^\circ) = \sin 30^\circ = \frac{1}{2}.$$

$$(6) \cos 150^\circ = -\cos(180^\circ - 150^\circ) = -\cos 30^\circ = -\frac{\sqrt{3}}{2}.$$

$$(7) \sin 225^\circ = \sin(180^\circ + 45^\circ) = -\sin 45^\circ = -\frac{1}{\sqrt{2}}.$$

$$(8) \sin 240^\circ = \sin(180^\circ + 60^\circ) = -\sin 60^\circ = -\frac{\sqrt{3}}{2}.$$

$$(9) \tan 300^\circ = \tan(360^\circ - 60^\circ) = -\tan 60^\circ = -\sqrt{3}.$$

$$(10) \operatorname{cosec} 300^\circ = \operatorname{cosec}(360^\circ - 60^\circ) = -\operatorname{cosec} 60^\circ = -\frac{2}{\sqrt{3}}.$$

$$(11) \sec 315^\circ = \sec(360^\circ - 45^\circ) = \sec 45^\circ = \sqrt{2}.$$

$$(12) \cot 330^\circ = \cot(360^\circ - 30^\circ) = -\cot 30^\circ = -\sqrt{3}.$$

EXAMPLES—XXIV. (p. 75).

$$(1) \sin \theta + \cos \theta = 0,$$

$$\sin \theta = -\cos \theta,$$

$$\sin^2 \theta = \cos^2 \theta,$$

$$\sin^2 \theta = 1 - \sin^2 \theta,$$

$$2 \sin^2 \theta = 1.$$

$$\text{Hence } \sin \theta = \pm \frac{1}{\sqrt{2}}, \text{ and } \therefore \theta = 45^\circ \text{ or } -45^\circ.$$

The latter of these values must be taken, because  $\sin \theta$  and  $\cos \theta$  must have different signs to satisfy the equation.

$$(2) \sin \theta - \cos \theta = 0,$$

$$\sin \theta = \cos \theta,$$

$$\text{and, as in Example (1), } \theta = 45^\circ \text{ or } -45^\circ.$$

The former of these values must be taken, because  $\sin \theta$  and  $\cos \theta$  must have the same sign to satisfy the equation.

$$(3) \sin \theta = \tan \theta,$$

$$\sin \theta = \frac{\sin \theta}{\cos \theta}, \text{ and, dividing by } \sin \theta,$$

$$1 = \frac{1}{\cos \theta}, \text{ or, } \cos \theta = 1, \text{ and } \therefore \theta = 0^\circ.$$

$$(4) \cos \theta = \cot \theta,$$

$$\cos \theta = \frac{\cos \theta}{\sin \theta}, \text{ and, dividing by } \cos \theta,$$

$$1 = \frac{1}{\sin \theta}, \text{ or, } \sin \theta = 1, \text{ and } \therefore \theta = 90^\circ.$$

$$(5) \quad 2 \sin \theta = \tan \theta,$$

$$2 \sin \theta = \frac{\sin \theta}{\cos \theta}, \text{ or, } 2 \cos \theta = 1, \text{ or, } \cos \theta = \frac{1}{2}, \text{ and, } \therefore, \theta = 60^\circ.$$

Also, since we divided by  $\sin \theta$ , one value of  $\theta$  to satisfy the original equation is given by  $\sin \theta = 0$ , or,  $\theta = 0^\circ$ .

(6)

$$3 \sin \theta = 2 \cos^2 \theta,$$

$$3 \sin \theta = 2(1 - \sin^2 \theta),$$

$$2 \sin^2 \theta + 3 \sin \theta = 2,$$

$$\sin^2 \theta + \frac{3}{2} \sin \theta = 1.$$

$$\sin^2 \theta + \frac{3}{2} \sin \theta + \frac{9}{16} = \frac{25}{16}.$$

$$\sin \theta + \frac{3}{4} = \pm \frac{5}{4}.$$

$$\text{Hence } \sin \theta = \frac{1}{2}, \text{ or, } -2.$$

The second value is inadmissible

$$\therefore \sin \theta = \frac{1}{2}, \text{ or, } \theta = 30^\circ.$$

(7)

$$\sin \theta + \cos^2 \theta \cdot \operatorname{cosec} \theta = 2,$$

$$\sin \theta + \frac{\cos^2 \theta}{\sin \theta} = 2,$$

$$\sin^2 \theta + \cos^2 \theta = 2 \sin \theta,$$

$$1 = 2 \sin \theta.$$

$$\text{Hence } \sin \theta = \frac{1}{2}, \text{ or, } \theta = 30^\circ.$$

(8)

$$\tan \theta = 4 - 3 \cot \theta,$$

$$\tan \theta + 3 \cot \theta = 4,$$

$$\tan \theta + \frac{3}{\tan \theta} = 4,$$

$$\tan^2 \theta + 3 = 4 \tan \theta,$$

$$\tan^2 \theta - 4 \tan \theta = -3,$$

$$\tan^2 \theta - 4 \tan \theta + 4 = 1,$$

$$\tan \theta - 2 = \pm 1.$$

Hence  $\tan \theta = 3$  or  $1$ , and the latter of these values of  $\tan \theta$  enables us to say that *one* value of  $\theta$  is  $45^\circ$ .



$$\begin{aligned}
 (9) \quad & 4 \sec^2 \theta - 7 \tan^2 \theta = 3, \\
 & 4(1 + \tan^2 \theta) - 7 \tan^2 \theta = 3, \\
 & 4 - 3 \tan^2 \theta = 3, \\
 & \tan^2 \theta = \frac{1}{3}, \text{ or, } \tan \theta = \frac{1}{\sqrt{3}}, \text{ and } \therefore \theta = 30^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (10) \quad & \cos \theta \cdot \operatorname{cosec} \theta + \sin \theta \cdot \sec \theta = \frac{4}{\sqrt{3}}, \\
 & \frac{\cos \theta}{\sin \theta} + \frac{\sin \theta}{\cos \theta} = \frac{4}{\sqrt{3}}, \\
 & \frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta \cdot \cos \theta} = \frac{4}{\sqrt{3}}, \\
 & \sqrt{3} = 4 \sin \theta \cdot \cos \theta, \\
 & 3 = 16 \sin^2 \theta (1 - \sin^2 \theta), \\
 & 16 \sin^4 \theta - 16 \sin^2 \theta = -3, \\
 & \sin^4 \theta - \sin^2 \theta = -\frac{3}{16}.
 \end{aligned}$$

$$\text{Hence } \sin^2 \theta = \frac{3}{4} \text{ or } \frac{1}{4},$$

$$\text{and } \therefore \sin \theta = \frac{\sqrt{3}}{2} \text{ or } \frac{1}{2}, \text{ and } \theta = 60^\circ \text{ or } 30^\circ.$$

$$\begin{aligned}
 (11) \quad & 3 \sin^2 \theta - \cos^2 \theta + (\sqrt{3} + 1)(1 - 2 \sin \theta) = 0, \\
 & 3 \sin^2 \theta - (1 - \sin^2 \theta) + \sqrt{3} + 1 - 2\sqrt{3} \sin \theta - 2 \sin \theta = 0, \\
 & 4 \sin^2 \theta - 2(\sqrt{3} + 1) \sin \theta = -\sqrt{3}, \\
 & \sin^2 \theta - \frac{\sqrt{3} + 1}{2} \sin \theta = -\frac{\sqrt{3}}{4}, \\
 & \sin^2 \theta - \frac{\sqrt{3} + 1}{2} \sin \theta + \frac{4 + 2\sqrt{3}}{16} = \frac{4 + 2\sqrt{3}}{16} - \frac{\sqrt{3}}{4} = \frac{4 - 2\sqrt{3}}{16}, \\
 & \sin \theta - \frac{\sqrt{3} + 1}{4} = \pm \frac{\sqrt{3} - 1}{4}.
 \end{aligned}$$

$$\text{Hence } \sin \theta = \frac{\sqrt{3}}{2} \text{ or } \frac{1}{2}, \text{ and } \theta = 60^\circ \text{ or } 30^\circ.$$

$$\begin{aligned}
 (12) \quad & 3 \cos^2 \theta - \sin^2 \theta + (\sqrt{3} + 1)(1 - 2 \cos \theta) = 0, \\
 & 3 \cos^2 \theta - (1 - \cos^2 \theta) + \sqrt{3} + 1 - 2\sqrt{3} \cos \theta - 2 \cos \theta = 0, \\
 & 4 \cos^2 \theta - 2(\sqrt{3} + 1) \cos \theta = -\sqrt{3}.
 \end{aligned}$$

Hence, by the same process as in Example (11),

$$\cos \theta = \frac{\sqrt{3}}{2} \text{ or } \frac{1}{2}, \text{ and } \theta = 30^\circ \text{ or } 60^\circ.$$

$$(13) \quad \sec \theta \cdot \operatorname{cosec} \theta + 2 \cot \theta = 4,$$

$$\frac{1}{\cos \theta \cdot \sin \theta} + \frac{2 \cos \theta}{\sin \theta} = 4,$$

$$1 + 2 \cos^2 \theta = 4 \sin \theta \cdot \cos \theta,$$

$$1 + 4 \cos^2 \theta + 4 \cos^4 \theta = 16 \sin^2 \theta \cdot \cos^2 \theta,$$

$$1 + 4 \cos^2 \theta + 4 \cos^4 \theta = 16 \cos^2 \theta - 16 \cos^4 \theta,$$

$$20 \cos^4 \theta - 12 \cos^2 \theta = -1,$$

$$\cos^4 \theta - \frac{3}{5} \cos^2 \theta = -\frac{1}{20}.$$

$$\text{Hence } \cos^2 \theta = \frac{1}{2}, \text{ and } \cos \theta = \frac{1}{\sqrt{2}}, \text{ and } \theta = 45^\circ$$

$$(14) \quad \sin \theta + \cos \theta = \sqrt{2}, \quad . \quad . \quad . \quad . \quad (1),$$

$$\sin^2 \theta + 2 \sin \theta \cdot \cos \theta + \cos^2 \theta = 2, \quad . \quad . \quad . \quad . \quad (2),$$

$$2 \sin \theta \cdot \cos \theta = 1,$$

$$4 \sin \theta \cdot \cos \theta = 2, \text{ and, subtracting this from (2),}$$

$$\sin^2 \theta - 2 \sin \theta \cdot \cos \theta + \cos^2 \theta = 0,$$

$$\sin \theta - \cos \theta = 0, \text{ and, adding this to (1),}$$

$$2 \sin \theta = \sqrt{2}, \therefore \sin \theta = \frac{1}{\sqrt{2}}, \text{ and } \theta = 45^\circ.$$

$$(15) \quad \cot^2 \theta + 4 \cos^2 \theta = 6,$$

$$\cos^2 \theta + 4 \cos^2 \theta \cdot \sin^2 \theta = 6 \sin^2 \theta,$$

$$\cos^2 \theta + 4 \cos^2 \theta - 4 \cos^4 \theta = 6 - 6 \cos^2 \theta,$$

$$\cos^4 \theta - \frac{11}{4} \cos^2 \theta = -\frac{3}{2}.$$

$$\text{Hence } \cos^2 \theta = \frac{3}{4} \text{ or } 2.$$

The latter value is inadmissible, and we must have

$$\cos^2 \theta = \frac{3}{4}, \text{ or, } \cos \theta = \frac{\sqrt{3}}{2}, \text{ and } \theta = 30^\circ.$$

$$(16) \quad \tan \theta + \cot \theta = 2.$$

$$\tan \theta + \frac{1}{\tan \theta} = 2,$$

$$\tan^2 \theta - 2 \tan \theta = -1;$$

$$\therefore \tan \theta = 1, \text{ and } \theta = 45^\circ.$$

$$(17) \quad \sin\theta - \cos\theta = \sqrt{2}, \quad . \quad . \quad . \quad . \quad (1)$$

$$\sin^2\theta - 2\sin\theta \cdot \cos\theta + \cos^2\theta = 2, \quad . \quad . \quad . \quad . \quad (2)$$

$$-2\sin\theta \cdot \cos\theta = 1,$$

$$4\sin\theta \cdot \cos\theta = -2, \text{ and adding this to (2)}$$

$$\sin^2\theta + 2\sin\theta \cdot \cos\theta + \cos^2\theta = 0,$$

$$\sin\theta + \cos\theta = 0, \text{ and adding this to (1)}$$

$$2\sin\theta = \sqrt{2}, \text{ or, } \sin\theta = \frac{1}{\sqrt{2}}, \text{ and } \theta = 45^\circ \text{ or } 135^\circ.$$

Now  $\cos\theta$  has to be of the same numerical value as  $\sin\theta$ , but with a different sign, and hence  $45^\circ$  is an inadmissible value of  $\theta$ ;

$$\therefore \theta = 135^\circ.$$

$$(18) \quad \sin\theta + \cos\theta = 2\sqrt{2} \cdot \sin\theta \cdot \cos\theta,$$

$$\sin^2\theta + 2\sin\theta \cdot \cos\theta + \cos^2\theta = 8\sin^2\theta \cdot \cos^2\theta,$$

$$8\sin^2\theta \cdot \cos^2\theta - 2\sin\theta \cdot \cos\theta = 1,$$

$$\sin^2\theta \cdot \cos^2\theta - \frac{1}{4} \cdot \sin\theta \cdot \cos\theta = \frac{1}{8},$$

$$\sin^2\theta \cdot \cos^2\theta - \frac{1}{4} \sin\theta \cdot \cos\theta + \frac{1}{64} = \frac{9}{64};$$

$$\therefore \sin\theta \cdot \cos\theta = \frac{1}{2} \text{ or } -\frac{1}{4}.$$

Taking the former of these values, we get

$$\sin^2\theta (1 - \sin^2\theta) = \frac{1}{4}.$$

$$\text{Whence } \sin^2\theta = \frac{1}{2}, \text{ or, } \sin\theta = \frac{1}{\sqrt{2}}, \text{ and } \theta = 45^\circ.$$

$$(19) \quad \sqrt{3} \cdot \sin\theta = \sqrt{3} - \cos\theta,$$

$$3\sin^2\theta = 3 - 2\sqrt{3} \cdot \cos\theta + \cos^2\theta,$$

$$3 - 3\cos^2\theta = 3 - 2\sqrt{3} \cos\theta + \cos^2\theta,$$

$$4\cos^2\theta = 2\sqrt{3} \cdot \cos\theta.$$

Dividing by  $\cos\theta$ , we get

$$4\cos\theta = 2\sqrt{3}, \text{ or, } \cos\theta = 0.$$

$$\text{Hence } \cos\theta = \frac{\sqrt{3}}{2}, \text{ or, } \cos\theta = 0;$$

$$\therefore \theta = 30^\circ \text{ or } 90^\circ.$$



$$\begin{aligned}
 (20) \quad & \tan^2 \theta + 4 \sin^2 \theta = 3, \\
 & \sin^2 \theta + 4 \sin^2 \theta \cdot \cos^2 \theta = 3 \cos^2 \theta, \\
 & 1 - \cos^2 \theta + 4 \cos^2 \theta - 4 \cos^4 \theta = 3 \cos^2 \theta, \\
 & 4 \cos^4 \theta = 1; \\
 & \therefore \text{one value of } \cos \theta \text{ is } \frac{1}{\sqrt{2}}, \text{ or, } \theta = 45^\circ.
 \end{aligned}$$

## EXAMPLES—XXV. (p. 81).

- (1)  $\sin 480^\circ = \sin(360^\circ + 120^\circ) = \sin 120^\circ = \sin 60^\circ = \frac{\sqrt{3}}{2}.$
- (2)  $\cos 480^\circ = \cos(360^\circ + 120^\circ) = \cos 120^\circ = -\cos 60^\circ = -\frac{1}{2}.$
- (3)  $\sin 495^\circ = \sin(360^\circ + 135^\circ) = \sin 135^\circ = \sin 45^\circ = \frac{1}{\sqrt{2}}.$
- (4)  $\cos 495^\circ = \cos(360^\circ + 135^\circ) = \cos 135^\circ = -\cos 45^\circ = -\frac{1}{\sqrt{2}}.$
- (5)  $\sin 870^\circ = \sin(720^\circ + 150^\circ) = \sin 150^\circ = \sin 30^\circ = \frac{1}{2}.$
- (6)  $\cos 870^\circ = \cos(720^\circ + 150^\circ) = \cos 150^\circ = -\cos 30^\circ = -\frac{\sqrt{3}}{2}.$
- (7)  $\sin 945^\circ = \sin(720^\circ + 225^\circ) = \sin 225^\circ = -\sin 45^\circ = -\frac{1}{\sqrt{2}}.$
- (8)  $\sin 960^\circ = \sin(720^\circ + 240^\circ) = \sin 240^\circ = -\sin 60^\circ = -\frac{\sqrt{3}}{2}.$
- (9)  $\tan 1020^\circ = \tan(720^\circ + 300^\circ) = \tan 300^\circ = -\tan 60^\circ = -\sqrt{3}.$
- (10)  $\operatorname{cosec} 1380^\circ = \operatorname{cosec}(1080^\circ + 300^\circ) = \operatorname{cosec} 300^\circ = -\operatorname{cosec} 60^\circ = -\frac{2}{\sqrt{3}}.$
- (11)  $\sec 1395^\circ = \sec(1080^\circ + 315^\circ) = \sec 315^\circ = \sec 45^\circ = \sqrt{2}.$
- (12)  $\cot 1410^\circ = \cot(1080^\circ + 330^\circ) = \cot 330^\circ = -\cot 30^\circ = -\sqrt{3}.$

$$(13) \cos 420^\circ = \cos(360^\circ + 60^\circ) = \cos 60^\circ = \frac{1}{2}.$$

$$(14) \sec 750^\circ = \sec(720^\circ + 30^\circ) = \sec 30^\circ = \frac{2}{\sqrt{3}}.$$

$$(15) \tan 945^\circ = \tan(720^\circ + 225^\circ) = \tan 225^\circ = \tan 45^\circ = 1.$$

$$(16) \sin 1200^\circ = \sin(1080^\circ + 120^\circ) = \sin 120^\circ = \sin 60^\circ = \frac{\sqrt{3}}{2}.$$

$$(17) \sin 1485^\circ = \sin(1440^\circ + 45^\circ) = \sin 45^\circ = \frac{1}{\sqrt{2}}.$$

$$(18) \cos 1470^\circ = \cos(1440^\circ + 30^\circ) = \cos 30^\circ = \frac{\sqrt{3}}{2}.$$

$$(19) \sin 7\pi = \sin(6\pi + \pi) = \sin \pi = 0.$$

$$(20) \sec 8\pi = \sec 2\pi = 1.$$

$$(21) \operatorname{cosec} 930^\circ = \operatorname{cosec}(720^\circ + 210^\circ) = \operatorname{cosec} 210^\circ = -\operatorname{cosec} 30^\circ = -2.$$

$$(22) \cot 1140^\circ = \cot(1080^\circ + 60^\circ) = \cot 60^\circ = \frac{1}{\sqrt{3}}.$$

$$(23) \tan 1305^\circ = \tan(1080^\circ + 225^\circ) = \tan 225^\circ = \tan 45^\circ = 1.$$

$$(24) \operatorname{cosec} 1740^\circ = \operatorname{cosec}(1440^\circ + 300^\circ) = \operatorname{cosec} 300^\circ = -\operatorname{cosec} 60^\circ = -\frac{2}{\sqrt{3}}.$$

$$(25) \sin(-240^\circ) = -\sin 240^\circ = -\sin(-60^\circ) = \sin 60^\circ = \frac{\sqrt{3}}{2}.$$

$$(26) \cot(-675^\circ) = \cot(720^\circ - 675^\circ) = \cot 45^\circ = 1.$$

$$(27) \sec(-135^\circ) = -\sec(180^\circ - 135^\circ) = -\sec 45^\circ = -\sqrt{2}.$$

$$(28) \tan(-225^\circ) = \tan(360^\circ - 225^\circ) = \tan 135^\circ = -\tan 45^\circ = -1.$$

$$(29) \operatorname{cosec}(-690^\circ) = \operatorname{cosec}(720^\circ - 690^\circ) = \operatorname{cosec} 30^\circ = 2.$$

$$(30) \cos(-120^\circ) = \cos 120^\circ = -\cos 60^\circ = -\frac{1}{2}.$$

EXAMPLES—XXVI. (p. 82).

(1)  $\sin\theta=1$  ;  $\therefore$  one value of  $\theta$  is  $\frac{\pi}{2}$  ;

$\therefore$  general value of  $\theta$  is  $n\pi + (-1)^n \cdot \frac{\pi}{2}$ .

(2)  $\cos\theta=1$  ;  $\therefore$  one value of  $\theta$  is 0 ;

$\therefore$  general value of  $\theta$  is  $2n\pi$ .

(3)  $\sin\theta=\frac{1}{\sqrt{2}}$  ;  $\therefore$  one value of  $\theta$  is  $\frac{\pi}{4}$  ;

$\therefore$  general value of  $\theta$  is  $n\pi + (-1)^n \cdot \frac{\pi}{4}$ .

(4)  $\tan\theta=\sqrt{3}$  ;  $\therefore$  one value of  $\theta$  is  $\frac{\pi}{3}$  ;

$\therefore$  general value of  $\theta$  is  $n\pi + \frac{\pi}{3}$ .

(5)  $3\sin\theta=2\cos^2\theta$

$3\sin\theta=2(1-\sin^2\theta)$

$\sin^2\theta + \frac{3}{2}\sin\theta = 1$

$\left(\sin\theta + \frac{3}{4}\right)^2 = \pm \frac{5}{4}$ , or,  $\sin\theta = \frac{1}{2}$  or  $-2$

$\therefore$  least positive value of  $\theta$  is  $\frac{\pi}{6}$  ;

$\therefore$  general value of  $\theta$  is  $n\pi + (-1)^n \cdot \frac{\pi}{6}$ .

(6)  $2\sin\theta=\tan\theta$ ,

$2\sin\theta=\frac{\sin\theta}{\cos\theta}$  ;

$\therefore \sin\theta=0$ , or,  $\cos\theta=\frac{1}{2}$  ;

$\therefore \theta=0$ , or,  $\theta=\frac{\pi}{3}$  ;

$\therefore$  general value of  $\theta$  is  $n\pi$  or  $2n\pi \pm \frac{\pi}{3}$ .

$$\begin{aligned}
 (7) \quad & \tan^2 \theta + 4 \sin^2 \theta = 3, \\
 & \sin^2 \theta + 4 \sin^2 \theta \cdot \cos^2 \theta = 3 \cos^2 \theta, \\
 & \sin^2 \theta + 4 \sin^2 \theta - 4 \sin^4 \theta = 3 - 3 \sin^2 \theta, \\
 & 4 \sin^4 \theta - 8 \sin^2 \theta = -3, \\
 & \sin^4 \theta - 2 \sin^2 \theta + 1 = \frac{1}{4},
 \end{aligned}$$

$$\sin^2 \theta - 1 = \pm \frac{1}{2}.$$

$$\text{Hence } \sin \theta = \pm \sqrt{\frac{3}{2}} \text{ or } \pm \frac{1}{\sqrt{2}};$$

$$\therefore \text{least positive value of } \theta \text{ is } \frac{\pi}{4};$$

$$\therefore \text{general value of } \theta \text{ is } n\pi + (-1)^n \cdot \frac{\pi}{4}.$$

$$\begin{aligned}
 (8) \quad & \cos^2 = \sin^2 \theta, \\
 & \cos^2 \theta = 1 - \cos^2 \theta,
 \end{aligned}$$

$$2 \cos^2 \theta = 1, \text{ and } \therefore \cos \theta = \pm \frac{1}{\sqrt{2}};$$

$$\therefore \text{the least positive values of } \theta \text{ are } \frac{\pi}{4} \text{ and } \frac{3\pi}{4};$$

$$\therefore \text{the general value of } \theta \text{ is } 2n\pi \pm \frac{\pi}{4} \text{ or } 2n\pi \pm \frac{3\pi}{4}.$$

$$(9) \quad \tan \theta = 4 - 3 \cot \theta,$$

$$\tan \theta + \frac{3}{\tan \theta} = 4,$$

$$\tan^2 \theta - 4 \tan \theta = -3,$$

$$\tan \theta = 3 \text{ or } 1;$$

$$\therefore \text{the least positive value of } \theta \text{ is } \frac{\pi}{4};$$

$$\therefore \text{general value of } \theta \text{ is } n\pi + \frac{\pi}{4}.$$

$$\begin{aligned}
 (10) \quad \sec^2\theta - \frac{5}{2}\sec\theta + 1 &= 0, \\
 \sec^2\theta - \frac{5}{2}\sec\theta + \frac{25}{16} &= \frac{9}{16}, \\
 \sec\theta - \frac{5}{4} &= \pm \frac{3}{4}; \\
 \therefore \sec\theta &= 2 \text{ or } \frac{1}{2}.
 \end{aligned}$$

Taking the value 2 for  $\sec\theta$  (the other value being impossible)  
the general value of  $\theta$  is  $2n\pi \pm \frac{\pi}{3}$ .

#### EXAMPLES—XXVII. (p. 87).

(1)

$$\begin{aligned}
 \sin(A+B).\sin(A-B) &= (\sin A.\cos B + \cos A.\sin B)(\sin A.\cos B - \cos A.\sin B) \\
 &= \sin^2 A.\cos^2 B - \cos^2 A.\sin^2 B \\
 &= \sin^2 A(1 - \sin^2 B) - (1 - \sin^2 A)\sin^2 B \\
 &= \sin^2 A - \sin^2 B.
 \end{aligned}$$

(2)

$$\begin{aligned}
 \sin(a+\beta).\sin(a-\beta) &= \sin a.\cos\beta + \cos a.\sin\beta)(\sin a.\cos\beta - \cos a.\sin\beta) \\
 &= \sin^2 a.\cos^2\beta - \cos^2 a.\sin^2\beta \\
 &= (1 - \cos^2 a)\cos^2\beta - \cos^2 a(1 - \cos^2\beta) \\
 &= \cos^2\beta - \cos^2 a.
 \end{aligned}$$

(3)

$$\begin{aligned}
 \cos(A+B).\cos(A-B) &= (\cos A.\cos B - \sin A.\sin B)(\cos B.\cos B + \sin A.\sin B) \\
 &= \cos^2 A.\cos^2 B - \sin^2 A.\sin^2 B \\
 &= \cos^2 A(1 - \sin^2 B) - (1 - \cos^2 A)\sin^2 B \\
 &= \cos^2 A - \sin^2 B.
 \end{aligned}$$

(4)

$$\begin{aligned}
 \cos(a+\beta).\cos(a-\beta) &= (\cos a.\cos\beta - \sin a.\sin\beta)(\cos a.\cos\beta + \sin a.\sin\beta) \\
 &= \cos^2 a.\cos^2\beta - \sin^2 a.\sin^2\beta \\
 &= (1 - \sin^2 a)\cos^2\beta - \sin^2 a(1 - \cos^2\beta) \\
 &= \cos^2\beta - \sin^2 a.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad 2 \sin(x+y) \cdot \cos(x-y) &= 2(\sin x \cdot \cos y + \cos x \cdot \sin y) \cdot (\cos x \cdot \cos y + \sin x \cdot \sin y) \\
 &= 2\{\sin x \cdot \cos x \cdot \cos^2 y + \sin^2 x \cdot \cos y \cdot \sin y + \cos^2 x \cdot \sin y \cdot \cos y + \\
 &\quad \sin x \cdot \cos x \cdot \sin^2 y\} \\
 &= 2\{\sin x \cdot \cos x (\cos^2 y + \sin^2 y) + \sin y \cdot \cos y (\sin^2 x + \cos^2 x)\} \\
 &= 2\{\sin x \cdot \cos x + \sin y \cdot \cos y\} \\
 &= (\sin x \cdot \cos x + \cos x \cdot \sin x) + (\sin y \cdot \cos y + \cos y \cdot \sin y) \\
 &= \sin(x+x) + \sin(y+y) \\
 &= \sin 2x + \sin 2y
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad 2 \cos(x+y) \cdot \sin(x-y) &= 2(\cos x \cdot \cos y - \sin x \cdot \sin y) \cdot (\sin x \cdot \cos y - \cos x \cdot \sin y) \\
 &= 2\{\sin x \cdot \cos x \cdot \cos^2 y - \sin y \cdot \cos y \cdot \cos^2 x - \sin y \cdot \cos y \cdot \sin^2 x + \\
 &\quad \sin x \cdot \cos x \cdot \sin^2 y\} \\
 &= 2\{\sin x \cdot \cos x \cdot (\cos^2 y + \sin^2 y) - \sin y \cdot \cos y \cdot (\cos^2 x + \sin^2 x)\} \\
 &= 2\{\sin x \cdot \cos x - \sin y \cdot \cos y\} \\
 &= (\sin x \cdot \cos x + \cos x \cdot \sin x) - (\sin y \cdot \cos y + \cos y \cdot \sin y) \\
 &= \sin 2x - \sin 2y.
 \end{aligned}$$

$$\begin{aligned}
 (7) \quad \tan A + \tan B &= \frac{\sin A}{\cos A} + \frac{\sin B}{\cos B} \\
 &= \frac{\sin A \cdot \cos B + \cos A \cdot \sin B}{\cos A \cdot \cos B} \\
 &= \frac{\sin(A+B)}{\cos A \cdot \cos B}.
 \end{aligned}$$

$$\begin{aligned}
 (8) \quad \tan a - \tan \beta &= \frac{\sin a}{\cos a} - \frac{\sin \beta}{\cos \beta} \\
 &= \frac{\sin a \cdot \cos \beta - \cos a \cdot \sin \beta}{\cos a \cdot \cos \beta} \\
 &= \frac{\sin(a-\beta)}{\cos a \cdot \cos \beta}.
 \end{aligned}$$

## EXAMPLES—XXVIII. (p. 88).

$$\begin{aligned}
 (1) \sin 15^\circ &= \sin(45^\circ - 30^\circ) \\
 &= \sin 45^\circ \cdot \cos 30^\circ - \cos 45^\circ \cdot \sin 30^\circ \\
 &= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \cdot \frac{1}{2} \\
 &= \frac{\sqrt{3}}{2\sqrt{2}} - \frac{1}{2\sqrt{2}} = \frac{\sqrt{3}-1}{2\sqrt{2}}.
 \end{aligned}$$

$$\begin{aligned}
 (2) \cos 75^\circ &= \cos(45^\circ + 30^\circ) \\
 &= \cos 45^\circ \cdot \cos 30^\circ - \sin 45^\circ \cdot \sin 30^\circ \\
 &= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \cdot \frac{1}{2} \\
 &= \frac{\sqrt{3}}{2\sqrt{2}} - \frac{1}{2\sqrt{2}} = \frac{\sqrt{3}-1}{2\sqrt{2}}.
 \end{aligned}$$

$$\begin{aligned}
 (3) \tan 75^\circ &= \tan(45^\circ + 30^\circ) \\
 &= \sin(45^\circ + 30^\circ) \div \cos(45^\circ + 30^\circ) \\
 &= \frac{\sqrt{3}+1}{2\sqrt{2}} \div \frac{\sqrt{3}-1}{2\sqrt{2}} \\
 &= \frac{\sqrt{3}+1}{\sqrt{3}-1} = \frac{(\sqrt{3}+1)^2}{(\sqrt{3}-1)(\sqrt{3}+1)} = \frac{4+2\sqrt{3}}{3-1} = 2+\sqrt{3}.
 \end{aligned}$$

$$\begin{aligned}
 (4) \cot 75^\circ &= \cos 75^\circ \div \sin 75^\circ \\
 &= \cos(45^\circ + 30^\circ) \div \sin(45^\circ + 30^\circ) \\
 &= \frac{\sqrt{3}-1}{2\sqrt{2}} \div \frac{\sqrt{3}+1}{2\sqrt{2}} = \frac{\sqrt{3}-1}{\sqrt{3}+1} \\
 &= \frac{(\sqrt{3}-1)^2}{(\sqrt{3}+1)(\sqrt{3}-1)} = \frac{4-2\sqrt{3}}{3-1} = 2-\sqrt{3}.
 \end{aligned}$$

$$(5) \text{ If } \sin \alpha = \frac{1}{3}, \quad \cos \alpha = \frac{\sqrt{8}}{3} = \frac{2\sqrt{2}}{3}.$$

$$\text{If } \sin \beta = \frac{2}{3}, \quad \cos \beta = \frac{\sqrt{5}}{3};$$

$$\therefore \sin(\alpha + \beta) = \frac{1}{3} \cdot \frac{\sqrt{5}}{3} + \frac{2\sqrt{2}}{3} \cdot \frac{2}{3} = \frac{\sqrt{5} + 4\sqrt{2}}{9}.$$

$$(6) \text{ If } \cos \alpha = \frac{3}{4}, \sin \alpha = \frac{\sqrt{7}}{4}.$$

$$\text{If } \cos \beta = \frac{2}{5}, \sin \beta = \frac{\sqrt{21}}{5}.$$

$$\therefore \sin(\alpha - \beta) = \frac{\sqrt{7}}{4} \cdot \frac{2}{5} - \frac{3}{4} \cdot \frac{\sqrt{21}}{5} = \frac{2\sqrt{7} - 3\sqrt{21}}{20}.$$

$$(7) \text{ If } \sin \alpha = \frac{1}{2}, \cos \alpha = \frac{\sqrt{3}}{2}.$$

$$\text{If } \cos \beta = \frac{1}{\sqrt{2}}, \sin \beta = \frac{1}{\sqrt{2}};$$

$$\therefore \cos(\alpha + \beta) = \frac{\sqrt{3}}{2} \cdot \frac{1}{\sqrt{2}} - \frac{1}{2} \cdot \frac{1}{\sqrt{2}} = \frac{\sqrt{3} - 1}{2\sqrt{2}}.$$

$$(8) \text{ If } \cos \alpha = \frac{1}{30}, \sin \alpha = \frac{\sqrt{899}}{30}.$$

$$\text{If } \sin \beta = \frac{1}{2}, \cos \beta = \frac{\sqrt{3}}{2};$$

$$\therefore \cos(\alpha - \beta) = \frac{1}{30} \cdot \frac{\sqrt{3}}{2} + \frac{\sqrt{899}}{30} \cdot \frac{1}{2} = \frac{\sqrt{3} + \sqrt{899}}{60}.$$

# EXAMPLES—XXIX. (p. 88).

$$(1) \cos(90^\circ + A) = \cos 90^\circ \cdot \cos A - \sin 90^\circ \cdot \sin A \\ = 0 \cdot \cos A - 1 \cdot \sin A = -\sin A.$$

$$(2) \sin(180^\circ + A) = \sin 180^\circ \cdot \cos A + \cos 180^\circ \cdot \sin A \\ = 0 \cdot \cos A - 1 \cdot \sin A = -\sin A.$$

$$(3) \cos(\pi + \theta) = \cos \pi \cdot \cos \theta - \sin \pi \cdot \sin \theta \\ = -1 \cdot \cos \theta - 0 \cdot \sin \theta = -\cos \theta.$$

$$(4) \sin\left(\frac{3\pi}{2} + \theta\right) = \sin \frac{3\pi}{2} \cdot \cos \theta + \cos \frac{3\pi}{2} \cdot \sin \theta \\ = -1 \cdot \cos \theta + 0 \cdot \sin \theta = -\cos \theta.$$



$$\begin{aligned}
 (5) \operatorname{cosec}\left(\frac{\pi}{2}+a\right) &= \frac{1}{\sin\left(\frac{\pi}{2}+a\right)} \\
 &= \frac{1}{\sin\frac{\pi}{2} \cdot \cos a + \cos\frac{\pi}{2} \cdot \sin a} \\
 &= \frac{1}{1 \cdot \cos a + 0 \cdot \sin a} = \frac{1}{\cos a} = \sec a.
 \end{aligned}$$

$$(6) \tan(\pi+a) = \frac{\sin(\pi+a)}{\cos(\pi+a)} = \frac{0 \cdot \cos a - 1 \cdot \sin a}{-1 \cdot \cos a - 0 \cdot \sin a} = \frac{-\sin a}{-\cos a} = \tan a.$$

$$\begin{aligned}
 (7) \sin(2\pi-\theta) &= \sin 2\pi \cdot \cos \theta - \cos 2\pi \cdot \sin \theta \\
 &= 0 \cdot \cos \theta - 1 \cdot \sin \theta = -\sin \theta.
 \end{aligned}$$

$$(8) \tan(2\pi-\theta) = \frac{\sin(2\pi-\theta)}{\cos(2\pi-\theta)} = \frac{0 \cdot \cos \theta - 1 \cdot \sin \theta}{1 \cdot \cos \theta + 0 \cdot \sin \theta} = \frac{-\sin \theta}{\cos \theta} = -\tan \theta.$$

$$(9) \sec(180^\circ-\theta) = \frac{1}{\cos(180^\circ-\theta)} = \frac{1}{-1 \cdot \cos \theta + 0 \cdot \sin \theta} = -\frac{1}{\cos \theta} = -\sec \theta.$$

$$(10) \operatorname{cosec}(\pi-\theta) = \frac{1}{\sin(\pi-\theta)} = \frac{1}{0 \cdot \cos \theta - (-1 \cdot \sin \theta)} = \frac{1}{\sin \theta} = \operatorname{cosec} \theta.$$

## EXAMPLES—XXX. (p. 89).

$$\begin{aligned}
 (1) \quad &\sin \theta - \cos \theta = 0. \\
 &\sin \theta \cdot \frac{1}{\sqrt{2}} - \cos \theta \cdot \frac{1}{\sqrt{2}} = 0 \\
 &\sin \theta \cdot \cos 45^\circ - \cos \theta \cdot \sin 45^\circ = 0; \\
 &\therefore \sin(\theta - 45^\circ) = 0, \therefore \theta - 45^\circ = 0^\circ, \text{ or } \theta = 45^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (2) \quad &\sin \theta + \cos \theta = 1 \\
 &\sin \theta \cdot \frac{1}{\sqrt{2}} + \cos \theta \cdot \frac{1}{\sqrt{2}} = \frac{1}{\sqrt{2}}; \\
 &\sin \theta \cdot \cos 45^\circ + \cos \theta \cdot \sin 45^\circ = \frac{1}{\sqrt{2}}; \\
 &\therefore \sin(\theta + 45^\circ) = \sin 45^\circ; \\
 &\therefore \theta + 45^\circ = 45^\circ, \text{ or, } \theta = 0^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (3) \quad \sin\theta - \cos\theta &= \sqrt{\frac{3}{2}} \\
 \sin\theta \cdot \frac{1}{\sqrt{2}} - \cos\theta \cdot \frac{1}{\sqrt{2}} &= \frac{\sqrt{3}}{2}; \\
 \therefore \sin(\theta - 45^\circ) &= \sin 60^\circ; \\
 \therefore \theta - 45^\circ &= 60^\circ, \text{ or, } \theta = 105^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (4) \quad \sin\theta + \cos\theta &= \frac{\sqrt{3} + 1}{2} \\
 \sin\theta \cdot \frac{1}{\sqrt{2}} + \cos\theta \cdot \frac{1}{\sqrt{2}} &= \frac{\sqrt{3} + 1}{2\sqrt{2}} \\
 \sin(\theta + 45^\circ) &= \sin 75^\circ, \text{ whence } \theta = 30^\circ, \text{ or,} \\
 \cos(\theta - 45^\circ) &= \cos 15^\circ, \text{ whence } \theta = 60^\circ, \text{ or,} \\
 \cos(45^\circ - \theta) &= \cos 15^\circ, \text{ whence } \theta = -30^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad \sin\theta + \cos\theta &= \sqrt{2} \\
 \sin\theta \cdot \frac{1}{\sqrt{2}} + \cos\theta \cdot \frac{1}{\sqrt{2}} &= 1 \\
 \sin(\theta + 45^\circ) &= \sin 90^\circ, \text{ or, } \theta = 45^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad \sin\theta - \cos\theta &= \frac{\sqrt{3} - 1}{2} \\
 \sin\theta \cdot \frac{1}{\sqrt{2}} - \cos\theta \cdot \frac{1}{\sqrt{2}} &= \frac{\sqrt{3} - 1}{2\sqrt{2}} \\
 \sin(\theta - 45^\circ) &= \sin 15^\circ, \text{ whence } \theta = 60^\circ.
 \end{aligned}$$

EXAMPLES—XXXI. (p. 92).

$$(1) \sin 6A + \sin 4A = 2 \sin \frac{6A + 4A}{2} \cdot \cos \frac{6A - 4A}{2} = 2 \sin 5A \cdot \cos A.$$

$$(2) \sin 5A - \sin 3A = 2 \cos \frac{5A + 3A}{2} \cdot \sin \frac{5A - 3A}{2} = 2 \cos 4A \cdot \sin A.$$

$$(3) \cos 7\theta + \cos 9\theta = 2 \cos \frac{7\theta + 9\theta}{2} \cdot \cos \frac{9\theta - 7\theta}{2} = 2 \cos 8\theta \cdot \cos \theta.$$

$$(4) \cos \theta - \cos 5\theta = 2 \sin \frac{\theta + 5\theta}{2} \cdot \sin \frac{5\theta - \theta}{2} = 2 \sin 3\theta \cdot \sin 2\theta.$$

$$(5) \sin a + \sin 4a = 2 \sin \frac{a+4a}{2} \cdot \cos \frac{4a-a}{2} = 2 \sin \frac{5a}{2} \cdot \cos \frac{3a}{2}.$$

$$(6) \cos 5a - \cos 8a = 2 \sin \frac{5a+8a}{2} \cdot \sin \frac{8a-5a}{2} = 2 \sin \frac{13a}{2} \cdot \sin \frac{3a}{2}.$$

$$(7) 2 \sin 5\theta \cdot \cos 7\theta = \sin(5\theta + 7\theta) - \sin(7\theta - 5\theta) = \sin 12\theta - \sin 2\theta.$$

$$(8) 2 \sin 3\theta \cdot \sin 5\theta = \cos(5\theta - 3\theta) - \cos(5\theta + 3\theta) = \cos 2\theta - \cos 8\theta.$$

$$(9) 2 \cos a \cdot \cos 4a = \cos(a + 4a) + \cos(4a - a) = \cos 5a + \cos 3a.$$

$$(10) 2 \cos a \cdot \sin 2a = \sin(a + 2a) + \sin(2a - a) = \sin 3a + \sin a.$$

$$(11) \frac{\sin A + \sin B}{\cos A + \cos B} = \frac{2 \sin \frac{A+B}{2} \cdot \cos \frac{A-B}{2}}{2 \cos \frac{A+B}{2} \cdot \cos \frac{A-B}{2}} = \frac{\sin \frac{A+B}{2}}{\cos \frac{A+B}{2}} = \tan \frac{A+B}{2}.$$

$$(12) \frac{\cos A - \cos 3A}{\sin 3A - \sin A} = \frac{2 \sin 2A \cdot \sin A}{2 \cos 2A \cdot \sin A} = \frac{\sin 2A}{\cos 2A} = \tan 2A.$$

$$(13) \frac{\sin 2A + \sin A}{\cos 2A + \cos A} = \frac{2 \sin \frac{3A}{2} \cdot \cos \frac{A}{2}}{2 \cos \frac{3A}{2} \cdot \cos \frac{A}{2}} = \frac{\sin \frac{3A}{2}}{\cos \frac{3A}{2}} = \tan \frac{3A}{2}.$$

$$(14) \cos(30^\circ - \theta) - \cos(30^\circ + \theta) = 2 \sin 30^\circ \cdot \sin \theta = 2 \times \frac{1}{2} \cdot \sin \theta = \sin \theta.$$

$$(15) \cos\left(\frac{\pi}{3} + \theta\right) + \cos\left(\frac{\pi}{3} - \theta\right) = 2 \cos \frac{\pi}{3} \cdot \cos \theta = 2 \times \frac{1}{2} \cdot \cos \theta = \cos \theta.$$

$$(16) \sin\left(\frac{\pi}{3} + a\right) - \sin\left(\frac{\pi}{3} - a\right) = 2 \cos \frac{\pi}{3} \cdot \sin a = 2 \times \frac{1}{2} \cdot \sin a = \sin a.$$

$$(17) \frac{\sin a - \sin \beta}{\cos \beta - \cos a} = \frac{2 \cos \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2}}{2 \sin \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2}} = \frac{\cos \frac{a+\beta}{2}}{\sin \frac{a+\beta}{2}} = \cot \frac{a+\beta}{2}.$$

$$(18) \frac{\sin a - \sin \beta}{\cos \beta + \cos a} = \frac{2 \cos \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2}}{2 \cos \frac{a+\beta}{2} \cdot \cos \frac{a-\beta}{2}} = \frac{\sin \frac{a-\beta}{2}}{\cos \frac{a-\beta}{2}} = \tan \frac{a-\beta}{2}.$$

$$(19) \frac{\sin 5\theta + \sin 3\theta}{\cos 3\theta - \cos 5\theta} = \frac{2 \sin 4\theta \cdot \cos \theta}{2 \sin 4\theta \cdot \sin \theta} = \frac{\cos \theta}{\sin \theta} = \cot \theta.$$

$$(20) \frac{\cos a + \cos \beta}{\cos \beta - \cos a} = \frac{2 \cos \frac{a+\beta}{2} \cdot \cos \frac{a-\beta}{2}}{2 \sin \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2}} = \frac{\cos \frac{a+\beta}{2}}{\sin \frac{a+\beta}{2}} \div \frac{\sin \frac{a-\beta}{2}}{\cos \frac{a-\beta}{2}} = \frac{\cot \frac{a+\beta}{2}}{\tan \frac{a-\beta}{2}}.$$

EXAMPLES—XXXII. (p. 93).

$$(1) \sin a - \cos \beta = \sin a - \sin \left( \frac{\pi}{2} - \beta \right) = 2 \cos \frac{1}{2} \left( a + \frac{\pi}{2} - \beta \right) \cdot \sin \frac{1}{2} \left( a - \frac{\pi}{2} + \beta \right).$$

$$(2) \sin \left( \frac{\pi}{2} + a \right) + \cos \left( \frac{\pi}{2} - a \right) = \sin \left( \frac{\pi}{2} + a \right) + \sin a = 2 \sin \left( \frac{\pi}{4} + a \right) \cdot \cos \frac{\pi}{4}.$$

$$(3) \sin a + \cos a = \sin a + \sin \left( \frac{\pi}{2} - a \right) = 2 \sin \frac{\pi}{4} \cdot \cos \left( a - \frac{\pi}{4} \right).$$

$$(4) \sin a - \cos a = \sin a - \sin \left( \frac{\pi}{2} - a \right) = 2 \cos \frac{\pi}{4} \cdot \sin \left( a - \frac{\pi}{4} \right).$$

$$(5) \sin 30^\circ + \cos 80^\circ = \sin 30^\circ + \sin 10^\circ = 2 \sin 20^\circ \cdot \cos 10^\circ.$$

$$(6) \sin 20^\circ - \cos 80^\circ = \sin 20^\circ - \sin 10^\circ = 2 \cos 15^\circ \cdot \sin 5^\circ.$$

$$(7) \sin \frac{\pi}{4} + \cos \frac{\pi}{6} = \sin \frac{\pi}{4} + \sin \frac{\pi}{3} = 2 \sin \frac{7\pi}{24} \cdot \cos \frac{\pi}{24}.$$

$$(8) \sin \frac{\pi}{3} - \cos \frac{\pi}{5} = \sin \frac{\pi}{3} - \sin \frac{3\pi}{10} = 2 \cos \frac{19\pi}{60} \cdot \sin \frac{\pi}{60}.$$

## EXAMPLES—XXXIII. (p. 96).

$$\begin{aligned}
 (1) \quad \frac{\tan \alpha + \tan \beta}{\cot \alpha + \cot \beta} &= \frac{\frac{\sin \alpha}{\cos \alpha} + \frac{\sin \beta}{\cos \beta}}{\frac{\cos \alpha}{\sin \alpha} + \frac{\cos \beta}{\sin \beta}} = \frac{\frac{\sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta}}{\frac{\cos \alpha \cdot \sin \beta + \sin \alpha \cdot \cos \beta}{\sin \alpha \cdot \sin \beta}} \\
 &= \frac{\sin \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta} = \tan \alpha \cdot \tan \beta.
 \end{aligned}$$

$$\begin{aligned}
 (2) \quad \frac{\tan \alpha + \tan \beta}{\cot \alpha - \tan \beta} &= \frac{\frac{\sin \alpha}{\cos \alpha} + \frac{\sin \beta}{\cos \beta}}{\frac{\cos \alpha}{\sin \alpha} - \frac{\sin \beta}{\cos \beta}} = \frac{\frac{\sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta}}{\frac{\cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta}{\sin \alpha \cdot \cos \beta}} \\
 &= \frac{\sin(\alpha + \beta) \cdot \sin \alpha \cdot \cos \beta}{\cos(\alpha + \beta) \cdot \cos \alpha \cdot \cos \beta} = \tan(\alpha + \beta) \cdot \tan \alpha.
 \end{aligned}$$

$$\begin{aligned}
 (3) \quad \frac{\tan \alpha - \tan \beta}{\cot \alpha + \tan \beta} &= \frac{\frac{\sin \alpha}{\cos \alpha} - \frac{\sin \beta}{\cos \beta}}{\frac{\cos \alpha}{\sin \alpha} + \frac{\sin \beta}{\cos \beta}} = \frac{\frac{\sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta}}{\frac{\cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta}{\sin \alpha \cdot \cos \beta}} \\
 &= \frac{\sin(\alpha - \beta) \cdot \sin \alpha \cdot \cos \beta}{\cos(\alpha - \beta) \cdot \cos \alpha \cdot \cos \beta} = \tan(\alpha - \beta) \cdot \tan \alpha.
 \end{aligned}$$

$$\begin{aligned}
 (4) \quad \tan \frac{\phi + \psi}{2} + \tan \frac{\phi - \psi}{2} &= \frac{\sin \frac{\phi + \psi}{2}}{\cos \frac{\phi + \psi}{2}} + \frac{\sin \frac{\phi - \psi}{2}}{\cos \frac{\phi - \psi}{2}} \\
 &= \frac{\sin \frac{\phi + \psi}{2} \cdot \cos \frac{\phi - \psi}{2} + \cos \frac{\phi + \psi}{2} \cdot \sin \frac{\phi - \psi}{2}}{\cos \frac{\phi + \psi}{2} \cdot \cos \frac{\phi - \psi}{2}} \\
 &= \frac{\sin\left(\frac{\phi + \psi}{2} + \frac{\phi - \psi}{2}\right)}{\frac{1}{2}(\cos \phi + \cos \psi)} = \frac{\sin \phi}{\frac{1}{2}(\cos \phi + \cos \psi)} = \frac{2 \sin \phi}{\cos \phi + \cos \psi}.
 \end{aligned}$$

$$(5) \sin \phi = \sin(\psi + (\phi - \psi)) = \sin \psi \cdot \cos(\phi - \psi) + \cos \psi \cdot \sin(\phi - \psi).$$

$$(6) \cos \phi = \cos((\phi + \psi) - \psi) = \cos(\phi + \psi) \cdot \cos \psi + \sin(\phi + \psi) \cdot \sin \psi.$$

$$\begin{aligned} (7) \quad & (\cos \alpha + \cos \beta)(1 - \cos(\alpha + \beta)) = (\cos \alpha + \cos \beta)(1 - \cos \alpha \cos \beta + \sin \alpha \sin \beta) \\ & = \cos \alpha + \cos \beta - \cos^2 \alpha \cos \beta - \cos \alpha \cos^2 \beta + \sin \alpha \sin \beta \cos \alpha + \sin \alpha \sin \beta \cos \beta \\ & = \cos \alpha (1 - \cos^2 \beta) + \cos \beta (1 - \cos^2 \alpha) + \sin \alpha \sin \beta \cos \alpha + \sin \alpha \sin \beta \cos \beta \\ & = \cos \alpha \sin^2 \beta + \cos \beta \sin^2 \alpha + \sin \alpha \sin \beta \cos \alpha + \sin \alpha \sin \beta \cos \beta \\ & = \sin \beta (\cos \alpha \sin \beta) + \sin \alpha (\cos \beta \sin \alpha) + \sin \alpha (\sin \beta \cos \alpha) + \sin \beta (\sin \alpha \cos \beta) \\ & = \sin \beta (\cos \alpha \sin \beta + \sin \alpha \cos \beta) + \sin \alpha (\cos \beta \sin \alpha + \sin \beta \cos \alpha) \\ & = \sin \beta \sin(\alpha + \beta) + \sin \alpha \sin(\alpha + \beta) \\ & = (\sin \alpha + \sin \beta) \sin(\alpha + \beta). \end{aligned}$$

(8)

$$\begin{aligned} \frac{\sin(\alpha + \beta)}{\sin \alpha + \sin \beta} &= \frac{\sin\left(\frac{\alpha + \beta}{2} + \frac{\alpha + \beta}{2}\right)}{\sin \alpha + \sin \beta} = \frac{\sin \frac{\alpha + \beta}{2} \cdot \cos \frac{\alpha + \beta}{2} + \cos \frac{\alpha + \beta}{2} \cdot \sin \frac{\alpha + \beta}{2}}{2 \sin \frac{\alpha + \beta}{2} \cdot \cos \frac{\alpha - \beta}{2}} \\ &= \frac{2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha + \beta}{2}}{2 \cos \frac{\alpha - \beta}{2} \cos \frac{\alpha - \beta}{2}} = \frac{\cos \frac{\alpha + \beta}{2}}{\cos \frac{\alpha - \beta}{2}}. \end{aligned}$$

(9)

$$\begin{aligned} \frac{\sin(\alpha + \beta)}{\sin \alpha - \sin \beta} &= \frac{\sin\left(\frac{\alpha + \beta}{2} + \frac{\alpha + \beta}{2}\right)}{\sin \alpha - \sin \beta} = \frac{\sin \frac{\alpha + \beta}{2} \cdot \cos \frac{\alpha + \beta}{2} + \cos \frac{\alpha + \beta}{2} \cdot \sin \frac{\alpha + \beta}{2}}{2 \cos \frac{\alpha + \beta}{2} \cdot \sin \frac{\alpha - \beta}{2}} \\ &= \frac{2 \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha + \beta}{2}}{2 \sin \frac{\alpha - \beta}{2} \sin \frac{\alpha - \beta}{2}} = \frac{\sin \frac{\alpha + \beta}{2}}{\sin \frac{\alpha - \beta}{2}}. \end{aligned}$$



$$\begin{aligned}
 (10) \quad \cot \frac{a+\beta}{2} + \cot \frac{a-\beta}{2} &= \frac{\cos \frac{a+\beta}{2}}{\sin \frac{a+\beta}{2}} + \frac{\cos \frac{a-\beta}{2}}{\sin \frac{a-\beta}{2}} \\
 &= \frac{\cos \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2} + \cos \frac{a-\beta}{2} \cdot \sin \frac{a+\beta}{2}}{\sin \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2}} \\
 &= \frac{\sin \left( \frac{a+\beta}{2} + \frac{a-\beta}{2} \right)}{\frac{1}{2}(\cos \beta - \cos a)} = \frac{\sin a}{\frac{1}{2}(\cos \beta - \cos a)} = \frac{2 \sin a}{\cos \beta - \cos a}.
 \end{aligned}$$

$$\begin{aligned}
 (11) \quad \tan \frac{a+\beta}{2} - \tan \frac{a-\beta}{2} &= \frac{\sin \frac{a+\beta}{2}}{\cos \frac{a+\beta}{2}} - \frac{\sin \frac{a-\beta}{2}}{\cos \frac{a-\beta}{2}} \\
 &= \frac{\sin \frac{a+\beta}{2} \cdot \cos \frac{a-\beta}{2} - \cos \frac{a+\beta}{2} \cdot \sin \frac{a-\beta}{2}}{\cos \frac{a+\beta}{2} \cdot \cos \frac{a-\beta}{2}} \\
 &= \frac{\sin \left( \frac{a+\beta}{2} - \frac{a-\beta}{2} \right)}{\frac{1}{2}(\cos a + \cos \beta)} = \frac{\sin \beta}{\frac{1}{2}(\cos a + \cos \beta)} = \frac{2 \sin \beta}{\cos a + \cos \beta}.
 \end{aligned}$$

$$(12) \quad \frac{\cos a - \cos \beta}{\sin a + \sin \beta} = \frac{2 \sin \frac{\beta+a}{2} \cdot \sin \frac{\beta-a}{2}}{2 \sin \frac{\beta+a}{2} \cdot \cos \frac{\beta-a}{2}} = \tan \frac{\beta-a}{2}.$$

$$(13) \quad \cot \beta - \tan a = \frac{\cos \beta}{\sin \beta} - \frac{\sin a}{\cos a} = \frac{\cos a \cdot \cos \beta - \sin a \cdot \sin \beta}{\cos a \cdot \sin \beta} = \frac{\cos(a+\beta)}{\cos a \cdot \sin \beta}.$$

$$(14) \quad \cot \theta + \tan \phi = \frac{\cos \theta}{\sin \theta} + \frac{\sin \phi}{\cos \phi} = \frac{\cos \theta \cdot \cos \phi + \sin \theta \cdot \sin \phi}{\sin \theta \cdot \cos \phi} = \frac{\cos(\phi - \theta)}{\sin \theta \cdot \cos \phi}.$$



$$\begin{aligned}
 (15) \tan^2 \alpha - \tan^2 \beta &= \frac{\sin^2 \alpha}{\cos^2 \alpha} - \frac{\sin^2 \beta}{\cos^2 \beta} = \frac{\sin^2 \alpha \cdot \cos^2 \beta - \cos^2 \alpha \cdot \sin^2 \beta}{\cos^2 \alpha \cdot \cos^2 \beta} \\
 &= \frac{(\sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta)(\sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta)}{\cos^2 \alpha \cdot \cos^2 \beta} \\
 &= \frac{\sin(\alpha + \beta) \cdot \sin(\alpha - \beta)}{\cos^2 \alpha \cdot \cos^2 \beta}.
 \end{aligned}$$

$$(16) 1 + \tan \alpha \cdot \tan \beta = 1 + \frac{\sin \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta} = \frac{\cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta} = \frac{\cos(\alpha - \beta)}{\cos \alpha \cdot \cos \beta}.$$

$$\begin{aligned}
 (17) 1 - \tan \alpha \cdot \tan \beta &= 1 - \frac{\sin \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta} = \frac{\cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta}{\cos \alpha \cdot \cos \beta} \\
 &= \frac{\cos(\alpha + \beta)}{\cos \alpha \cdot \cos \beta}.
 \end{aligned}$$

$$\begin{aligned}
 (18) \cot \alpha + \tan \beta &= \frac{\cos \alpha}{\sin \alpha} + \frac{\sin \beta}{\cos \beta} = \frac{\cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta}{\sin \alpha \cdot \cos \beta} \\
 \tan \alpha + \cot \beta &= \frac{\sin \alpha}{\cos \alpha} + \frac{\cos \beta}{\sin \beta} = \frac{\sin \alpha \cdot \sin \beta + \cos \alpha \cdot \cos \beta}{\cos \alpha \cdot \sin \beta} \\
 &= \frac{\cos \alpha \cdot \sin \beta}{\sin \alpha \cdot \cos \beta} = \cot \alpha \cdot \tan \beta.
 \end{aligned}$$

$$\begin{aligned}
 (19) \frac{\tan^2 x - \tan^2 y}{1 - \tan^2 x \cdot \tan^2 y} &= \frac{\frac{\sin^2 x}{\cos^2 x} - \frac{\sin^2 y}{\cos^2 y}}{1 - \frac{\sin^2 x \cdot \sin^2 y}{\cos^2 x \cdot \cos^2 y}} = \frac{\sin^2 x \cdot \cos^2 y - \cos^2 x \cdot \sin^2 y}{\cos^2 x \cdot \cos^2 y - \sin^2 x \cdot \sin^2 y} \\
 &= \frac{(\sin x \cdot \cos y + \cos x \cdot \sin y)(\sin x \cdot \cos y - \cos x \cdot \sin y)}{(\cos x \cdot \cos y + \sin x \cdot \sin y)(\cos x \cdot \cos y - \sin x \cdot \sin y)} \\
 &= \frac{\sin(x + y) \cdot \sin(x - y)}{\cos(x - y) \cdot \cos(x + y)} = \tan(x + y) \cdot \tan(x - y).
 \end{aligned}$$

$$\begin{aligned}
 (20) \cot(\theta + 45^\circ) &= \frac{\cos(\theta + 45^\circ)}{\sin(\theta + 45^\circ)} = \frac{\cos \theta \cdot \frac{1}{\sqrt{2}} - \sin \theta \cdot \frac{1}{\sqrt{2}}}{\sin \theta \cdot \frac{1}{\sqrt{2}} + \cos \theta \cdot \frac{1}{\sqrt{2}}} = \frac{\cos \theta - \sin \theta}{\sin \theta + \cos \theta} \\
 &= \frac{\frac{\cos \theta}{\sin \theta} - 1}{1 + \frac{\cos \theta}{\sin \theta}} = \frac{\cot \theta - 1}{\cot \theta + 1}.
 \end{aligned}$$

$$(21) \sin \theta + \cos \theta = \sqrt{2} \cdot \left( \sin \theta \cdot \frac{1}{\sqrt{2}} + \cos \theta \cdot \frac{1}{\sqrt{2}} \right) = \sqrt{2} \cdot \sin(45^\circ + \theta).$$

$$(22) \cos \theta - \sin \theta = \sqrt{2} \left( \cos \theta \cdot \frac{1}{\sqrt{2}} - \sin \theta \cdot \frac{1}{\sqrt{2}} \right) = \sqrt{2} \cdot \sin \left( \frac{\pi}{4} - \theta \right).$$

$$(23) \frac{\tan \alpha - \tan \beta}{\tan \alpha + \tan \beta} = \frac{\frac{\sin \alpha}{\cos \alpha} - \frac{\sin \beta}{\cos \beta}}{\frac{\sin \alpha}{\cos \alpha} + \frac{\sin \beta}{\cos \beta}} = \frac{\sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta}{\sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta} = \frac{\sin(\alpha - \beta)}{\sin(\alpha + \beta)}.$$

$$(24) \frac{\cot x - \cot y}{\cot x + \cot y} = \frac{\frac{\cos x}{\sin x} - \frac{\cos y}{\sin y}}{\frac{\cos x}{\sin x} + \frac{\cos y}{\sin y}} = \frac{\cos x \cdot \sin y - \sin x \cdot \cos y}{\cos x \cdot \sin y + \sin x \cdot \cos y} = \frac{\sin(y - x)}{\sin(y + x)}.$$

$$(25) \begin{aligned} \cos(A - B) + \sin(A + B) &= \cos(A - B) + \cos(90^\circ - A - B) \\ &= 2 \cos(45^\circ - B) \cdot \cos(45^\circ - A) \\ &= 2 \cos(B - 45^\circ) \cdot \sin(45^\circ + A). \end{aligned}$$

$$(26) \begin{aligned} \cos(A - B) - \sin(A + B) &= \sin(90^\circ - A - B) - \sin(A + B) \\ &= 2 \cos(45^\circ + B) \cdot \sin(45^\circ - A). \end{aligned}$$

$$(27) \begin{aligned} \cos(A + B) + \sin(A - B) &= \cos(A + B) + \cos(90^\circ - A + B) \\ &= 2 \cos(45^\circ + B) \cdot \cos(45^\circ - A) \\ &= 2 \cos(45^\circ + B) \cdot \sin(45^\circ + A). \end{aligned}$$

$$(28) \begin{aligned} \cos(A + B) - \sin(A - B) &= \sin(90^\circ - A - B) - \sin(A - B) \\ &= 2 \cos(45^\circ - B) \cdot \sin(45^\circ - A). \end{aligned}$$

$$(29) \frac{\cos \alpha + \cos \beta}{\cos \alpha - \cos \beta} = \frac{\cos \alpha + \cos \beta}{-(\cos \beta - \cos \alpha)}$$

$$= - \frac{2 \cos \frac{\alpha + \beta}{2} \cdot \cos \frac{\alpha - \beta}{2}}{2 \sin \frac{\alpha + \beta}{2} \cdot \sin \frac{\alpha - \beta}{2}} = - \frac{\cot \frac{\alpha + \beta}{2}}{\tan \frac{\alpha - \beta}{2}}.$$

$$(30) \begin{aligned} \sec 72^\circ - \sec 36^\circ &= \frac{1}{\cos 72^\circ} - \frac{1}{\cos 36^\circ} = \frac{\cos 36^\circ - \cos 72^\circ}{\cos 72^\circ \cdot \cos 36^\circ} \\ &= \frac{2 \sin 54^\circ \cdot \sin 18^\circ}{\sin 18^\circ \cdot \sin 54^\circ} = 2 = \sec 60^\circ. \end{aligned}$$

$$\begin{aligned}
 (31) \quad & (\sin 81^\circ + \sin 9^\circ)(\sin 81^\circ - \sin 9^\circ) \\
 &= (2 \sin 45^\circ \cdot \cos 36^\circ) \cdot (2 \cos 45^\circ \cdot \sin 36^\circ) \\
 &= 2 \cdot \frac{1}{\sqrt{2}} \cdot \sin 54^\circ \cdot 2 \cdot \frac{1}{\sqrt{2}} \cdot \cos 54^\circ \\
 &= 2 \sin 54^\circ \cdot \cos 54^\circ \\
 &= \sin 108^\circ.
 \end{aligned}$$

$$(32) \quad \frac{\cos 3^\circ - \cos 33^\circ}{\sin 3^\circ + \sin 33^\circ} = \frac{2 \sin 18^\circ \cdot \sin 15^\circ}{2 \sin 18^\circ \cdot \cos 15^\circ} = \tan 15^\circ.$$

$$(33) \quad \frac{\sin 33^\circ + \sin 3^\circ}{\cos 33^\circ + \cos 3^\circ} = \frac{2 \sin 18^\circ \cdot \cos 15^\circ}{2 \cos 18^\circ \cdot \cos 15^\circ} = \tan 18^\circ.$$

$$(34) \quad \frac{\cos 9^\circ + \sin 9^\circ}{\cos 9^\circ - \sin 9^\circ} = \frac{\sin 81^\circ + \sin 9^\circ}{\sin 81^\circ - \sin 9^\circ} = \frac{2 \sin 45^\circ \cdot \cos 36^\circ}{2 \cos 45^\circ \cdot \sin 36^\circ} = \cot 36^\circ = \tan 54^\circ.$$

$$(35) \quad \frac{\cos 27^\circ - \sin 27^\circ}{\cos 27^\circ + \sin 27^\circ} = \frac{\sin 63^\circ - \sin 27^\circ}{\sin 63^\circ + \sin 27^\circ} = \frac{2 \cos 45^\circ \cdot \sin 18^\circ}{2 \sin 45^\circ \cdot \cos 18^\circ} = \tan 18^\circ.$$

$$\begin{aligned}
 (36) \quad & \tan 50^\circ + \cot 50^\circ = \tan 50^\circ + \tan 40^\circ \\
 &= \frac{\sin 50^\circ \cdot \cos 40^\circ + \cos 50^\circ \cdot \sin 40^\circ}{\cos 50^\circ \cdot \cos 40^\circ} = \frac{\sin 90^\circ}{\frac{1}{2} \{\cos 90^\circ + \cos 10^\circ\}} \\
 &= \frac{2 \sin 90^\circ}{\cos 10^\circ} = \frac{2}{\cos 10^\circ} = 2 \sec 10^\circ.
 \end{aligned}$$

## EXAMPLES—XXXIV. (p. 100).

$$(1) \quad \frac{2 \cot A}{1 + \cot^2 A} = \frac{2 \cot A}{\operatorname{cosec}^2 A} = \frac{2 \cos A}{\sin A} \cdot \sin^2 A = 2 \cos A \cdot \sin A = \sin 2A.$$

$$\begin{aligned}
 (2) \quad & \frac{\sin 2A}{1 + \cos 2A} \cdot \frac{\cos A}{1 + \cos A} = \frac{2 \sin A \cdot \cos A}{2 \cos^2 A} \cdot \frac{\cos A}{1 + \cos A} = \frac{\sin A}{1 + \cos A} \\
 &= \frac{2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}}{2 \cos^2 \frac{A}{2}} = \tan \frac{A}{2}.
 \end{aligned}$$

$$(3) \quad \operatorname{cosec} A + \cot A = \frac{1}{\sin A} + \frac{\cos A}{\sin A} = \frac{1 + \cos A}{\sin A} = \frac{2 \cos^2 \frac{A}{2}}{2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}} = \cot \frac{A}{2}.$$

(4)

$$\tan\theta + \cot\theta = \frac{\sin\theta}{\cos\theta} + \frac{\cos\theta}{\sin\theta} = \frac{\sin^2\theta + \cos^2\theta}{\sin\theta \cdot \cos\theta} = \frac{1}{\sin\theta \cdot \cos\theta} = \frac{2}{2\sin\theta \cdot \cos\theta} = \frac{2}{\sin 2\theta}.$$

$$(5) \frac{2 \tan\theta}{1 + \tan^2\theta} = \frac{2 \frac{\sin\theta}{\cos\theta}}{\sec^2\theta} = \frac{2 \sin\theta}{\cos\theta} \cdot \cos^2\theta = 2 \sin\theta \cdot \cos\theta = \sin 2\theta.$$

$$(6) 2 \operatorname{cosec} 2A = \frac{2}{\sin 2A} = \frac{2}{2 \sin A \cdot \cos A} = \operatorname{cosec} A \cdot \sec A.$$

$$(7) \frac{1 - \tan^2\theta}{1 + \tan^2\theta} = \frac{1 - \frac{\sin^2\theta}{\cos^2\theta}}{1 + \frac{\sin^2\theta}{\cos^2\theta}} = \frac{\cos^2\theta - \sin^2\theta}{\cos^2\theta + \sin^2\theta} = \cos^2\theta - \sin^2\theta = \cos 2\theta.$$

$$(8) \frac{2 \sec 2\theta}{1 + \sec 2\theta} = \frac{\frac{2}{\cos 2\theta}}{1 + \frac{1}{\cos 2\theta}} = \frac{2}{\cos 2\theta + 1} = \frac{2}{2 \cos^2\theta} = \sec^2\theta.$$

$$(9) \frac{1 - \tan A}{1 + \tan A} = \frac{\cos A - \sin A}{\cos A + \sin A} = \frac{\cos^2 A - \sin^2 A}{(\cos A + \sin A)^2} = \frac{1 - 2 \sin^2 A}{1 + \sin 2A}.$$

$$(10) \cot\theta - 2 \cot 2\theta = \frac{\cos\theta}{\sin\theta} - \frac{2 \cos 2\theta}{\sin 2\theta} = \frac{\cos\theta}{\sin\theta} - \frac{\cos 2\theta}{\sin\theta \cdot \cos\theta} \\ = \frac{\cos^2\theta - \cos 2\theta}{\sin\theta \cdot \cos\theta} = \frac{\cos^2\theta - 2 \cos^2\theta + 1}{\sin\theta \cos\theta} = \frac{\sin^2\theta}{\sin\theta \cdot \cos\theta} = \tan\theta.$$

$$(11) \frac{1 - \cos a}{\sin a} = \frac{2 \sin^2 \frac{a}{2}}{2 \sin \frac{a}{2} \cdot \cos \frac{a}{2}} = \frac{\sin \frac{a}{2}}{\cos \frac{a}{2}} = \tan \frac{a}{2}.$$

$$(12) \frac{2\sqrt{\operatorname{cosec}^2\phi - 1}}{\operatorname{cosec}^2\phi} = \frac{2 \cdot \cot\phi}{\operatorname{cosec}^2\phi} = \frac{2 \cdot \cos\phi \cdot \sin^2\phi}{\sin\phi} = 2 \sin\phi \cdot \cos\phi = \sin 2\phi.$$

$$(13) \frac{2 - \sec^2\phi}{\sec^2\phi} = 2 \cos^2\phi - 1 = \cos 2\phi.$$



$$(14) \frac{2 \cot \phi}{\cot^2 \phi - 1} = \frac{2 \cos \phi \cdot \sin \phi}{\cos^2 \phi - \sin^2 \phi} = \frac{\sin 2\phi}{\cos 2\phi} = \tan 2\phi.$$

$$(15) \sqrt{\left(\frac{\sec 2a - 1}{2 \sec 2a}\right)} = \sqrt{\left(\frac{1 - \cos 2a}{2}\right)} = \sqrt{\left(\frac{1 - 1 + 2 \sin^2 a}{2}\right)} = \sin a.$$

$$(16) \sqrt{\left(\frac{\sec 2a + 1}{2 \sec 2a}\right)} = \sqrt{\left(\frac{1 + \cos 2a}{2}\right)} = \sqrt{\left(\frac{1 + 2 \cos^2 a - 1}{2}\right)} = \cos a.$$

$$(17) \operatorname{cosec} 2a - \cot 2a = \frac{1 - \cos 2a}{\sin 2a} = \frac{2 \sin^2 a}{2 \sin a \cdot \cos a} = \tan a.$$

$$(18) \operatorname{cosec} 2\beta + \cot 2\beta = \frac{1 + \cos 2\beta}{\sin 2\beta} = \frac{2 \cos^2 \beta}{2 \sin \beta \cdot \cos \beta} = \cot \beta.$$

$$(19) \tan(45^\circ + A) = \frac{\tan 45^\circ + \tan A}{1 - \tan 45^\circ \cdot \tan A} = \frac{1 + \tan A}{1 - \tan A} = \frac{\cos A + \sin A}{\cos A - \sin A} \\ = \frac{\cos^2 A - \sin^2 A}{(\cos A - \sin A)^2} = \frac{\cos 2A}{1 - \sin 2A}.$$

$$(20) \cot(45^\circ - A) = \frac{1}{\tan(45^\circ - A)} = \frac{1}{\frac{\tan 45^\circ - \tan A}{1 + \tan 45^\circ \cdot \tan A}} = \frac{1 + \tan A}{1 - \tan A} \\ = \frac{\cos A + \sin A}{\cos A - \sin A} = \frac{(\cos A + \sin A)^2}{\cos^2 A - \sin^2 A} = \frac{1 + \sin 2A}{\cos 2A} = \sec 2A + \tan 2A.$$

$$(21) \frac{1 + \sin a}{1 + \cos a} = \frac{\cos^2 \frac{a}{2} + \sin^2 \frac{a}{2} + 2 \sin \frac{a}{2} \cdot \cos \frac{a}{2}}{2 \cos^2 \frac{a}{2}} = \frac{1}{2} + \frac{1}{2} \tan^2 \frac{a}{2} + \tan \frac{a}{2} \\ = \frac{1}{2} \left(1 + \tan^2 \frac{a}{2} + 2 \tan \frac{a}{2}\right) = \frac{1}{2} \left(1 + \tan \frac{a}{2}\right)^2.$$

$$(22) \frac{1 - \sin a}{1 - \cos a} = \frac{\cos^2 \frac{a}{2} + \sin^2 \frac{a}{2} - 2 \sin \frac{a}{2} \cdot \cos \frac{a}{2}}{2 \sin^2 \frac{a}{2}} = \frac{1}{2} \cot^2 \frac{a}{2} + \frac{1}{2} - \cot \frac{a}{2} \\ = \frac{1}{2} \left(\cot^2 \frac{a}{2} + 1 - 2 \cot \frac{a}{2}\right) = \frac{1}{2} \left(\cot \frac{a}{2} - 1\right)^2.$$

$$\begin{aligned}
 (23) \quad \tan \frac{\theta}{2} + \frac{1}{2} \tan \theta \cdot \sec^2 \frac{\theta}{2} &= \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}} + \frac{\sin \theta}{2 \cos \theta \cdot \cos^2 \frac{\theta}{2}} \\
 &= \frac{2 \cos \theta \cdot \cos \frac{\theta}{2} \cdot \sin \frac{\theta}{2} + \sin \theta}{2 \cos \theta \cdot \cos^2 \frac{\theta}{2}} = \frac{\cos \theta \cdot \sin \theta + \sin \theta}{2 \cos \theta \cdot \cos^2 \frac{\theta}{2}} \\
 &= \frac{\sin \theta (\cos \theta + 1)}{2 \cos \theta \cdot \cos^2 \frac{\theta}{2}} = \frac{\sin \theta \cdot 2 \cos^2 \frac{\theta}{2}}{2 \cos \theta \cdot \cos^2 \frac{\theta}{2}} = \tan \theta.
 \end{aligned}$$

$$(24) \quad \frac{1 + \sin \theta}{1 - \sin \theta} = \frac{(1 + \sin \theta)^2}{1 - \sin^2 \theta} = \left( \frac{1 + \sin \theta}{\cos \theta} \right)^2 = (\sec \theta + \tan \theta)^2.$$

$$\begin{aligned}
 (25) \quad \frac{1 - \tan^2(45^\circ - A)}{1 + \tan^2(45^\circ - A)} &= \frac{\cos^2(45^\circ - A) - \sin^2(45^\circ - A)}{\cos^2(45^\circ - A) + \sin^2(45^\circ - A)} \\
 &= \frac{\cos 2(45^\circ - A)}{1} = \cos(90^\circ - 2A) = \sin 2A.
 \end{aligned}$$

$$\begin{aligned}
 (26) \quad \frac{\tan\left(\frac{\pi}{4} + \theta\right) - \tan\left(\frac{\pi}{4} - \theta\right)}{\tan\left(\frac{\pi}{4} + \theta\right) + \tan\left(\frac{\pi}{4} - \theta\right)} &= \frac{\frac{1 + \tan \theta}{1 - \tan \theta} - \frac{1 - \tan \theta}{1 + \tan \theta}}{\frac{1 + \tan \theta}{1 - \tan \theta} + \frac{1 - \tan \theta}{1 + \tan \theta}} \\
 &= \frac{1 + 2 \tan \theta + \tan^2 \theta - 1 + 2 \tan \theta - \tan^2 \theta}{1 + 2 \tan \theta + \tan^2 \theta + 1 - 2 \tan \theta + \tan^2 \theta} \\
 &= \frac{4 \tan \theta}{2 + 2 \tan^2 \theta} = \frac{2 \tan \theta}{1 + \tan^2 \theta} = 2 \sin \theta \cdot \cos \theta = \sin 2\theta.
 \end{aligned}$$

## EXAMPLES—XXXV. (p. 103).

$$\begin{aligned}
 1. \quad (1) \quad \frac{\cos 3\theta - \sin 3\theta}{\sin \theta + \cos \theta} &= \frac{4 \cos^3 \theta - 3 \cos \theta - 3 \sin \theta + 4 \sin^3 \theta}{\sin \theta + \cos \theta} \\
 &= \frac{4(\sin^3 \theta + \cos^3 \theta) - 3(\sin \theta + \cos \theta)}{\sin \theta + \cos \theta} \\
 &= 4(\sin^2 \theta - \sin \theta \cdot \cos \theta + \cos^2 \theta) - 3 \\
 &= 1 - 4 \sin \theta \cdot \cos \theta = 1 - 2 \sin 2\theta.
 \end{aligned}$$

$$(2) \frac{2 \tan \theta + \sec \theta}{1 + \tan^2 \theta} = \frac{2 \tan \theta + \sec \theta}{\sec^2 \theta} = 2 \tan \theta \cdot \cos^2 \theta + \cos \theta = \sin 2\theta + \cos \theta.$$

$$(3) \tan \frac{A}{2} + 2 \sin^2 \frac{A}{2} \cot A = \sin \frac{A}{2} \left\{ \frac{1}{\cos \frac{A}{2}} + 2 \sin \frac{A}{2} \cdot \frac{\cos A}{\sin A} \right\}$$

$$= \sin \frac{A}{2} \left\{ \frac{1}{\cos \frac{A}{2}} + \frac{\cos A}{\cos \frac{A}{2}} \right\} = \sin \frac{A}{2} \left( \frac{2 \cos^2 \frac{A}{2}}{\cos \frac{A}{2}} \right) = 2 \sin \frac{A}{2} \cdot \cos \frac{A}{2} = \sin A.$$

$$(4) \frac{\cot A}{\cot A - \cot 3A} + \frac{\tan A}{\tan A - \tan 3A}$$

$$= \frac{\frac{1}{\tan A}}{\frac{1}{\tan A} - \frac{1 - 3 \tan^2 A}{\tan A (3 - \tan^2 A)}} + \frac{\tan A}{\tan A - \frac{\tan A (3 - \tan^2 A)}{1 - 3 \tan^2 A}}$$

$$= \frac{1}{3 - \tan^2 A - 1 + 3 \tan^2 A} + \frac{1}{1 - 3 \tan^2 A - 3 + \tan^2 A}$$

$$= \frac{3 - \tan^2 A}{2(1 + \tan^2 A)} + \frac{1 - 3 \tan^2 A}{-2(1 + \tan^2 A)}$$

$$= \frac{3 - \tan^2 A - 1 + 3 \tan^2 A}{2(1 + \tan^2 A)} = \frac{2 + 2 \tan^2 A}{2(1 + \tan^2 A)} = 1.$$

$$(5) \cos 4A + \cos 4B = 2 \cos 2(A+B) \cdot \cos 2(A-B)$$

$$= 2 \cdot \{1 - 2 \sin^2(A+B)\} \cdot \{1 - 2 \sin^2(A-B)\}.$$

$$(6) \tan(45^\circ + \theta) - \tan(45^\circ - \theta) = \frac{1 + \tan \theta}{1 - \tan \theta} - \frac{1 - \tan \theta}{1 + \tan \theta} = \frac{4 \tan \theta}{1 - \tan^2 \theta}$$

$$= \frac{\frac{4 \sin \theta}{\cos \theta}}{1 - \frac{\sin^2 \theta}{\cos^2 \theta}} = \frac{4 \sin \theta \cdot \cos \theta}{\cos^2 \theta - \sin^2 \theta} = \frac{2 \sin 2\theta}{\cos 2\theta} = \frac{2 \sin^2 2\theta}{\cos 2\theta \cdot \sin 2\theta}$$

$$= \frac{2(1 - \cos^2 2\theta)}{\cos 2\theta \cdot \sin 2\theta}$$

$$= 2 \cdot \frac{1 - \cos^2 2\theta}{\sin 2\theta} = 2 \cdot \frac{\sec 2\theta - \cos 2\theta}{\sin 2\theta}.$$



$$\begin{aligned}
 (7) \quad \cot^2 \theta - \tan^2 \theta &= \frac{\cos^4 \theta - \sin^4 \theta}{\cos^2 \theta \cdot \sin^2 \theta} = \frac{(\cos^2 \theta + \sin^2 \theta)(\cos^2 \theta - \sin^2 \theta)}{\cos^2 \theta \cdot \sin^2 \theta} \\
 &= \frac{\cos^2 \theta - \sin^2 \theta}{\cos^2 \theta \cdot \sin^2 \theta} = \frac{\cos 2\theta}{\cos^2 \theta \cdot \sin^2 \theta} = \frac{4 \cos 2\theta}{4 \cos^2 \theta \cdot \sin^2 \theta} = \frac{4 \cos 2\theta}{\sin^2 2\theta} \\
 &= \frac{8 \cos 2\theta}{2 \sin^2 2\theta} = \frac{8 \cos 2\theta}{1 - \cos 4\theta}.
 \end{aligned}$$

(8)

$$2 \sin A \cdot \cos 2A = 2 \sin A (1 - 2 \sin^2 A) = 2 \sin A - 4 \sin^3 A = \sin 3A - \sin A.$$

$$(9) \quad \frac{\cos nA - \cos(n+2)A}{\sin(n+2)A - \sin nA} = \frac{2 \sin(n+1)A \cdot \sin A}{2 \cos(n+1)A \cdot \sin A} = \tan(n+1)A.$$

(10)

$$\begin{aligned}
 \cos 9A + 3 \cos 7A + 3 \cos 5A + \cos 3A &= \cos 9A + \cos 3A + 3(\cos 7A + \cos 5A) \\
 &= 2 \cos 6A \cdot \cos 3A + 6 \cos 6A \cdot \cos A \\
 &= 2 \cos 6A (\cos 3A + 3 \cos A) \\
 &= 2 \cos 6A \cdot 4 \cos^3 A = 8 \cos^3 A \cdot \cos 6A.
 \end{aligned}$$

$$(11) \quad \frac{\operatorname{cosec} 2A - \cot 2A}{\operatorname{cosec} 2A + \cot 2A} = \frac{1 - \cos 2A}{1 + \cos 2A} = \frac{2 \sin^2 A}{2 \cos^2 A} = \tan^2 A.$$

$$\begin{aligned}
 (12) \quad \frac{1 - \sin A}{1 + \cos A} &= \frac{\sin^2 \frac{A}{2} + \cos^2 \frac{A}{2} - 2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}}{1 + 2 \cos^2 \frac{A}{2} - 1} = \frac{\left(\cos \frac{A}{2} - \sin \frac{A}{2}\right)^2}{2 \cos^2 \frac{A}{2}} \\
 &= \frac{1}{2} \left( \frac{\cos \frac{A}{2} - \sin \frac{A}{2}}{\cos \frac{A}{2}} \right)^2 = \frac{1}{2} \left( 1 - \tan \frac{A}{2} \right)^2.
 \end{aligned}$$

$$\begin{aligned}
 (13) \quad \frac{\cos 3A - 2 \cos A}{\sin 3A + 2 \sin A} \cdot \tan A &= \frac{4 \cos^2 A - 3 \cos A - 2 \cos A}{3 \sin A - 4 \sin^3 A + 2 \sin A} \cdot \frac{\sin A}{\cos A} \\
 &= \frac{4 \cos^2 A - 3 - 2}{3 - 4 \sin^2 A + 2} = \frac{2(2 \cos^2 A - 1) - 3}{2(1 - 2 \sin^2 A) + 3} = \frac{2 \cos 2A - 3}{2 \cos 2A + 3}.
 \end{aligned}$$

$$(14) \tan(45^\circ - A) + \tan(45^\circ + A) = \frac{1 - \tan A}{1 + \tan A} + \frac{1 + \tan A}{1 - \tan A}$$

$$= \frac{2(1 + \tan^2 A)}{1 - \tan^2 A} = \frac{2 \sec^2 A}{(\cos^2 A - \sin^2 A) \sec^2 A} = \frac{2}{\cos^2 A - \sin^2 A} = 2 \sec 2A.$$

$$(15) \cos 2a + \tan \frac{a}{2} \sin 2a = \cos 2a + \frac{\sin \frac{a}{2}}{\cos \frac{a}{2}} \cdot 2 \sin a \cdot \cos a$$

$$= \cos 2a + 4 \sin^2 \frac{a}{2} \cdot \cos a = 2 \cos^2 a - 1 + 4 \sin^2 \frac{a}{2} \cdot \cos a$$

$$= 2 \cos a \left( \cos a + 2 \sin^2 \frac{a}{2} \right) - 1 = 2 \cos a \cdot 1 - 1 = 2 \cos a - 1$$

$$= \cos a + \cos a - 1 = \cos a - 2 \sin^2 \frac{a}{2} = \cos a - \frac{2 \cdot \sin^2 \frac{a}{2} \cdot \cos \frac{a}{2}}{\cos \frac{a}{2}}$$

$$= \cos a - \tan \frac{a}{2} \cdot \sin a.$$

$$(16) \cot^2 A - \tan^2 A = \frac{\cos^4 A - \sin^4 A}{\cos^2 A \cdot \sin^2 A} = \frac{(\cos^2 A + \sin^2 A)(\cos^2 A - \sin^2 A)}{\cos^2 A \cdot \sin^2 A}$$

$$= \frac{\cos^2 A - \sin^2 A}{\cos^2 A \cdot \sin^2 A} = \frac{4(\cos^2 A - \sin^2 A)}{4 \cos^2 A \cdot \sin^2 A} = \frac{4 \cos 2A}{\sin^2 2A} = 4 \cot 2A \cdot \operatorname{cosec} 2A.$$

$$(17) \operatorname{cosec} a \cdot \cot a - \sec a \cdot \tan a = \frac{\cos a}{\sin^2 a} - \frac{\sin a}{\cos^2 a} = \frac{\cos^3 a - \sin^3 a}{\sin^2 a \cdot \cos^2 a}$$

$$= \frac{4(\cos^3 a - \sin^3 a)}{4 \sin^2 a \cdot \cos^2 a} = \frac{4(\cos^3 a - \sin^3 a)}{\sin^2 2a} = 4 \operatorname{cosec}^2 2a \cdot (\cos^3 a - \sin^3 a).$$

$$(18) \cot^2 a - \tan^2 a = \frac{\cos^2 a - \sin^2 a}{\cos^2 a \cdot \sin^2 a} = \frac{4(\cos^2 a - \sin^2 a)}{4 \cos^2 a \cdot \sin^2 a} = \frac{4 \cos 2a}{\sin^2 2a}.$$

$$(19) \operatorname{cosec}^2 b - \sec^2 b = \frac{1}{\sin^2 b} - \frac{1}{\cos^2 b} = \frac{\cos^2 b - \sin^2 b}{\sin^2 b \cdot \cos^2 b} = \frac{4(\cos^2 b - \sin^2 b)}{4 \sin^2 b \cdot \cos^2 b}$$

$$= \frac{4 \cos 2b}{\sin^2 2b} = 4 \cos 2b \cdot \operatorname{cosec}^2 2b.$$

$$\begin{aligned}
 (20) \quad \frac{2 \operatorname{cosec} 2A - \sec A}{2 \operatorname{cosec} 2A + \sec A} &= \frac{2 - \sec A \cdot \sin 2A}{2 + \sec A \cdot \sin 2A} = \frac{2 - 2 \sin A}{2 + 2 \sin A} = \frac{1 - \sin A}{1 + \sin A} \\
 &= \frac{\sin^2 \frac{A}{2} + \cos^2 \frac{A}{2} - 2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}}{\sin^2 \frac{A}{2} + \cos^2 \frac{A}{2} + 2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}} = \left( \frac{\cos \frac{A}{2} - \sin \frac{A}{2}}{\cos \frac{A}{2} + \sin \frac{A}{2}} \right)^2 \\
 &= \left( \frac{1 - \tan \frac{A}{2}}{1 + \tan \frac{A}{2}} \right)^2 = \cot^2 \left( 45^\circ + \frac{A}{2} \right).
 \end{aligned}$$

$$\begin{aligned}
 (21) \quad \sin \left( \frac{5\pi}{2} + \theta \right) - \sin \left( \frac{3\pi}{2} - \theta \right) &= 2 \cos 2\pi \cdot \sin \left( \frac{\pi}{2} + \theta \right) \\
 &= 2 \cos 2\pi \cdot \cos \theta = 2 \cos 2\pi \cdot \sin \left( \frac{\pi}{2} - \theta \right) \\
 &= \sin \left( \frac{5\pi}{2} - \theta \right) - \sin \left( \frac{3\pi}{2} + \theta \right). \quad (\text{Art. 122.})
 \end{aligned}$$

$$\begin{aligned}
 (22) \quad \cot \left( \frac{\pi}{2} + \theta \right) - \tan \left( \frac{\pi}{2} + \theta \right) &= \frac{\cos^2 \left( \frac{\pi}{2} + \theta \right) - \sin^2 \left( \frac{\pi}{2} + \theta \right)}{\sin \left( \frac{\pi}{2} + \theta \right) \cdot \cos \left( \frac{\pi}{2} + \theta \right)} \\
 &= \frac{\sin^2 \theta - \cos^2 \theta}{-\cos \theta \cdot \sin \theta} = \frac{\cos^2 \theta - \sin^2 \theta}{\sin \theta \cdot \cos \theta} = \frac{2 \cdot \cos 2\theta}{\sin 2\theta} = 2 \cot 2\theta.
 \end{aligned}$$

$$\begin{aligned}
 (23) \quad \frac{(\operatorname{cosec} a + \sec a)^2}{\operatorname{cosec}^2 a + \sec^2 a} &= \frac{\left( \frac{\cos a + \sin a}{\sin a \cdot \cos a} \right)^2}{\frac{1}{\sin^2 a \cdot \cos^2 a}} = (\cos a + \sin a)^2 = 1 + 2 \sin a \cdot \cos a \\
 &= 1 + \sin 2a.
 \end{aligned}$$

$$\begin{aligned}
 (24) \quad \frac{\tan \theta}{\tan 2\theta - \tan \theta} &= \frac{\tan \theta}{\frac{2 \tan \theta}{1 - \tan^2 \theta} - \tan \theta} = \frac{1}{\frac{2}{1 - \tan^2 \theta} - 1} = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \\
 &= \cos^2 \theta - \sin^2 \theta = \cos 2\theta.
 \end{aligned}$$

$$\begin{aligned}
 (25) \quad \frac{\tan 2\theta \cdot \tan \theta}{\tan 2\theta - \tan \theta} &= \frac{\frac{2 \tan \theta}{1 - \tan^2 \theta} \cdot \tan \theta}{\frac{2 \tan \theta}{1 - \tan^2 \theta} - \tan \theta} = \frac{\frac{2 \tan^2 \theta}{1 - \tan^2 \theta}}{\frac{2}{1 - \tan^2 \theta} - 1} = \frac{2 \tan^2 \theta}{1 + \tan^2 \theta} \\
 &= \frac{2 \tan \theta}{\sec^2 \theta} = 2 \sin \theta \cdot \cos \theta = \sin 2\theta.
 \end{aligned}$$

$$\begin{aligned}
 (26) \quad \frac{\sin^2 \alpha - \sin^2 \beta}{\sin \alpha \cdot \cos \alpha - \sin \beta \cdot \cos \beta} &= \frac{\sin(\alpha + \beta) \cdot \sin(\alpha - \beta)}{\sin \alpha \cdot \cos \alpha - \sin \beta \cdot \cos \beta} \\
 &\quad \text{(Ex. XXVII. 1.)} \\
 &= \frac{2 \sin(\alpha + \beta) \cdot \sin(\alpha - \beta)}{2 \sin \alpha \cdot \cos \alpha - 2 \sin \beta \cdot \cos \beta} = \frac{2 \sin(\alpha + \beta) \cdot \sin(\alpha - \beta)}{\sin 2\alpha - \sin 2\beta} \\
 &= \frac{2 \sin(\alpha + \beta) \cdot \sin(\alpha - \beta)}{2 \cos(\alpha + \beta) \cdot \sin(\alpha - \beta)} = \tan(\alpha + \beta).
 \end{aligned}$$

$$\begin{aligned}
 (27) \quad 4 \sin A \cdot \sin(60^\circ + A) \cdot \sin(60^\circ - A) &= 4 \sin A \cdot (\sin^2 60^\circ - \sin^2 A). \\
 &\quad \text{(Ex. XXVII. 1.)} \\
 &= 4 \sin A \left( \frac{3}{4} - \sin^2 A \right) = 3 \sin A - 4 \sin^3 A = \sin 3A.
 \end{aligned}$$

$$\begin{aligned}
 (28) \quad \operatorname{cosec} 2\theta + \cot 4\theta + \operatorname{cosec} 4\theta &= \frac{1}{\sin 2\theta} + \frac{\cos 4\theta}{\sin 4\theta} + \frac{1}{\sin 4\theta} \\
 &= \frac{2 \cos 2\theta + \cos 4\theta + 1}{2 \sin 2\theta \cdot \cos 2\theta} = \frac{2 \cos 2\theta + 2 \cos^2 2\theta}{2 \sin 2\theta \cdot \cos 2\theta} = \frac{1 + \cos 2\theta}{\sin 2\theta} \\
 &= \frac{2 \cos^2 \theta}{2 \sin \theta \cdot \cos \theta} = \frac{\cos \theta}{\sin \theta} = \cot \theta.
 \end{aligned}$$

$$\begin{aligned}
 2. \quad (1) \quad \sin 2\theta + \sqrt{3} \cdot \cos 2\theta &= 1, \\
 \sqrt{3} \cdot \cos 2\theta &= 1 - \sin 2\theta, \\
 3 \cdot \cos^2 2\theta &= 1 - 2 \sin 2\theta + \sin^2 2\theta, \\
 3 - 3 \sin^2 2\theta &= 1 - 2 \sin 2\theta + \sin^2 2\theta.
 \end{aligned}$$

Solving this quadratic, we obtain  $\sin 2\theta = -\frac{1}{2}$ , or, 1 ;

$$\begin{aligned}
 \therefore 2\theta &= -30^\circ, \text{ or, } 90^\circ; \\
 \therefore \theta &= -15^\circ, \text{ or, } 45^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (2) \quad \sin^2 2\theta - \sin^2 \theta &= \sin^2 \frac{\pi}{4}, \\
 4 \sin^2 \theta \cdot \cos^2 \theta - \sin^2 \theta &= \frac{1}{2}, \\
 4 \sin^2 \theta - 4 \sin^4 \theta - \sin^2 \theta &= \frac{1}{2}.
 \end{aligned}$$

Solving this quadratic, we obtain  $\sin^2 \theta = \frac{1}{2}$ , or,  $\frac{1}{4}$  ;

$$\begin{aligned}
 \therefore \sin \theta &= \frac{1}{\sqrt{2}}, \text{ or, } \frac{1}{2}; \\
 \therefore \theta &= 45^\circ, \text{ or, } 30^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (3) \quad & \sin 5x \cdot \cos 3x = \sin 9x \cdot \cos 7x; \\
 & \therefore \sin 8x + \sin 2x = \sin 16x + \sin 2x; \\
 & \therefore \sin 8x = \sin 16x, \\
 & \sin 8x = 2 \sin 8x \cdot \cos 8x. \\
 & \text{Hence } \sin 8x = 0, \text{ or, } 2 \cos 8x = 1, \\
 & \sin 8x = 0, \text{ or, } \cos 8x = \frac{1}{2}; \\
 & \therefore x = 0^\circ, \text{ or, } 8x = 60^\circ, \text{ and } \therefore x = 7\frac{1}{2}^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (4) \quad & 2 \sin^2 3\theta + \sin^2 6\theta = 2, \\
 & \sin^2 6\theta = 2(1 - \sin^2 3\theta), \\
 & 4 \sin^2 3\theta \cdot \cos^2 3\theta = 2 \cos^2 3\theta, \\
 & 2 \sin^2 3\theta \cdot \cos 3\theta = \sqrt{2} \cos 3\theta. \\
 & \text{Hence } \cos 3\theta = 0, \text{ or, } \sin 3\theta = \frac{1}{\sqrt{2}}; \\
 & \therefore 3\theta = 90^\circ, \text{ or, } 3\theta = 45^\circ; \\
 & \therefore \theta = 30^\circ, \text{ or, } 15^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad & \cos 2A + \sin^2 A = \frac{3}{4} \\
 & 1 - 2\sin^2 A + \sin^2 A = \frac{3}{4}, \\
 & \sin^2 A = \frac{1}{4}, \text{ and } \therefore \sin A = \pm \frac{1}{2}. \\
 & \text{Hence } A = 30^\circ, \text{ or, } 150^\circ.
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad & \cos 3\theta - \cos 5\theta = \sin \theta, \\
 & 2 \sin 4\theta \cdot \sin \theta = \sin \theta. \\
 & \text{Hence } \sin \theta = 0, \text{ or, } \sin 4\theta = \frac{1}{2}; \\
 & \therefore \theta = 0^\circ, \text{ or, } 4\theta = 30^\circ; \\
 & \therefore \theta = 0^\circ, \text{ or, } \theta = 7\frac{1}{2}^\circ.
 \end{aligned}$$

$$\begin{aligned}(7) \quad & \sin 5\theta - \cos 3\theta = \sin \theta, \\ & \sin 5\theta - \sin \theta = \cos 3\theta, \\ & 2 \cos 3\theta \cdot \sin 2\theta = \cos 3\theta.\end{aligned}$$

$$\text{Hence } \cos 3\theta = 0, \text{ or, } \sin 2\theta = \frac{1}{2};$$

$$\therefore 3\theta = 90^\circ, \text{ or, } 2\theta = 30^\circ,$$

$$\therefore \theta = 30^\circ, \text{ or, } \theta = 15^\circ.$$

$$\begin{aligned}(8) \quad & \tan 2a = 3 \tan a, \\ & \frac{2 \tan a}{1 - \tan^2 a} = 3 \tan a.\end{aligned}$$

$$\text{Hence } \tan a = 0, \text{ and } \therefore a = 0^\circ,$$

$$\text{or, } 2 = 3 - 3 \tan^2 a,$$

$$\tan^2 a = \frac{1}{3}, \text{ or, } \tan a = \frac{1}{\sqrt{3}}, \text{ or, } a = 30^\circ.$$

$$\begin{aligned}(9) \quad & \sin 2\theta + \sin \theta = \cos 2\theta + \cos \theta, \\ & 2 \sin \frac{3\theta}{2} \cdot \cos \frac{\theta}{2} = 2 \cos \frac{3\theta}{2} \cdot \cos \frac{\theta}{2}.\end{aligned}$$

$$\therefore \cos \frac{\theta}{2} = 0, \text{ or, } \frac{\theta}{2} = 90^\circ, \text{ or, } \theta = 180^\circ;$$

$$\text{or, } \sin \frac{3\theta}{2} = \cos \frac{3\theta}{2}, \text{ or, } \tan \frac{3\theta}{2} = 1, \text{ or, } \frac{3\theta}{2} = 45^\circ, \text{ or, } \theta = 30^\circ.$$

$$\begin{aligned}(10) \quad & \sin 7a - \sin a = \sin 3a, \\ & 2 \cos 4a \cdot \sin 3a = \sin 3a.\end{aligned}$$

$$\text{Hence } \sin 3a = 0, \text{ or, } 3a = 0^\circ, \text{ or, } a = 0^\circ \left. \vphantom{\begin{matrix} \text{or, } 3a = 180^\circ, \text{ or, } a = 60^\circ \end{matrix}} \right\},$$

$$\text{or, } 2 \cos 4a = 1, \text{ or, } 4a = 60^\circ, \text{ or, } a = 15^\circ.$$

$$(11) \quad \operatorname{cosec}^2 \theta - \sec^2 \theta = 2 \operatorname{cosec}^2 \theta + 3,$$

$$\frac{\operatorname{cosec}^2 \theta}{3} = \sec^2 \theta, \text{ or, } \cos^2 \theta = 3 \sin^2 \theta;$$

$$\therefore 4 \sin^2 \theta = 1, \text{ or, } \sin \theta = \frac{1}{2}, \text{ and } \therefore \theta = 30^\circ.$$



$$(12) \quad \sin 6\theta = \sin 4\theta - \sin 2\theta,$$

$$\sin 6\theta + \sin 2\theta = \sin 4\theta,$$

$$2 \sin 4\theta \cdot \cos 2\theta = \sin 4\theta.$$

$$\text{Hence } \sin 4\theta = 0, \text{ or, } 4\theta = 0^\circ, \text{ or, } \theta = 0^\circ,$$

$$\text{or, } 2 \cos 2\theta = 1, \text{ or, } \cos 2\theta = \frac{1}{2}, \text{ or, } \theta = 30^\circ.$$

## EXAMPLES—XXXVI. (p. 106).

$$\begin{aligned} 1. \quad (1) \quad \sin 36^\circ &= 2 \sin 18^\circ \cdot \cos 18^\circ = 2 \cdot \frac{\sqrt{5}-1}{4} \cdot \frac{\sqrt{(10+2\sqrt{5})}}{4} \\ &= \frac{2\sqrt{(40-8\sqrt{5})}}{16} = \frac{\sqrt{(10-2\sqrt{5})}}{4}. \end{aligned}$$

$$(2) \quad \cos 36^\circ = 1 - 2\sin^2 18^\circ = 1 - 2 \cdot \left(\frac{\sqrt{5}-1}{4}\right)^2 = 1 - \frac{6-2\sqrt{5}}{8} = \frac{1+\sqrt{5}}{4}$$

$$(3) \quad \sin 54^\circ = \cos 36^\circ = \frac{1+\sqrt{5}}{4}.$$

$$(4) \quad \cos 54^\circ = \sin 36^\circ = \frac{\sqrt{(10-2\sqrt{5})}}{4}.$$

$$(5) \quad \sin 72^\circ = \cos 18^\circ = \frac{\sqrt{(10+2\sqrt{5})}}{4}.$$

$$(6) \quad \tan 72^\circ = \frac{\sin 72^\circ}{\cos 72^\circ} = \frac{\cos 18^\circ}{\sin 18^\circ} = \frac{\sqrt{(10+2\sqrt{5})}}{4} \div \frac{\sqrt{5}-1}{4} = \frac{\sqrt{(10+2\sqrt{5})}}{\sqrt{5}-1}$$

$$\begin{aligned} (7) \quad \sin 90^\circ &= \sin(18^\circ + 72^\circ) = \sin 18^\circ \cdot \cos 72^\circ + \cos 18^\circ \cdot \sin 72^\circ \\ &= \sin 18^\circ \cdot \sin 18^\circ + \cos 18^\circ \cdot \cos 18^\circ \\ &= \left(\frac{\sqrt{5}-1}{4}\right)^2 + \left(\frac{\sqrt{(10+2\sqrt{5})}}{4}\right)^2 \\ &= \frac{6-2\sqrt{5}+10+2\sqrt{5}}{16} = \frac{16}{16} = 1 \end{aligned}$$

$$\begin{aligned} (8) \quad \cos 90^\circ &= \cos(18^\circ + 72^\circ) = \cos 18^\circ \cdot \cos 72^\circ - \sin 18^\circ \cdot \sin 72^\circ \\ &= \cos 18^\circ \cdot \cos 72^\circ - \cos 72^\circ \cdot \cos 18^\circ = 0. \end{aligned}$$



$$\begin{aligned}
 2. \quad & \sin(36^\circ + A) + \sin(72^\circ - A) - \sin(36^\circ - A) - \sin(72^\circ + A) \\
 &= \{\sin(36^\circ + A) - \sin(36^\circ - A)\} - \{\sin(72^\circ + A) - \sin(72^\circ - A)\} \\
 &= 2 \cos 36^\circ \cdot \sin A - 2 \cos 72^\circ \cdot \sin A \\
 &= \sin A \{2 \cos 36^\circ - 2 \cos 72^\circ\} = \sin A \left\{ \frac{1 + \sqrt{5}}{2} - \frac{\sqrt{5} - 1}{2} \right\} = \sin A.
 \end{aligned}$$

Also,

$$\begin{aligned}
 & \{\sin(54^\circ + A) + \sin(54^\circ - A)\} - \{\sin(18^\circ + A) + \sin(18^\circ - A)\} \\
 &= 2 \sin 54^\circ \cdot \cos A - 2 \sin 18^\circ \cdot \cos A \\
 &= \cos A \{2 \sin 54^\circ - 2 \sin 18^\circ\} = \cos A \left\{ \frac{1 + \sqrt{5}}{2} - \frac{\sqrt{5} - 1}{2} \right\} = \cos A.
 \end{aligned}$$

EXAMPLES—XXXVII (p. 110).

(1) At  $7\frac{1}{2}^\circ$  the cosine is greater than the sine, and both are positive;

$$\begin{aligned}
 \therefore \cos \frac{A}{2} + \sin \frac{A}{2} &= +\sqrt{1 + \sin A}, \\
 \cos \frac{A}{2} - \sin \frac{A}{2} &= +\sqrt{1 - \sin A}.
 \end{aligned}$$

(2) At  $150^\circ$  the cosine (negative) is greater than the sine (positive);

$$\begin{aligned}
 \therefore \cos \frac{A}{2} + \sin \frac{A}{2} &= -\sqrt{1 + \sin A}, \\
 \cos \frac{A}{2} - \sin \frac{A}{2} &= -\sqrt{1 - \sin A}.
 \end{aligned}$$

(3)  $\cos 189^\circ + \sin 189^\circ = -\sqrt{1 + \sin 378^\circ}$ ,

$$\cos 189^\circ - \sin 189^\circ = -\sqrt{1 - \sin 378^\circ};$$

$$\begin{aligned}
 \therefore \cos 189^\circ &= -\frac{1}{2} \cdot \left\{ \sqrt{1 + \frac{\sqrt{5}-1}{4}} + \sqrt{1 - \frac{\sqrt{5}-1}{4}} \right\} \\
 &= -\frac{1}{2} \cdot \left\{ \frac{\sqrt{3+\sqrt{5}}}{2} + \frac{\sqrt{5-\sqrt{5}}}{2} \right\} \\
 &= -\frac{1}{4} \left\{ \sqrt{3+\sqrt{5}} + \sqrt{5-\sqrt{5}} \right\},
 \end{aligned}$$

$$\begin{aligned}\text{and } \sin 189^\circ &= \frac{1}{2} \left\{ \sqrt{1 - \frac{\sqrt{5}-1}{4}} - \sqrt{1 + \frac{\sqrt{5}-1}{4}} \right\} \\ &= \frac{1}{4} \left\{ \sqrt{5} - \sqrt{5} - \sqrt{3 + \sqrt{5}} \right\}.\end{aligned}$$

$$\begin{aligned}(4) \quad 2 \sin 9^\circ.44'.30'' &= \sqrt{1 + \frac{1}{3}} - \sqrt{1 - \frac{1}{3}} \\ &= \sqrt{\frac{4}{3}} - \sqrt{\frac{2}{3}} = \frac{2 - \sqrt{2}}{\sqrt{3}}; \\ \therefore \sin 9^\circ.44'.30'' &= \frac{2 - \sqrt{2}}{2\sqrt{3}}.\end{aligned}$$

$$\begin{aligned}(5) \quad \cos 157^\circ.30' &= -\sqrt{\frac{1 + \cos 315^\circ}{2}} = -\sqrt{\frac{1 + \frac{1}{\sqrt{2}}}{2}} = -\sqrt{\frac{\sqrt{2} + 1}{2\sqrt{2}}} \\ &= -\sqrt{\frac{2 + \sqrt{2}}{4}} = -\frac{\sqrt{2 + \sqrt{2}}}{2}.\end{aligned}$$

## EXAMPLES—XXXVIII. (p. 111).

$$\begin{aligned}(1) \quad \sin A &= \frac{3}{5} \text{ and } \sin B = \frac{4}{5}, \\ \cos A &= \frac{4}{5} \text{ and } \cos B = \frac{3}{5}; \\ \therefore \sin(A+B) &= \frac{3}{5} \cdot \frac{3}{5} + \frac{4}{5} \cdot \frac{4}{5} = \frac{25}{25} = 1; \\ \therefore A+B &= 90^\circ.\end{aligned}$$

$$\begin{aligned}(2) \quad \tan A &= \frac{1}{7}; \tan B = \frac{1}{3}, \\ \tan 2B &= \frac{2 \tan B}{1 - \tan^2 B} = \frac{2}{3} \div \left(1 - \frac{1}{9}\right) = \frac{2 \times 9}{3 \times 8} = \frac{3}{4}; \\ \therefore \tan(A+2B) &= \frac{\tan A + \tan 2B}{1 - \tan A \cdot \tan 2B} = \frac{\frac{1}{7} + \frac{3}{4}}{1 - \frac{1}{7} \cdot \frac{3}{4}} = 1; \\ \therefore A+2B &= 45^\circ.\end{aligned}$$

- (3) Let  $\sin A = \frac{1}{\sqrt{5}}$  and  $\cot B = 3$ .

Then  $\tan A = \frac{1}{2}$  and  $\tan B = \frac{1}{3}$ ;

$$\therefore \tan(A+B) = \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \cdot \frac{1}{3}} = 1;$$

$$\therefore A+B=45^\circ,$$

$$\text{that is } \sin^{-1} \frac{1}{\sqrt{5}} + \cot^{-1} 3 = 45^\circ.$$

- (4) Let  $A, B, C, D$  be the four angles whose tangents are

$$\frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \frac{1}{8}.$$

Then  $\tan\{(A+B)+(C+D)\}$

$$= \frac{\tan(A+B) + \tan(C+D)}{1 - \tan(A+B) \cdot \tan(C+D)}$$

$$= \left( \frac{\frac{1}{3} + \frac{1}{5}}{1 - \frac{1}{15}} + \frac{\frac{1}{7} + \frac{1}{8}}{1 - \frac{1}{7} \cdot \frac{1}{8}} \right) \div \left( 1 - \frac{\frac{1}{3} + \frac{1}{5}}{1 - \frac{1}{15}} \cdot \frac{\frac{1}{7} + \frac{1}{8}}{1 - \frac{1}{7} \cdot \frac{1}{8}} \right)$$

$$= \left( \frac{4}{7} + \frac{3}{11} \right) \div \left( 1 - \frac{12}{77} \right) = 1;$$

$$\therefore A+B+C+D=45^\circ.$$

- (5) Let  $\cot A = \frac{3}{4}$  and  $\cot B = \frac{1}{7}$ .

Then  $\tan A = \frac{4}{3}$  and  $\tan B = 7$ ;

$$\therefore \tan(A+B) = \frac{\frac{4}{3} + 7}{1 - \frac{28}{3}} = -1;$$

$$\therefore A+B=135^\circ, \text{ or, } \cot^{-1} \frac{3}{4} + \cot^{-1} \frac{1}{7} = 135^\circ.$$

$$(6) \quad \text{Let } \tan A = \frac{3}{5} \text{ and } \tan B = \frac{3}{7}.$$

$$\text{Then } \tan(A+B) = \frac{\frac{3}{5} + \frac{3}{7}}{1 - \frac{9}{35}} = \frac{18}{13};$$

$$\therefore \cot(A+B) = \frac{13}{18}, \text{ or, } A+B = \cot^{-1} \frac{13}{18}.$$

$$(7) \quad \text{Let } \tan A = x \text{ and } \tan B = y.$$

$$\text{Then } \tan(A-B) = \frac{x-y}{1+xy};$$

$$\therefore \tan^{-1}x - \tan^{-1}y = \tan^{-1} \frac{x-y}{1+xy}.$$

$$(8) \quad \text{Let } \sin A = x \text{ and } \cos B = x.$$

$$\text{Then } \cos A = \sqrt{1-x^2} \text{ and } \sin B = \sqrt{1-x^2};$$

$$\therefore \sin(A+B) = x \cdot x + \sqrt{1-x^2} \cdot \sqrt{1-x^2} \\ = x^2 + 1 - x^2 = 1;$$

$$\therefore A+B = 90^\circ, \text{ or, } \sin^{-1}x + \cos^{-1}x = 90^\circ.$$

$$(9) \quad \text{Let } \sin A = \frac{4}{5}, \sin B = \frac{5}{13}, \sin C = \frac{16}{65};$$

$$\therefore \cos A = \frac{3}{5}, \cos B = \frac{12}{13}, \cos C = \frac{63}{65}.$$

$$\text{Then } \sin(A+B+C) = \sin(A+B) \cdot \cos C + \cos(A+B) \cdot \sin C$$

$$= (\sin A \cdot \cos B + \cos A \cdot \sin B) \frac{63}{65} + (\cos A \cdot \cos B - \sin A \cdot \sin B) \frac{16}{65}$$

$$= \left( \frac{4}{5} \cdot \frac{12}{13} + \frac{3}{5} \cdot \frac{5}{13} \right) \cdot \frac{63}{65} + \left( \frac{3}{5} \cdot \frac{12}{13} - \frac{4}{5} \cdot \frac{5}{13} \right) \frac{16}{65}$$

$$= \frac{63}{65} \cdot \frac{63}{65} + \frac{16}{65} \cdot \frac{16}{65} = \frac{4225}{4225} = 1.$$

$$\therefore A+B+C = 90^\circ, \text{ or, } \sin^{-1} \frac{4}{5} + \sin^{-1} \frac{5}{13} + \sin^{-1} \frac{16}{65} = \frac{\pi}{2}.$$

(10) Let  $\tan A = \frac{1}{5}$ , and  $\tan B = \frac{1}{239}$ .

Then  $\tan(4A - B) = \frac{\tan 4A - \tan B}{1 + \tan 4A \cdot \tan B}$

$$= \frac{\frac{120}{119} - \frac{1}{239}}{1 + \frac{120}{119} \cdot \frac{1}{239}} = 1;$$

$\therefore 4A - B = 45^\circ$ , or,  $4 \tan^{-1} \frac{1}{5} - \tan^{-1} \frac{1}{239} = \frac{\pi}{4}$ .

EXAMPLES—XXXIX. (p. 120).

(1)  $\begin{array}{r} \bar{3} \cdot 1651553 \\ \bar{4} \cdot 7505855 \\ \bar{6} \cdot 6879746 \\ \bar{2} \cdot 6150026 \\ \hline \bar{1} \cdot 2187180 \end{array}$

(2)  $\begin{array}{r} \bar{4} \cdot 6843785 \\ \bar{5} \cdot 6650657 \\ \bar{3} \cdot 8905196 \\ \bar{3} \cdot 4675284 \\ \hline \bar{7} \cdot 7074922 \end{array}$

(3)  $\begin{array}{r} 2 \cdot 5324716 \\ \bar{3} \cdot 6650657 \\ \bar{5} \cdot 8905196 \\ \hline \bar{3} \cdot 156215 \\ \hline 2 \cdot 4036784 \end{array}$

(4)  $\begin{array}{r} \bar{2} \cdot 483269 \\ \bar{3} \cdot 742891 \\ \hline 4 \cdot 740378 \end{array}$

(5)  $\begin{array}{r} \bar{2} \cdot 352678 \\ \bar{5} \cdot 428619 \\ \hline 2 \cdot 924059 \end{array}$

(6)  $\begin{array}{r} \bar{5} \cdot 349162 \\ \bar{3} \cdot 624329 \\ \hline \bar{3} \cdot 724833 \end{array}$

(7)  $\begin{array}{r} \bar{2} \cdot 4596721 \\ \quad \quad \quad 3 \\ \hline \bar{5} \cdot 3790163 \end{array}$

(8)  $\begin{array}{r} \bar{7} \cdot 429683 \\ \quad \quad \quad 6 \\ \hline \bar{40} \cdot 578098 \end{array}$

(9)  $\begin{array}{r} \bar{9} \cdot 2843617 \\ \quad \quad \quad 7 \\ \hline \bar{62} \cdot 9905319 \end{array}$

(10)  $3 \mid \begin{array}{r} \bar{6} \cdot 3725409 \\ \hline \bar{2} \cdot 1241803 \end{array}$

(11)  $6 \mid \begin{array}{r} \bar{14} \cdot 432962 \\ \hline \bar{3} \cdot 738827 \end{array}$

(12)  $9 \mid \begin{array}{r} \bar{4} \cdot 53627188 \\ \hline \bar{1} \cdot 61514132 \end{array}$

## EXAMPLES—XL. (p. 123).

$$1. \log 128 = \log 2^7 = 7 \log 2 = 2.1072100$$

$$\begin{aligned} \log 125 &= \log \frac{1000}{8} = \log 1000 - \log 8 = 3 - \log 2^3 \\ &= 3 - 3 \log 2 = 3 - .90309000 = 2.0969100 \end{aligned}$$

$$\begin{aligned} \log 2500 &= \log \frac{10000}{4} = \log 10000 - \log 4 = 4 - 2 \log 2 \\ &= 4 - .6020600 = 3.3979400. \end{aligned}$$

$$2. \log 50 = \log \frac{100}{2} = \log 100 - \log 2 = 2 - .3010300 = 1.6989700$$

$$\log .005 = \log \frac{5}{1000} = \log 10 - \log 2 - 3 = -\log 2 - 2 = \bar{3}.6989700$$

$$\log 196 = \log (49 \times 4) = 2 \log 7 + 2 \log 2 = 2.2922560.$$

$$3. \log 6 = \log 3 + \log 2 = .7781513$$

$$\log 27 = 3 \log 3 = 1.4313639$$

$$\log 54 = \log (27 \times 2) = 3 \log 3 + \log 2 = 1.7323939$$

$$\log 576 = \log (9 \times 64) = 2 \log 3 + 6 \log 2 = 2.7604226.$$

$$4. \log 60 = \log (2 \times 3 \times 10) = \log 2 + \log 3 + \log 10 = 1.7781513$$

$$\log .03 = \log \frac{3}{100} = \log 3 - 2 = .4771213 - 2 = \bar{2}.4771213$$

$$\log 1.05 = \log \frac{105}{100} = \log \frac{21}{20} = \log 3 + \log 7 - \log 2 - 1 = .0211893$$

$$\log .0000432 = \log \frac{16 \times 27}{10000000} = 4 \log 2 + 3 \log 3 - 7 = \bar{5}.6354839$$

$$5. \log .00075 = \log 75 - 5 = \log 3 + \log 25 - 5 = \log \left(\frac{18}{2}\right)^{\frac{1}{2}} + \log 25 - 5$$

$$= \frac{1}{2} \left\{ \log 18 - \log 2 \right\} + \log 100 - \log 4 - 5$$

$$= \frac{1}{2} \left\{ 1.2552725 - .3010300 \right\} + 2 - .6020600 - 5$$

$$= .4771213 - .6020600 - 3 = \bar{4}.8750613.$$

$$\begin{aligned}\log 31\cdot5 &= \log (21 \times 3 \times 5) - 1 = \log 21 + \log 3 + 1 - \log 2 - 1 \\ &= \log 21 + \frac{1}{2}(\log 18 - \log 2) - \log 2 \\ &= 1\cdot3222193 + \cdot4771212 - \cdot3010300 = 1\cdot4983105.\end{aligned}$$

$$6. \quad \log 2 = \log \frac{10}{5} = 1 - \log 5 = \cdot3010300.$$

$$\begin{aligned}\log \cdot064 &= \log \frac{2^6}{1000} = 6 \log 2 - 3 = 6 - 6 \log 5 - 3 = \bar{2}\cdot8061800 \\ \log \left\{ \frac{2^{80}}{5^{20}} \right\}^{\frac{1}{14}} &= \frac{1}{14} (60 \log 2 - 20 \log 5) \\ &= \frac{1}{7} (30 - 30 \log 5 - 10 \log 5) = \frac{1}{7} (30 - 27\cdot9588000) \\ &= \frac{1}{7} (2\cdot0412000) = \cdot2916000.\end{aligned}$$

$$7. \quad \log 5 = \log \frac{10}{2} = 1 - \cdot3010300 = \cdot6989700,$$

$$\begin{aligned}\log \cdot125 &= \log \frac{5^3}{1000} = 3 \log 5 - 3 = 2\cdot0969100 - 3 = \bar{1}\cdot0969100 \\ \log \left( \frac{5^{80}}{2^{40}} \right)^{\frac{1}{15}} &= \log 5^{\frac{80}{15}} - \log 2^{\frac{40}{15}} = \log 5^6 - \log 2^{\frac{8}{3}} \\ &= 6 \log 5 - \frac{8}{3} \log 2 = 6 (\log 10 - \log 2) - \frac{8}{3} \log 2 \\ &= 4\cdot1938200 - \cdot8027467 = 3\cdot3910733.\end{aligned}$$

$$\begin{aligned}8. \quad \left. \begin{aligned} \cdot01 &= \frac{1}{100} = \frac{1}{10^2} = 10^{-2} \\ 1 &= 10^0 \\ 100 &= 10^2 \end{aligned} \right\} \therefore \text{the logarithms are } -2, 0, 2; \\ \left. \begin{aligned} \cdot01 &= (\cdot01)^1 \\ 1 &= (\cdot01)^0 \\ 100 &= \frac{1}{\cdot01} = (\cdot01)^{-1} \end{aligned} \right\} \therefore \text{the logarithms are } 1, 0, -1.\end{aligned}$$



9. 1593 is greater than  $10^3$  and less than  $10^4$ ; characteristic 3.  
 1593 is greater than  $12^2$  and less than  $12^3$ ; characteristic 2.

$$10. \frac{4^{2y}}{2^{4y}} = 8; \frac{2^{6y}}{2^{4y}} = 2^3; 2^{2y} = 2^3; 2y = 3.$$

$$\text{Hence } y = \frac{3}{2} \text{ and } x = \frac{9}{2}.$$

$$11. (a) \log 2 = \frac{1}{2} \log 4 = \cdot 3010300,$$

$$\log 25 = \log 100 - \log 4 = 2 - \cdot 6020600 = 1\cdot 3979400$$

$$\log 83\cdot 2 = \log (80 \times 1\cdot 04) = \frac{3}{2} \log 4 + \log 10 + \log 1\cdot 04$$

$$= \cdot 9030900 + 1 + \cdot 0170333 = 1\cdot 9201233$$

$$\log (625)^{\frac{1}{100}} = \frac{1}{100} \left\{ \log 625 - \log 1000 \right\} = \frac{1}{100} \left\{ 2 \log 25 - 3 \right\}$$

$$= \frac{1}{100} \left\{ 2 \log 100 - 2 \log 4 - 3 \right\} = \frac{1}{100} \left\{ 4 - 1\cdot 2041200 - 3 \right\}$$

$$= -\cdot 0020412 = \bar{1}\cdot 9979588.$$

$$(b) \log (1\cdot 04)^{6000} = 6000 \log 1\cdot 04 = 6000 \times \cdot 0170333$$

$$= 102\cdot 1998000; \therefore \text{number of digits is } 103.$$

$$12. (a) \log 5 = \frac{1}{2} \log 25 = \cdot 6989700$$

$$\log 4 = 2 - \log 25 = \cdot 6020600$$

$$\log 51\cdot 5 = \log 5 + \log 10\cdot 3 = \cdot 6989700 + 1\cdot 0128372 = 1\cdot 7118072$$

$$\log (064)^{\frac{1}{100}} = \frac{1}{100} \left\{ \log 64 - \log 1000 \right\} = \frac{1}{100} \left\{ 3 \log 4 - 3 \right\}$$

$$= \frac{1}{100} \left\{ 1\cdot 8061800 - 3 \right\} = -\cdot 0119382 = \bar{1}\cdot 9880618.$$

$$(b) \log (1\cdot 03)^{600} = 600 \log 1\cdot 03 = 600 \times \cdot 0128372$$

$$= 7\cdot 7023200; \therefore \text{number of digits is } 8.$$

$$\begin{aligned}
 13. \log 7623 &= \log (9 \times 121 \times 7) = 2 \log 3 + 2 \log 11 + \log 7 \\
 &= .9542426 + 2.0827854 + .8450980 = 3.8821260 \\
 \log \frac{77}{300} &= \log 7 + \log 11 - \log 3 - \log 100 \\
 &= .8450980 + 1.0413927 - .4771213 - 2 = \bar{1}.4093694 \\
 \log \frac{3}{539} &= \log 3 - \log 11 - 2 \log 7 \\
 &= .4771213 - 1.0413927 - 1.6901960 = \bar{3}.7455326.
 \end{aligned}$$

$$\begin{aligned}
 14. \quad (1) \quad & x \log 4096 = \log 8 - x \log 64 \\
 & 4x \log 8 = \log 8 - 2x \log 8 \\
 & 4x = 1 - 2x; 6x = 1; x = \frac{1}{6}.
 \end{aligned}$$

$$(2) \quad (2.5)^x = 6.25 = (2.5)^2; \therefore x = 2.$$

$$\begin{aligned}
 (3) \quad & (ab)^x = m; x \log (ab) = \log m; \\
 & \therefore x = \frac{\log m}{\log a + \log b}.
 \end{aligned}$$

$$\begin{aligned}
 (4) \quad & x(m \log a + 2 \log b) = \log c; \\
 & \therefore x = \frac{\log c}{m \log a + 2 \log b}.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad & 3x \log a + (4-x) \log b = (2x-1) \log c \\
 & x(3 \log a - \log b - 2 \log c) = -4 \log b - \log c; \\
 & \therefore x = \frac{4 \log b + \log c}{2 \log c + \log b - 3 \log a}.
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad & x(\log a + m \log b) = \log c - 3x \log c \\
 & x(\log a + m \log b + 3 \log c) = \log c; \\
 & \therefore x = \frac{\log c}{\log a + m \log b + 3 \log c}.
 \end{aligned}$$

EXAMPLES—XLI. (p. 127).

$$(1) \quad \log 525030 = 5.7201841$$

$$\log 525020 = 5.7201758$$

$$\text{Difference for } 10 = .0000083$$

$$\therefore 10 : 5 = .0000083 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000041 ;$$

$$\therefore \log 52502.5 = 4.7201799.$$

$$(2) \quad \log 300430 = 5.4777433$$

$$\log 300420 = 5.4777288$$

$$\text{Difference for } 10 = .0000145$$

$$\therefore 10 : 5 = .0000145 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000072 ;$$

$$\therefore \log 300.425 = 2.4777360.$$

$$(3) \quad \log 32026000 = 7.5055027$$

$$\log 32025000 = 7.5054891$$

$$\text{Difference for } 1000 = .0000136$$

$$\therefore 1000 : 613 = .0000136 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000083 ;$$

$$\therefore \log 32.025613 = 1.5054974.$$

$$(4) \quad \log 236610 = 5.3740331$$

$$\log 236600 = 5.3740147$$

$$\text{Difference for } 10 = .0000184$$

$$\therefore 10 : 1 = .0000184 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000018 ;$$

$$\therefore \log 236.601 = 2.3740165.$$

$$(5) \quad \log 675030 = 5.8293231$$

$$\log 675020 = 5.8293166$$

$$\text{Difference for } 10 = .0000065$$

$$\therefore 10:1 = .0000065 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000007 \text{ (see end of Art. 162) ;}$$

$$\therefore \log 675021 = 1.8293173.$$

$$(6) \quad \log 7333600 = 6.8653172$$

$$\log 7333500 = 6.8653113$$

$$\text{Difference for } 100 = .0000059$$

$$\therefore 100:33 = .0000059 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000019 ;$$

$$\therefore \log 7333533 = 6.8653132.$$

$$(7) \quad \log 6593200 = 6.8190962$$

$$\log 6593100 = 6.8190897$$

$$\text{Difference for } 100 = .0000065$$

$$\therefore 100:71 = .0000065 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000046 ;$$

$$\therefore \log 6593171 = 6.8190943.$$

$$(8) \quad \log 340780 = 5.5324741$$

$$\log 340770 = 5.5324614$$

$$\text{Difference for } 10 = .0000127$$

$$\therefore 10:8 = .0000127 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000102 ;$$

$$\therefore \log 340778 = 5.5324716.$$

$$(9) \quad \log 390980 = 5.5921545$$

$$\log 390970 = 5.5921434$$

$$\text{Difference for } 10 = .0000111$$

$$\therefore 10:4 = .0000111 : \text{what we must add ;}$$

$$\therefore \text{we must add } .0000044 ;$$

$$\therefore \log 390974 = 5.5921478.$$

$$(10) \quad \log 2582000 = 6.4119562$$

$$\log 2581900 = 6.4119394$$

Difference for 100 = .0000168

$\therefore 100 : 26 :: .0000168 : \text{what we must add ;}$

$\therefore \text{we must add } .0000044 ;$

$\therefore \log 2581926 = 6.4119438.$

EXAMPLES—XLII. (p. 129).

$$(1) \quad \log 12955 = 4.1124374$$

$$\log 12954 = 4.1124039$$

Difference for 1 = .0000335

$\therefore .0000335 : .0000271 = 1 : \text{what has to be added ;}$

$\therefore \text{we must add } .8 ;$

$\therefore 4.112431 \text{ is the logarithm of } 12954.8.$

$$(2) \quad \log 46246 = 4.6650742$$

$$\log 46245 = 4.6650648$$

Difference for 1 = .0000094

$\therefore .0000094 : .0000009 = 1 : \text{what has to be added ;}$

$\therefore \text{we must add } .0957 \dots, \text{ or, } .096 ;$

$\therefore 4.6650657 \text{ is the logarithm of } 46245.096.$

$$(3) \quad \log 34573 = 4.5387371$$

$$\log 34572 = 4.5387245$$

Difference for 1 = .0000126

$\therefore .0000126 : .0000114 = 1 : \text{what we must add ;}$

$\therefore \text{we must add } .9047 \dots, \text{ or, } .91 ;$

$\therefore 4.5387359 \text{ is the logarithm of } 34572.91.$

$$\begin{aligned}
 (4) \quad & \log 39376 = 4.5952316 \\
 & \log 39375 = 4.5952206 \\
 & \text{Difference for } 1 = .0000110 \\
 \therefore .0000110 : .0000076 = 1 : & \text{what we must add;} \\
 \therefore & \text{we must add } .69; \\
 \therefore 5.5952282 & \text{ is the logarithm of } 393756.9.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad & \log 37160 = 4.5700757 \\
 & \log 37159 = 4.5700640 \\
 & \text{Difference for } 1 = .0000117 \\
 \therefore .0000117 : .0000062 = 1 : & \text{what we must add;} \\
 \therefore & \text{we must add } .529, \text{ or, } .53; \\
 \therefore 3.5700702 & \text{ is the logarithm of } 3715.953.
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad & \log 96462 = 4.9843563 \\
 & \log 96461 = 4.9843518 \\
 & \text{Difference for } 1 = .0000045 \\
 \therefore .0000045 : .0000024 = 1 : & \text{what we must add;} \\
 \therefore & \text{we must add } .53; \\
 \therefore 3.9843542 & \text{ is the logarithm of } .009646153.
 \end{aligned}$$

$$\begin{aligned}
 (7) \quad & \log 25726 = 4.4103723 \\
 & \log 25725 = 4.4103554 \\
 & \text{Difference for } 1 = .0000169 \\
 \therefore .0000169 : .0000166 = 1 : & \text{what must be added;} \\
 \therefore & \text{we must add } .982; \\
 \therefore 7.4103720 & \text{ is the logarithm of } .00000025725982.
 \end{aligned}$$

$$\begin{aligned}
 (8) \quad & \log 60196 = 4.7795604 \\
 & \log 60195 = 4.7795532 \\
 & \text{Difference for } 1 = .0000072 \\
 \therefore .0000072 : .0000029 = 1 : & \text{what must be added} \\
 \therefore & \text{we must add } .4027, \text{ or, } .403; \\
 \therefore 2.7795561 & \text{ is the logarithm of } 601.95403.
 \end{aligned}$$



$$(9) \quad \log 10906 = 4.0376655$$

$$\log 10905 = 4.0376257$$

Difference for 1 = .0000398

$\therefore .0000398 : .0000114 = 1$  : what must be added ;

$\therefore$  we must add .286 ;

$\therefore 3.0376371$  is the logarithm of 1090.5286.

$$(10) \quad \log 26202 = 4.4183344$$

$$\log 26201 = 4.4183179$$

Difference for 1 = .0000165

$\therefore .0000165 : .0000135 = 1$  : what must be added ;

$\therefore$  we must add .818 ;

$\therefore 2.4183314$  is the logarithm of 262.01818.

#### EXAMPLES—XLIII. (p. 132).

$$(1) \quad \sin 42^\circ.16' = .6725821$$

$$\sin 42^\circ.15' = .6723668$$

Difference for 1' = .0002153

$\therefore 60'' : 16'' = .0002153$  : what we must *add* ;

$\therefore$  we must add .0000574 ;

$\therefore \sin 42^\circ.15'.16'' = .6724242.$

$$(2) \quad \sin 72^\circ.15' = .9523958$$

$$\sin 72^\circ.14' = .9523071$$

Difference for 1' = .0000887

$\therefore 60'' : 6'' = .0000887$  : what we must *add* ;

$\therefore$  we must add .0000088 ;

$\therefore \sin 72^\circ.14'.6'' = .9523159.$



$$(3) \quad \sin 54^{\circ}.36' = .8151278$$

$$\sin 54^{\circ}.35' = .8149593$$

Difference for  $1' = .0001685$

$\therefore 60'' : 45'' = .0001685$  : what we must *add*;

$\therefore$  we must add  $.0001263$ ;

$$\therefore \sin 54^{\circ}.35'.45'' = .8150856.$$

$$(4) \quad \sin 87^{\circ}.27' = .9990098$$

$$\sin 87^{\circ}.26' = .9989968$$

Difference for  $1' = .0000130$

$\therefore 60'' : 15'' = .0000130$  : what we must *add*;

$\therefore$  we must add  $.0000032$ ;

$$\therefore \sin 87^{\circ}.26'.15'' = .9990000.$$

$$(5) \quad \sin 43^{\circ}.15' = .6851830$$

$$\sin 43^{\circ}.14' = .6849711$$

Difference for  $1' = .0002119$

$\therefore 60'' : 20'' = .0002119$  : what we must *add*;

$\therefore$  we must add  $.0000706$ ;

$$\therefore \sin 43^{\circ}.14'.20'' = .6850417.$$

$$(6) \quad \cos 41^{\circ}.13' = .7522233$$

$$\cos 41^{\circ}.14' = .7520316$$

Difference for  $1' = .0001917$

$\therefore 60'' : 26'' = .0001917$  : what we must *subtract*;

$\therefore$  we must subtract  $.0000830$ ;

$$\therefore \cos 41^{\circ}.13'.26'' = .7521403.$$

$$(7) \quad \tan 1^{\circ}.23' = .0241484$$

$$\tan 1^{\circ}.22' = .0238573$$

Difference for  $1' = .0002911$

$\therefore 60'' : 30'' = .0002911$  : what we must *add*;

$\therefore$  we must add  $.0001455$ ;

$$\therefore \tan 1^{\circ}.22'.30'' = .0240028.$$

$$\begin{array}{r} (8) \quad \cot 35^{\circ}. 6' = 1.4228561 \\ \cot 35^{\circ}. 7' = 1.4219766 \end{array}$$

Difference for  $1' = .0008795$

$$\begin{array}{l} \therefore 60'' : 23'' = .0008795 : \text{what we must subtract;} \\ \therefore \text{we must subtract } .0003371; \\ \therefore \cot 35^{\circ}. 6'. 23'' = 1.4225190. \end{array}$$

$$\begin{array}{r} (9) \quad \sin 67^{\circ}. 23' = .9230984 \\ \sin 67^{\circ}. 22' = .9229865 \end{array}$$

Difference for  $1' = .0001119$

$$\begin{array}{l} \therefore 60'' : 48''.5 = .0001119 : \text{what we must add;} \\ \therefore \text{we must add } .0000904; \\ \therefore \sin 67^{\circ}. 22'. 48''.5 = .9230769. \end{array}$$

$$\begin{array}{r} (10) \quad \cos 34^{\circ}. 12' = .8270806 \\ \cos 34^{\circ}. 13' = .8269170 \end{array}$$

Difference for  $1' = .0001636$

$$\begin{array}{l} \therefore 60'' : 19''.6 = .0001636 : \text{what we must subtract;} \\ \therefore \text{we must subtract } .0000534; \\ \therefore \cos 34^{\circ}. 12'. 19''.6 = .8270272. \end{array}$$

#### EXAMPLES—XLIV. (p. 135).

$$\begin{array}{r} (1) \quad \sin 48^{\circ}. 47' = .7522233 \\ \sin 48^{\circ}. 46' = .7520316 \end{array}$$

Difference for  $1' = .0001917$

$$\begin{array}{l} \therefore .0001917 : .0001084 = 60'' : \text{what we must add to } 48^{\circ}. 46'; \\ \therefore \text{we must add } 34''; \\ \therefore \text{the angle is } 48^{\circ}. 46'. 34''. \end{array}$$

$$(2) \quad \cos 2^{\circ}.33' = \cdot 9990098$$

$$\cos 2^{\circ}.34' = \cdot 9989968$$

Difference for  $1' = \cdot 0000130$

$\therefore \cdot 0000130 : \cdot 0000098 = 60''$  : what we must add to  $2^{\circ}.33'$ ;

$\therefore$  we must add  $45''$ ;

$\therefore$  the angle is  $2^{\circ}.33'.45''$ .

$$(3) \quad \sin 43^{\circ}.15' = \cdot 6851830$$

$$\sin 43^{\circ}.14' = \cdot 6849711$$

Difference for  $1' = \cdot 0002119$

$\therefore \cdot 0002119 : \cdot 0000289 = 60''$  : what we must add to  $43^{\circ}.14'$ ;

$\therefore$  we must add  $8''.18$ ;

$\therefore$  the angle is  $43^{\circ}.14'.8''.18$ .

$$(4) \quad \cos 32^{\circ}.31' = \cdot 8432351$$

$$\cos 32^{\circ}.32' = \cdot 8430787$$

Difference for  $1' = \cdot 0001564$

$\therefore \cdot 0001564 : \cdot 0000351 = 60''$  : what we must add to  $32^{\circ}.31'$ ;

$\therefore$  we must add  $13''.46$ , or, approximately,  $13''.5$ ;

$\therefore$  the angle is  $32^{\circ}.31'.13''.5$ .

$$(5) \quad \sin 24^{\circ}.12' = \cdot 4099230$$

$$\sin 24^{\circ}.11' = \cdot 4096577$$

Difference for  $1' = \cdot 0002653$

$\therefore \cdot 0002653 : \cdot 0000982 = 60''$  : what we must add to  $24^{\circ}.11'$ ;

$\therefore$  we must add  $22''.2$ ;

$\therefore$  the angle is  $24^{\circ}.11'.22''.2$ .

$$(6) \quad \sec 82^{\circ}.23' = 7.552169$$

$$\sec 82^{\circ}.22' = 7.528249$$

Difference for  $1' = \cdot 023920$

$\therefore \cdot 023920 : \cdot 005084 = 60''$  : what we must add to  $82^{\circ}.22'$ ;

$\therefore$  we must add  $12''.8$  nearly;

$\therefore$  the angle is  $82^{\circ}.22'.12''.8$ .

$$(7) \quad \begin{array}{l} \cos 53^{\circ}.7' = .6001876 \\ \cos 53^{\circ}.8' = .5999549 \end{array}$$

Difference for  $1' = .0002327$

$$\begin{aligned} \therefore .0002327 : .0001876 = 60'' : \text{what we must add to } 53^{\circ}.7'; \\ \therefore \text{we must add } 48''.4 \text{ nearly;} \\ \therefore \text{the angle is } 53^{\circ}.7'.48''.4. \end{aligned}$$

$$(8) \quad \begin{array}{l} \operatorname{cosec} 25^{\circ}.3' = 2.36179 \\ \operatorname{cosec} 25^{\circ}.4' = 2.36029 \end{array}$$

Difference for  $1' = .00150$

$$\begin{aligned} \therefore .00150 : .00068 = 60'' : \text{what we must add to } 25^{\circ}.3'; \\ \therefore \text{we must add } 27''.2; \\ \therefore \text{the angle is } 25^{\circ}.3'.27''.2. \end{aligned}$$

$$(9) \quad \begin{array}{l} \sin 73^{\circ}.45' = .9600499 \\ \sin 73^{\circ}.44' = .9599684 \end{array}$$

Difference for  $1' = .0000815$

$$\begin{aligned} \therefore .0000815 : .0000316 = 60'' : \text{what we must add to } 73^{\circ}.44'; \\ \therefore \text{we must add } 23''.2; \\ \therefore \text{the angle is } 73^{\circ}.44'.23''.2. \end{aligned}$$

$$(10) \quad \begin{array}{l} \tan 77^{\circ}.20' = 4.44942 \\ \tan 77^{\circ}.19' = 4.44338 \end{array}$$

Difference for  $1' = .00604$

$$\begin{aligned} \therefore .00604 : .00106 = 60'' : \text{what we must add to } 77^{\circ}.19'; \\ \therefore \text{we must add } 10''.5; \\ \therefore \text{the angle is } 77^{\circ}.19'.10''.5. \end{aligned}$$

#### EXAMPLES—XLV. (p. 138).

$$(1) \quad \begin{array}{l} L \sin 55^{\circ}.34' = 9.9163406 \\ L \sin 55^{\circ}.33' = 9.9162539 \end{array}$$

Difference for  $1' = .0000867$

$$\begin{aligned} \therefore 60'' : 54'' = .0000867 : \text{what we have to add;} \\ \therefore \text{we must add } .0000780; \\ \therefore L \sin 55^{\circ}.33'.54'' = 9.9163319. \end{aligned}$$

$$\begin{aligned}
 (2) \quad & L \sin 29^\circ. 26' = 9.6914445 \\
 & L \sin 29^\circ. 25' = 9.6912205 \\
 & \text{Difference for } 1' = .0002240 \\
 \therefore 60'' : 2'' &= .0002240 : \text{what we have to add;} \\
 \therefore & \text{we must add } .0000075; \\
 \therefore L \sin 29^\circ. 25'. 2'' &= 9.6912280.
 \end{aligned}$$

$$\begin{aligned}
 (3) \quad & L \cos 37^\circ. 28' = 9.8996604 \\
 & L \cos 37^\circ. 29' = 9.8995636 \\
 & \text{Difference for } 1' = .0000968 \\
 \therefore 60'' : 36'' &= .0000968 : \text{what we have to subtract;} \\
 \therefore & \text{we must subtract } .0000581; \\
 \therefore L \cos 37^\circ. 28'. 36'' &= 9.8996023.
 \end{aligned}$$

$$\begin{aligned}
 (4) \quad & L \sin 54^\circ. 14' = 9.9092371 \\
 & L \sin 54^\circ. 13' = 9.9091461 \\
 & \text{Difference for } 1' = .0000910 \\
 \therefore 60'' : 19'' &= .0000910 : \text{what we have to add;} \\
 \therefore & \text{we must add } .0000288; \\
 \therefore L \sin 54^\circ. 13'. 19'' &= 9.9091749.
 \end{aligned}$$

$$\begin{aligned}
 (5) \quad & L \tan 27^\circ. 43' = 9.7204759 \\
 & L \tan 27^\circ. 42' = 9.7201690 \\
 & \text{Difference for } 1' = .0003069 \\
 \therefore 60'' : 34'' &= .0003069 : \text{what we have to add;} \\
 \therefore & \text{we must add } .0001739; \\
 \therefore L \tan 27^\circ. 42'. 34'' &= 9.7203429.
 \end{aligned}$$

$$\begin{aligned}
 (6) \quad & L \tan 5^\circ. 14' = 8.9618659 \\
 & L \tan 5^\circ. 13' = 8.9604728 \\
 & \text{Difference for } 1' = .0013931 \\
 \therefore 60'' : 23'' &= .0013931 : \text{what we have to add;} \\
 \therefore & \text{we must add } .0005340; \\
 \therefore L \tan 5^\circ. 13'. 23'' &= 8.9610068.
 \end{aligned}$$



$$\begin{array}{r}
 (7) \quad L \cot 3^{\circ}. 37' = 11.1992368 \\
 \quad \quad L \cot 3^{\circ}. 38' = 11.1972347 \\
 \text{Difference for } 1' = .0020021 \\
 \therefore 60'' : 50'' = .0020021 : \text{what we have to subtract;} \\
 \therefore \text{we must subtract } .0016684; \\
 \therefore L \cot 3^{\circ}. 37'. 50'' = 11.1975684.
 \end{array}$$

$$\begin{array}{r}
 (8) \quad L \sin 39^{\circ}. 26' = 9.8028968 \\
 \quad \quad L \sin 39^{\circ}. 25' = 9.8027431 \\
 \text{Difference for } 1' = .0001537 \\
 \therefore 60'' : 10'' = .0001537 : \text{what we have to add;} \\
 \therefore \text{we must add } .0000256; \\
 \therefore L \sin 39^{\circ}. 25'. 10'' = 9.8027687.
 \end{array}$$

$$\begin{array}{r}
 (9) \quad L \sin 70^{\circ}. 35' = 9.9745697 \\
 \quad \quad L \sin 70^{\circ}. 34' = 9.9745252 \\
 \text{Difference for } 1' = .0000445 \\
 \therefore 60'' : 17'' = .0000445 : \text{what we must add;} \\
 \therefore \text{we must add } .0000126; \\
 \therefore L \sin 70^{\circ}. 34'. 17'' = 9.9745378.
 \end{array}$$

$$\begin{array}{r}
 (10) \quad L \cos 88^{\circ}. 54' = 8.2832434 \\
 \quad \quad L \cos 88^{\circ}. 55' = 8.2766136 \\
 \text{Difference for } 1' = .0066298 \\
 \therefore 60'' : 16'' = .0066298 : \text{what we must subtract;} \\
 \therefore \text{we must subtract } .0017679; \\
 \therefore L \cos 88^{\circ}. 54'. 16'' = 8.2814755.
 \end{array}$$

## EXAMPLES—XLVI. (p. 140).

$$\begin{array}{r}
 (1) \quad L \sin 14^{\circ}. 25' = 9.3961499 \\
 \quad \quad L \sin 14^{\circ}. 24' = 9.3956581 \\
 \text{Difference for } 1' = .0004918 \\
 \therefore .0004918 : .0002868 = 60'' : \text{what we have to add;} \\
 \therefore \text{we must add } 35'' \text{ nearly;} \\
 \therefore \text{the angle is } 14^{\circ}. 24'. 35''.
 \end{array}$$

$$(2) \quad L \sin 54^{\circ}.14' = 9.9092371$$

$$L \sin 54^{\circ}.13' = 9.9091461$$

Difference for  $1' = .0000910$

$\therefore .0000910 : .0000299 = 60''$ : what we have to add;

$\therefore$  we must add  $19''$ ;

$\therefore$  the angle is  $54^{\circ}.13'.19''$ .

$$(3) \quad L \sin 71^{\circ}.41' = 9.9774191$$

$$L \sin 71^{\circ}.40' = 9.9773772$$

Difference for  $1' = .0000419$

$\therefore .0000419 : .0000125 = 60''$ : what we must add;

$\therefore$  we must add  $18''$  nearly;

$\therefore$  the angle is  $71^{\circ}.40'.18''$ .

$$(4) \quad L \cos 29^{\circ}.25' = 9.9400535$$

$$L \cos 29^{\circ}.26' = 9.9399823$$

Difference for  $1' = .0000712$

$\therefore .0000712 : .0000023 = 60''$ : what we must add;

$\therefore$  we must add  $2''$  nearly;

$\therefore$  the angle is  $29^{\circ}.25'.2''$ .

$$(6) \quad L \tan 30^{\circ}.51' = 9.7761947$$

$$L \tan 30^{\circ}.50' = 9.7759077$$

Difference for  $1' = .0002870$

$\therefore .0002870 : .0001320 = 60''$ : what we must add;

$\therefore$  we must add  $27''.6$  nearly;

$\therefore$  the angle is  $30^{\circ}.50'.27''.6$ .

$$(6) \quad L \cot 86^{\circ}.32' = 8.7823199$$

$$L \cot 86^{\circ}.33' = 8.7802218$$

Difference for  $1' = .0020981$

$\therefore .0020981 : .0008556 = 60''$ : what we must add;

$\therefore$  we must add  $24''.5$  nearly;

$\therefore$  the angle is  $86^{\circ}.32'.24''.5$ .



$$(7) \quad L \sin 24^{\circ}.9' = 9.6118580$$

$$L \sin 24^{\circ}.8' = 9.6115762$$

$$\text{Difference for } 1' = .0002818$$

$$\therefore .0002818 : .0002114 = 60'' : \text{what we must add ;}$$

$$\therefore \text{ we must add } 45'' ;$$

$$\therefore \text{ the angle is } 24^{\circ}.8'.45''.$$

$$(8) \quad L \tan 11^{\circ}.40' = 9.3148851$$

$$L \tan 11^{\circ}.39' = 9.3142468$$

$$\text{Difference for } 1' = .0006383$$

$$\therefore .0006383 : .0005543 = 60'' : \text{what we must add ;}$$

$$\therefore \text{ we must add } 52'' ;$$

$$\therefore \text{ the angle is } 11^{\circ}.39'.52''.$$

$$(9) \quad L \operatorname{cosec} 46^{\circ}.23' = 10.1402787$$

$$L \operatorname{cosec} 46^{\circ}.24' = 10.1401584$$

$$\text{Difference for } 1' = .0001203$$

$$\therefore .0001203 : .0000220 = 60'' : \text{what we must add ;}$$

$$\therefore \text{ we must add } 11'' \text{ nearly ;}$$

$$\therefore \text{ the angle is } 46^{\circ}.23'.11''.$$

$$(10) \quad L \sec 29^{\circ}.55' = 10.0621053$$

$$L \sec 29^{\circ}.54' = 10.0620326$$

$$\text{Difference for } 1' = .0000727$$

$$\therefore .0000727 : .0000359 = 60'' : \text{what we must add ;}$$

$$\therefore \text{ we must add } 29''.6 \text{ nearly ;}$$

$$\therefore \text{ the angle is } 29^{\circ}.54'.29''.6.$$

#### EXAMPLES—XLVII. (p. 149).

$$(1) \quad \sin(A+B) = \sin(180^{\circ} - C) = \sin C.$$

$$(2) \quad \cos(A+B) = \cos(180^{\circ} - C) = -\cos C.$$

$$(3) \quad \sin \frac{A+B}{2} = \sin \left( 90^{\circ} - \frac{C}{2} \right) = \cos \frac{C}{2}.$$

$$(4) \cos \frac{A+B}{2} = \cos \left( 90^\circ - \frac{C}{2} \right) = \sin \frac{C}{2}.$$

$$(5) \tan \frac{A+B}{2} = \tan \left( 90^\circ - \frac{C}{2} \right) = \cot \frac{C}{2}.$$

$$(6) \cot \frac{A+B}{2} = \cot \left( 90^\circ - \frac{C}{2} \right) = \tan \frac{C}{2}.$$

## EXAMPLES—XLVIII. (p. 150).

$$\begin{aligned} 1. \quad (1) \sin 2A + \sin 2B + \sin 2C &= 2 \sin(A+B) \cdot \cos(A-B) + \sin 2C \\ &= 2 \sin C \cdot \cos(A-B) + 2 \sin C \cdot \cos C \\ &= 2 \sin C \cdot \{ \cos(A-B) + \cos C \} \\ &= 2 \sin C \{ \cos(A-B) - \cos(A+B) \} \\ &= 2 \sin C \cdot (2 \sin A \cdot \sin B) \\ &= 4 \sin A \cdot \sin B \cdot \sin C. \end{aligned}$$

$$\begin{aligned} (2) \sin(-A+B+C) + \sin(A-B+C) + \sin(A+B-C) \\ &= 2 \sin C \cdot \cos(A-B) + \sin(A+B) \cdot \cos C - \cos(A+B) \cdot \sin C \\ &= 2 \sin C \cdot \cos(A-B) + \sin C \cdot \cos C + \cos C \cdot \sin C \\ &= 2 \sin C \cdot \{ \cos(A-B) + \cos C \} \\ &= 2 \sin C \cdot \{ \cos(A-B) - \cos(A+B) \} \\ &= 4 \sin A \cdot \sin B \cdot \sin C. \end{aligned}$$

$$\begin{aligned} (3) \quad \frac{\cot \frac{A}{2} + \cot \frac{C}{2}}{\cot \frac{B}{2} + \cot \frac{C}{2}} &= \frac{\frac{\cos \frac{A}{2} \cdot \sin \frac{C}{2} + \sin \frac{A}{2} \cdot \cos \frac{C}{2}}{\sin \frac{A}{2} \cdot \sin \frac{C}{2}}}{\frac{\cos \frac{B}{2} \cdot \sin \frac{C}{2} + \sin \frac{B}{2} \cdot \cos \frac{C}{2}}{\sin \frac{B}{2} \cdot \sin \frac{C}{2}}} = \frac{\sin \left( \frac{A}{2} + \frac{C}{2} \right) \cdot \sin \frac{B}{2}}{\sin \left( \frac{B}{2} + \frac{C}{2} \right) \cdot \sin \frac{A}{2}} \\ &= \frac{\cos \frac{B}{2} \cdot \sin \frac{B}{2}}{\cos \frac{A}{2} \cdot \sin \frac{A}{2}} = \frac{2 \cos \frac{B}{2} \cdot \sin \frac{B}{2}}{2 \cos \frac{A}{2} \cdot \sin \frac{A}{2}} = \frac{\sin B}{\sin A}. \end{aligned}$$

$$(4) \tan(A+B+C) = \tan 180^\circ = 0;$$

$$\therefore \frac{\tan A + \tan B + \tan C - \tan A \cdot \tan B \cdot \tan C}{1 - \tan A \cdot \tan B - \tan B \cdot \tan C - \tan C \cdot \tan A} = 0;$$

$$\therefore \tan A + \tan B + \tan C - \tan A \cdot \tan B \cdot \tan C = 0;$$

$$\therefore \tan A + \tan B + \tan C = \tan A \cdot \tan B \cdot \tan C.$$

(5) As in Example (4),

$$\tan A + \tan B + \tan C = \tan A \cdot \tan B \cdot \tan C,$$

and dividing both sides by  $\tan A \cdot \tan B \cdot \tan C$ ,

$$\cot B \cdot \cot C + \cot A \cdot \cot C + \cot A \cdot \cot B = 1.$$

$$(6) \cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \frac{\cos \frac{A}{2} \cdot \sin \frac{B}{2} + \sin \frac{A}{2} \cdot \cos \frac{B}{2}}{\sin \frac{A}{2} \cdot \sin \frac{B}{2}} + \frac{\cos \frac{C}{2}}{\sin \frac{C}{2}}.$$

$$= \frac{\sin \left( \frac{A}{2} + \frac{B}{2} \right)}{\sin \frac{A}{2} \cdot \sin \frac{B}{2}} + \frac{\cos \frac{C}{2}}{\sin \frac{C}{2}}$$

$$= \cos \frac{C}{2} \left\{ \frac{1}{\sin \frac{A}{2} \cdot \sin \frac{B}{2}} + \frac{1}{\sin \frac{C}{2}} \right\}$$

$$= \cos \frac{C}{2} \left\{ \frac{\sin \frac{C}{2} + \sin \frac{A}{2} \cdot \sin \frac{B}{2}}{\sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}} \right\}$$

$$= \cos \frac{C}{2} \left\{ \frac{\cos \left( \frac{A}{2} + \frac{B}{2} \right) + \sin \frac{A}{2} \cdot \sin \frac{B}{2}}{\sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}} \right\}$$

$$= \frac{\cos \frac{C}{2} \cdot \cos \frac{A}{2} \cdot \cos \frac{B}{2}}{\sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}} = \cot \frac{A}{2} \cdot \cot \frac{B}{2} \cdot \cot \frac{C}{2}.$$

(7)

$$\begin{aligned}
 1 + \cos 2A + \cos 2B + \cos 2C &= 1 + (2 \cos^2 A - 1) + 2 \cos(B+C) \cdot \cos(B-C) \\
 &= 2 \cos^2 A - 2 \cos A \cdot \cos(B-C) \\
 &= -2 \cos A \cdot \{\cos(B+C) + \cos(B-C)\} \\
 &= -2 \cos A \cdot 2 \cos B \cdot \cos C = -4 \cos A \cdot \cos B \cdot \cos C.
 \end{aligned}$$

$$\begin{aligned}
 (8) \quad \cos A + \cos B + \cos C &= 2 \cos \frac{A+B}{2} \cdot \cos \frac{A-B}{2} + 1 - 2 \sin^2 \frac{C}{2} \\
 &= 2 \sin \frac{C}{2} \cdot \cos \frac{A-B}{2} - 2 \sin^2 \frac{C}{2} + 1 \\
 &= 2 \sin \frac{C}{2} \left\{ \cos \frac{A-B}{2} - \cos \frac{A+B}{2} \right\} + 1 \\
 &= 2 \sin \frac{C}{2} \cdot 2 \sin \frac{A}{2} \cdot \sin \frac{B}{2} + 1 = 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} + 1.
 \end{aligned}$$

$$\begin{aligned}
 (9) \quad -\sin 2A + \sin 2B + \sin 2C &= 2 \sin(B+C) \cdot \cos(B-C) - 2 \sin A \cdot \cos A \\
 &= 2 \sin A \cdot \{\cos(B-C) - \cos A\} \\
 &= 2 \sin A \cdot \{\cos(B-C) + \cos(B+C)\} \\
 &= 4 \sin A \cdot \cos B \cdot \cos C.
 \end{aligned}$$

$$\begin{aligned}
 (10) \quad \sin A + \sin B - \sin C &= 2 \sin \frac{A+B}{2} \cdot \cos \frac{A-B}{2} - 2 \sin \frac{C}{2} \cdot \cos \frac{C}{2} \\
 &= 2 \cos \frac{C}{2} \cdot \left\{ \cos \frac{A-B}{2} - \sin \frac{C}{2} \right\} \\
 &= 2 \cos \frac{C}{2} \cdot \left\{ \cos \frac{A-B}{2} - \cos \frac{A+B}{2} \right\} \\
 &= 2 \cos \frac{C}{2} \cdot 2 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \\
 &= 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \cos \frac{C}{2}.
 \end{aligned}$$

$$\begin{aligned}
 (11) \quad \sin 2A + \sin 2B - \sin 2C &= 2 \sin(A+B) \cdot \cos(A-B) - 2 \sin C \cdot \cos C \\
 &= 2 \sin C \cdot \{\cos(A-B) - \cos C\} \\
 &= 2 \sin C \cdot \{\cos(A-B) + \cos(A+B)\} \\
 &= 4 \sin C \cdot \cos A \cdot \cos B.
 \end{aligned}$$

$$\begin{aligned}
 (12) \quad \cos A + \cos B - \cos C &= 2 \cos \frac{A+B}{2} \cdot \cos \frac{A-B}{2} - \left(1 - 2 \sin^2 \frac{C}{2}\right) \\
 &= 2 \sin \frac{C}{2} \cdot \cos \frac{A-B}{2} + 2 \sin \frac{C}{2} \cdot \cos \frac{A+B}{2} - 1 \\
 &= 2 \sin \frac{C}{2} \cdot \left\{ \cos \frac{A-B}{2} + \cos \frac{A+B}{2} \right\} - 1 \\
 &= 4 \sin \frac{C}{2} \cdot \cos \frac{A}{2} \cdot \cos \frac{B}{2} - 1.
 \end{aligned}$$

$$\begin{aligned}
 (13) \quad \cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} + \cos^2 \frac{C}{2} &= \frac{1}{2} \left\{ \cos A + 1 + \cos B + 1 + \cos C + 1 \right\} \\
 &= \frac{1}{2} \cdot \left\{ \cos A + \cos B + \cos C + 3 \right\} \\
 &= \frac{1}{2} \cdot \left\{ 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} + 1 + 3 \right\}, \text{ as in Ex. 8.} \\
 &= 2 + 2 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}.
 \end{aligned}$$

$$\begin{aligned}
 (14) \quad \sin^2 \frac{A}{2} + \sin^2 \frac{B}{2} + \sin^2 \frac{C}{2} &= \frac{1}{2} \cdot \left\{ 1 - \cos A + 1 - \cos B + 1 - \cos C \right\} \\
 &= \frac{1}{2} \cdot \left\{ 3 - 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} - 1 \right\}, \text{ as in Ex. 8.} \\
 &= 1 - 2 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}.
 \end{aligned}$$

2.

$$(1) \quad \frac{b+c}{a} = \cot A + \operatorname{cosec} A = \frac{\cos A + 1}{\sin A} = \frac{2 \cos^2 \frac{A}{2}}{2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}} = \frac{\cos \frac{A}{2}}{\sin \frac{A}{2}} = \cot \frac{A}{2}.$$

$$\begin{aligned}
 (2) \quad 2 \operatorname{cosec} 2A \cdot \cot B &= \frac{2}{\sin 2A} \cdot \frac{\cos B}{\sin B} = \frac{2 \cos B}{2 \sin A \cdot \cos A \cdot \sin B} \\
 &= \frac{\cos B}{\cos B \cdot \sin B \cdot \sin B} = \frac{1}{\sin^2 B} = \frac{c^2}{b^2}.
 \end{aligned}$$

$$(3) \quad 2 \sin^2 \frac{B}{2} = 1 - \cos B = 1 - \frac{a}{c} = \frac{c-a}{c};$$

$$\therefore \sin \frac{B}{2} = \sqrt{\left(\frac{c-a}{2c}\right)}.$$

$$(4) \quad 2 \cos^2 \frac{B}{2} = 1 + \cos B = 1 + \frac{a}{c} = \frac{a+c}{c};$$

$$\therefore \cos \frac{B}{2} = \sqrt{\left(\frac{a+c}{2c}\right)}.$$

$$\begin{aligned} (5) \quad \frac{\cos 2B - \cos 2A}{\sin 2A} &= \frac{\cos^2 B - \sin^2 B - \cos^2 A + \sin^2 A}{2 \sin A \cdot \cos A} \\ &= \frac{\sin^2 A - \sin^2 B - \sin^2 B + \sin^2 A}{2 \sin A \cdot \cos A} = \frac{2 \sin^2 A - 2 \sin^2 B}{2 \sin A \cdot \cos A} \\ &= \frac{\sin A}{\cos A} - \frac{\sin B}{\cos B} = \tan A - \tan B. \end{aligned}$$

$$\begin{aligned} (6) \quad \tan 2A - \sec 2B &= \frac{2 \tan A}{1 - \tan^2 A} - \frac{1}{\cos^2 B - \sin^2 B} \\ &= \frac{2ab}{b^2 - a^2} - \frac{c^2}{a^2 - b^2} = \frac{2ab + c^2}{b^2 - a^2} \\ &= \frac{2ab + a^2 + b^2}{b^2 - a^2} = \frac{b+a}{b-a}. \end{aligned}$$

$$\begin{aligned} (7) \quad (\sin A - \sin B)^2 + (\cos A + \cos B)^2 &= \sin^2 A - 2 \sin A \cdot \sin B + \sin^2 B + \cos^2 A + 2 \cos A \cdot \cos B + \cos^2 B \\ &= 2 + 2(\cos A \cdot \cos B - \sin A \cdot \sin B) \\ &= 2 + 2 \cos(A+B) = 2 - 2 \cos C = 4 \sin^2 \frac{C}{2}. \end{aligned}$$

$$(8) \quad \sec 2A = \frac{1}{\cos^2 A - \sin^2 A} = \frac{1}{b^2 - a^2} = \frac{c^2}{b^2 - a^2}.$$



$$(9) a^3 \cdot \cos A + b^3 \cdot \cos B = a^3 \cdot \frac{b}{c} + b^3 \cdot \frac{a}{c} = \frac{ab(a^2 + b^2)}{c} = \frac{abc^2}{c} = abc.$$

(10)

$$\begin{aligned} \cot(B-A) + \cot 2\left(A + \frac{C}{2}\right) &= \frac{\cos B \cdot \cos A + \sin B \cdot \sin A}{\sin B \cdot \cos A - \cos B \cdot \sin A} + \cot(2A + 90^\circ) \\ &= \frac{\sin A \cdot \sin B + \sin B \cdot \sin A}{\sin B \cdot \sin B - \sin A \cdot \sin A} - \tan 2A \\ &= \frac{2\sin A \sin B}{b^2 - a^2} = \frac{2 \tan A}{1 - \tan^2 A} = \frac{2ab}{b^2 - a^2} = \frac{2ab}{b^2 - a^2} = 0. \end{aligned}$$

$$3. (1) \frac{\sin A - \sin B}{a - b} = \frac{\frac{a \sin C}{c} - \frac{b \sin C}{c}}{a - b} = \frac{(a - b) \sin C}{(a - b)c} = \frac{\sin C}{c}.$$

$$(2) \frac{\sin(A-B)}{\sin C} = \frac{\sin(A-B) \cdot \sin(A+B)}{\sin C \cdot \sin C} = \frac{\sin^2 A - \sin^2 B}{\sin^2 C} = \frac{a^2 - b^2}{c^2}.$$

$$(3) \frac{a \cdot \sin C}{b - a \cos C} = \frac{a \sin C}{a \cos C + c \cos A - a \cos C} = \frac{a \cdot \sin C}{c \cdot \cos A} = \frac{c \cdot \sin A}{c \cdot \cos A} = \tan A.$$

$$\begin{aligned} (4) \frac{c}{a} \cdot \operatorname{cosec} B - \cot B &= \frac{c}{a \cdot \sin B} - \frac{\cos B}{\sin B} = \frac{c - a \cdot \cos B}{a \cdot \sin B} \\ &= \frac{b \cos A + a \cos B - a \cos B}{a \sin B} = \frac{b \cos A}{b \sin A} = \cot A. \end{aligned}$$

$$\begin{aligned} (5) a + b + c &= (b \cos C + c \cos B) + (a \cos C + c \cos A) + (a \cos B + b \cos A) \\ &= (a + b) \cos C + (a + c) \cos B + (b + c) \cos A. \end{aligned}$$

$$(6) \frac{a + b}{c} = \frac{\sin A + \sin B}{\sin C} = \frac{2 \sin \frac{A+B}{2} \cdot \cos \frac{A-B}{2}}{2 \sin \frac{C}{2} \cdot \cos \frac{C}{2}} = \frac{\cos \frac{A-B}{2}}{\sin \frac{C}{2}};$$

$$\therefore (a + b) \cdot \sin \frac{C}{2} = c \cdot \cos \frac{A-B}{2}.$$



$$(7) \frac{a-b}{c} = \frac{\sin A - \sin B}{\sin C} = \frac{2 \cos \frac{A+B}{2} \cdot \sin \frac{A-B}{2}}{2 \sin \frac{C}{2} \cdot \cos \frac{C}{2}} = \frac{\sin \frac{A-B}{2}}{\cos \frac{C}{2}};$$

$$\therefore (a-b) \cos \frac{C}{2} = c \cdot \sin \frac{A-B}{2}.$$

$$(8) \frac{\tan B}{\tan C} = \frac{\sin B \cdot \cos C}{\sin C \cdot \cos B} = \frac{b \cdot \left( \frac{a^2 + b^2 - c^2}{2ab} \right)}{c \cdot \left( \frac{a^2 + c^2 - b^2}{2ac} \right)} = \frac{a^2 + b^2 - c^2}{a^2 - b^2 + c^2}.$$

$$(9) c = a \cos B + b \cos A = a \cos B + \frac{a \sin B}{\sin A} \cdot \cos A = a(\cos B + \sin B \cdot \cot A).$$

$$(10) 2(ab \cdot \cos C + ac \cdot \cos B + bc \cdot \cos A) \\ = (a^2 + b^2 - c^2) + (a^2 + c^2 - b^2) + (b^2 + c^2 - a^2) = a^2 + b^2 + c^2.$$

$$(11) \cos^2 A + \cos^2 B + \cos^2 C + 2 \cos A \cdot \cos B \cdot \cos C \\ = \frac{1}{2} \left\{ 1 + \cos 2A + 1 + \cos 2B + 1 + \cos 2C + 4 \cos A \cdot \cos B \cdot \cos C \right\} \\ = \frac{1}{2} \left\{ 3 + (-1 - 4 \cos A \cdot \cos B \cdot \cos C) + 4 \cos A \cdot \cos B \cdot \cos C \right\},$$

by Example XLVIII. 1. (7).

$$= \frac{1}{2} \times 2 = 1.$$

$$(12) \frac{a-b}{c} \cdot 2 \cos^2 \frac{C}{2} = \frac{\sin A - \sin B}{\sin C} \cdot 2 \cos^2 \frac{C}{2} = \frac{2 \cos \frac{A+B}{2} \cdot \sin \frac{A-B}{2}}{\sin \frac{C}{2}} \cdot \cos \frac{C}{2} \\ = 2 \sin \frac{A-B}{2} \cdot \sin \frac{A+B}{2} = \cos B - \cos A.$$

$$(13) \frac{a+b}{c} \cdot 2 \sin^2 \frac{C}{2} = \frac{\sin A + \sin B}{\sin C} \cdot 2 \sin^2 \frac{C}{2} = \frac{2 \sin \frac{A+B}{2} \cdot \cos \frac{A-B}{2}}{\cos \frac{C}{2}} \cdot \sin \frac{C}{2} \\ = 2 \cos \frac{A-B}{2} \cdot \cos \frac{A+B}{2} = \cos A + \cos B.$$

$$(14) \quad a^2 \cdot \sin A + ab \cdot \sin B + ac \cdot \sin C = a^2 \sin A + b \cdot b \sin A + c \cdot c \sin A \\ = (a^2 + b^2 + c^2) \sin A.$$

(15) By Art. 184, page 149,

$$\cot \frac{A}{2} = \sqrt{\frac{s \cdot (s-a)}{(s-b)(s-c)}} \quad \text{and} \quad \cot \frac{B}{2} = \sqrt{\frac{s \cdot (s-b)}{(s-a)(s-c)}}; \\ \therefore \cot \frac{A}{2} : \cot \frac{B}{2} = s-a : s-b \\ = b+c-a : a+c-b.$$

(16)

$$\cot \frac{A}{2} \cdot \cot \frac{B}{2} = \sqrt{\frac{s \cdot (s-a)}{(s-b)(s-c)}} \cdot \sqrt{\frac{s \cdot (s-b)}{(s-a)(s-c)}} = \frac{s}{s-c} = \frac{a+b+c}{a+b-c}.$$

$$(17) \quad a \sin(B-C) + b \sin(C-A) + c \cdot \sin(A-B) \\ = a (\sin B \cdot \cos C - \cos B \cdot \sin C) + b (\sin C \cdot \cos A - \cos C \cdot \sin A) \\ \quad + c (\sin A \cdot \cos B - \cos A \cdot \sin B) \\ = \cos C (a \sin B - b \sin A) + \cos B (c \sin A - a \sin C) \\ \quad + \cos A (b \sin C - c \sin B) \\ = 0 + 0 + 0 = 0.$$

4. If the sides are in arithmetical progression, so also are the sines of the angles ;

$$\therefore \sin A + \sin C = 2 \sin B, \\ \text{or } \sin A + \sin(A+B) = 2 \sin B, \\ \text{or } 2 \sin \left( A + \frac{B}{2} \right) \cos \frac{B}{2} = 4 \sin \frac{B}{2} \cdot \cos \frac{B}{2}; \\ \therefore \sin \left( A + \frac{B}{2} \right) = 2 \sin \frac{B}{2}.$$

$$5. \quad (b+c) \cdot AD = b \cdot AD + c \cdot AD \\ = b \cdot b \sin C + c \cdot c \sin B \\ = b^2 \sin C + c^2 \sin B.$$




FIG. 20.

6. Let  $AB=4$ ,  $AC=9$ ,  $BC=12$ , and let  $AD$  be the line bisecting  $\angle BAC$ .




FIG. 21.

Then, by EUCLID VI. B,

$$BD \cdot DC + DA^2 = BA \cdot AC$$

$$AD \cdot \frac{\sin \frac{A}{2}}{\sin B} \times AD \cdot \frac{\sin \frac{A}{2}}{\sin C} + DA^2 = 36$$

$$AD^2 \left( \frac{\sin^2 \frac{A}{2}}{\sin B \cdot \sin C} + 1 \right) = 36$$

$$AD^2 \left\{ \frac{\frac{(s-b)(s-c)}{bc}}{\frac{4}{a^2 bc} \cdot s \cdot (s-a) \cdot (s-b) \cdot (s-c)} + 1 \right\} = 36$$

$$AD^2 \left\{ \frac{a^2}{4 \cdot s \cdot (s-a)} + 1 \right\} = 36$$

$$AD^2 \times \frac{169}{25} = 36, \text{ or, } AD = \frac{6 \times 5}{13} = 2\frac{4}{13}.$$

7. If  $\sin A = 2 \cos B \cdot \sin C$

$$\sin(B+C) = 2 \cos B \cdot \sin C$$

$$\sin B \cdot \cos C + \cos B \cdot \sin C = 2 \cos B \cdot \sin C$$

$$\sin B \cdot \cos C - \cos B \sin C = 0$$

$$\sin(B-C) = 0, \text{ and } \therefore B=C.$$

$$8. \text{ If } \cos A \cdot \cos B \cdot \sin C = \frac{\sin A + \sin B}{\cos A + \cos B} \\ \cos A \cdot \cos B$$

$$\sin C = \frac{\sin A + \sin B}{\cos A + \cos B} = \frac{2 \cdot \sin \frac{A+B}{2} \cdot \cos \frac{A-B}{2}}{2 \cdot \cos \frac{A+B}{2} \cdot \cos \frac{A-B}{2}};$$

$$\therefore 2 \sin \frac{C}{2} \cdot \cos \frac{C}{2} = \frac{\cos \frac{C}{2}}{\sin \frac{C}{2}};$$

$$\therefore \sin^2 \frac{C}{2} = \frac{1}{2}, \text{ or, } \sin \frac{C}{2} = \frac{1}{\sqrt{2}};$$

$$\therefore \frac{C}{2} = 45^\circ, \text{ and } \therefore C = 90^\circ.$$

$$9. \text{ If } \sin^2 A = \sin^2 B + \sin^2 C$$

$$\sin^2 A = \frac{b^2}{a^2} \cdot \sin^2 A + \frac{c^2}{a^2} \cdot \sin^2 A;$$

$$\therefore a^2 = b^2 + c^2, \text{ and } \therefore A = 90^\circ.$$

$$10. \text{ If } \frac{\sin A}{\sin C} = \frac{\sin C}{\sin B}, \text{ then } \frac{a}{c} = \frac{c}{b}, \text{ or, } ab = c^2.$$

$$\text{Then } \frac{a^3 + b^3 + c^3}{a + b + c} = ab$$

$$a^3 + b^3 + c^3 = ab(a + b) + abc \\ = ab(a + b) + c^3;$$

$$\therefore a^3 + b^3 = ab(a + b);$$

$$\therefore a^2 - ab + b^2 = ab, \text{ or, } (a - b)^2 = 0, \text{ or, } a = b.$$

Hence  $a, b, c$  are all equal.

$$11. \quad c^2 = a^2 + b^2 - 2ab \cdot \cos C$$

$$= a^2 + b^2 - 2ab \times \left(-\frac{1}{2}\right)$$

$$= a^2 + b^2 + ab.$$

$$12. \frac{\sin A}{\sin B} = \frac{a}{b};$$

$$\therefore \frac{\sin A + \sin B}{\sin A - \sin B} = \frac{a+b}{a-b};$$

$$\therefore \frac{\sin(B+C) + \sin B}{\sin(B+C) - \sin B} = \frac{a+b}{a-b};$$

$$\therefore \frac{\sin\left(B + \frac{C}{2}\right) \cdot \cos \frac{C}{2}}{\cos\left(B + \frac{C}{2}\right) \cdot \sin \frac{C}{2}} = \frac{a+b}{a-b}.$$

Now  $\angle ADC = B + \frac{C}{2}$ , by EUCLID I. 32

$$\therefore \tan ADC \cdot \cot \frac{C}{2} = \frac{a+b}{a-b};$$

$$\therefore \tan ADC = \frac{a+b}{a-b} \cdot \tan \frac{C}{2}.$$




FIG. 22.

(13) Draw  $CE$  perpendicular to  $AB$ .

Then by EUCLID II. XII. and XIII.

$$CB^2 = CD^2 + DB^2 + 2DB, DE,$$

$$CA^2 = CD^2 + DA^2 - 2AD, DE,$$

and  $DB = AD$ .




FIG. 23.

$$\therefore CB^2 + CA^2 = 2CD^2 + DB^2 + DA^2;$$

$$\therefore a^2 + b^2 = 2CD^2 + \frac{c^2}{4} + \frac{c^2}{4};$$

$$\therefore CD^2 = \frac{a^2}{2} + \frac{b^2}{2} - \frac{c^2}{4}.$$

### EXAMPLES—XLIX. (p. 157).

$$(1) \quad a = \sqrt{c^2 - b^2} = \sqrt{16} = 4$$

$$\sin A = \frac{a}{c} = \frac{4}{5} = .8.$$

Hence, as in Art. 168, we find  $A = 53^\circ. 7'. 48''. 4$ ;

and  $\therefore B = 36^\circ. 52'. 11''. 6$ .

$$(2) \quad a = \sqrt{c^2 - b^2} = \sqrt{64} = 8,$$

$$\sin A = \frac{a}{c} = \frac{8}{17} = .4705882.$$

Hence  $A = 28^\circ. 4'. 20''.9$ , and  $B = 61^\circ. 55'. 39''.1$ .

$$(3) \quad a = \sqrt{c^2 - b^2} = \sqrt{400} = 20,$$

$$\sin A = \frac{a}{c} = \frac{20}{29} = .6896552.$$

Hence  $A = 43^\circ. 36'. 10''.1$ , and  $B = 46^\circ. 23'. 49''.9$ .

$$(4) \quad a = \sqrt{c^2 - b^2} = \sqrt{576} = 24,$$

$$\cos A = \frac{b}{c} = \frac{7}{25} = .28.$$

Hence  $A = 73^\circ. 44'. 23''.3$ , and  $B = 16^\circ. 15'. 36''.7$ .

$$(5) \quad a = \sqrt{c^2 - b^2} = \sqrt{3136} = 56,$$

$$\cos A = \frac{b}{c} = \frac{33}{65} = .5076923.$$

$\therefore A = 59^\circ. 29'. 23''.2$ , and  $B = 30^\circ. 30'. 36''.8$ .

$$(6) \quad a = c \cdot \sin A = 13 \times .9230770 = 12 \text{ very nearly,}$$

$$b = \sqrt{c^2 - a^2} = \sqrt{25} = 5,$$

$$B = 22^\circ. 37'. 11''.5.$$

$$(7) \quad a = c \cdot \sin A = 41 \times .9756098 = 40 \text{ very nearly,}$$

$$b = \sqrt{c^2 - a^2} = \sqrt{81} = 9,$$

$$B = 12^\circ. 40'. 49''.4.$$

$$(8) \quad a = c \cdot \cos B = 73 \times .6575341 = 48 \text{ very nearly,}$$

$$b = \sqrt{c^2 - a^2} = \sqrt{3025} = 55,$$

$$A = 41^\circ. 6'. 43''.5.$$

$$(9) \quad a = c \cdot \cos B = 89 \times .4382021 = 39 \text{ very nearly,}$$

$$b = \sqrt{c^2 - a^2} = \sqrt{6400} = 80,$$

$$A = 25^\circ. 59'. 21''.2.$$

$$(10) \quad b = a \div \tan A = 40 \div 4.4444442 = 9 \text{ very nearly,}$$

$$c = \sqrt{a^2 + b^2} = \sqrt{1681} = 41,$$

$$B = 12^\circ. 40'. 49''. 4.$$

EXAMPLES—L. (p. 159).

- (1)  $b = \sqrt{c^2 - a^2} = \sqrt{289 \times 81} = 17 \times 9 = 153,$   
 $\sin A = \frac{a}{c},$   
 $L \sin A = 10 + 2.0170333 - 2.2671717 = 9.7498616;$   
 $\therefore A = 34^\circ. 12'. 19''. 6,$  and  $B = 55^\circ. 47'. 40''. 4.$
- (2)  $b = \sqrt{c^2 - a^2} = \sqrt{729 \times 121} = 27 \times 11 = 297,$   
 $\sin A = \frac{a}{c};$   
 $\therefore L \sin A = 10 + 2.4828736 - 2.6283889 = 9.8544847;$   
 $\therefore A = 45^\circ. 40'. 2''. 3,$  and  $B = 44^\circ. 19'. 57''. 7.$
- (3)  $b = \sqrt{c^2 - a^2} = \sqrt{1681 \times 1} = 41,$   
 $\sin A = \frac{a}{c};$   
 $\therefore L \sin A = 10 + 2.9242793 - 2.9247960 = 9.9994833;$   
 $\therefore A = 87^\circ. 12'. 20''. 3,$  and  $B = 2^\circ. 47'. 39''. 7.$
- (4)  $b = \sqrt{c^2 - a^2} = \sqrt{961 \times 289} = 31 \times 17 = 527,$   
 $\sin A = \frac{a}{c};$   
 $\therefore L \sin A = 10 + 2.5263393 - 2.7958800 = 9.7304593;$   
 $\therefore A = 32^\circ. 31'. 13''. 5,$  and  $B = 57^\circ. 28'. 46''. 5.$
- (5)  $b = \sqrt{c^2 - a^2} = \sqrt{2209 \times 9} = 47 \times 3 = 141,$   
 $\sin A = \frac{a}{c};$   
 $\therefore L \sin A = 10 + 3.0413927 - 3.0449315 = 9.9964612;$   
 $\therefore A = 82^\circ. 41'. 44'',$  and  $B = 7^\circ. 18'. 16''.$



$$\begin{aligned}
 (6) \quad a &= \sqrt{c^2 - b^2} = \sqrt{968 \times 578}; \\
 \therefore \log a &= \frac{1}{2} \{ \log 968 + \log 578 \} = 2.8739016; \\
 \therefore a &= 748, \text{ and } \cos A = \frac{b}{c}; \\
 \therefore L \cos A &= 10 + 2.2900346 - 2.8881795 = 9.4018551. \\
 \therefore A &= 75^\circ. 23'. 18''.5, \text{ and } B = 14^\circ. 36'. 41''.5.
 \end{aligned}$$

$$\begin{aligned}
 (7) \quad a &= \sqrt{c^2 - b^2} = \sqrt{1058 \times 512}; \\
 \therefore \log a &= \frac{1}{2} \{ \log 1058 + 9 \log 2 \} = 2.8668778; \\
 \therefore a &= 736, \text{ and } \cos A = \frac{b}{c}; \\
 \therefore L \cos A &= 10 + 2.4361626 - 2.8948697 = 9.5412929; \\
 \therefore A &= 69^\circ. 38'. 56''.3, \text{ and } B = 20^\circ. 21'. 3''.7.
 \end{aligned}$$

$$\begin{aligned}
 (8) \quad a &= \sqrt{c^2 - b^2} = \sqrt{1250 \times 32} = 200, \\
 \cos A &= \frac{b}{c}; \\
 \therefore L \cos A &= 10 + 2.7846173 - 2.8068580 = 9.9777593; \\
 \therefore A &= 18^\circ. 10'. 50'', \text{ and } B = 71^\circ. 49'. 10''.
 \end{aligned}$$

$$\begin{aligned}
 (9) \quad c &= \sqrt{a^2 + b^2} = \sqrt{76176 + 243049} = 565, \\
 \tan A &= \frac{a}{b}; \\
 \therefore L \tan A &= 10 + 2.4409091 - 2.6928469 = 9.7480622; \\
 \therefore A &= 29^\circ. 14'. 30''.3, \text{ and } B = 60^\circ. 45'. 29''.7.
 \end{aligned}$$

$$\begin{aligned}
 (10) \quad c &= \sqrt{a^2 + b^2} = \sqrt{156816 + 162409} = 565, \\
 \tan A &= \frac{a}{b}; \\
 \therefore L \tan A &= 10 + 2.5976952 - 2.6053050 = 9.9923902; \\
 \therefore A &= 44^\circ. 29'. 53'', \text{ and } B = 45^\circ. 30'. 7''.
 \end{aligned}$$

EXAMPLES—LI. (p. 161).

- (1)  $\frac{\text{Height of steeple in feet}}{220} = \tan 46^\circ. 30'$ , and if  $h$  be put for height of steeple,

$$\begin{aligned}\log h &= \log 220 + L \tan 46^\circ. 30' - 10 \\ &= 2.3424227 + .0227500 = 2.3651727 ; \\ \therefore h &= 231.835 \text{ feet.}\end{aligned}$$

- (2)  $\frac{BC}{AC} = \tan 25^\circ. 10'$ , and if  $h$  be the height of the tower in feet,

$$\begin{aligned}\frac{h}{200} &= \tan 25^\circ. 10' ; \\ \therefore \log h &= \log 200 + L \tan 25^\circ. 10' - 10 \\ &= \log 1000 - \log 5 + 9.6719628 - 10 \\ &= 3 - .6989700 + 9.6719628 - 10 \\ &= 1.9729928 ; \\ \therefore h &= 93.97 \text{ feet.}\end{aligned}$$

- (3)  $BC = 50$  feet ;  $\angle BAC = 45^\circ$  ;  $\angle BDC = 30^\circ$ .

Then  $AC = BC = 50$  feet.

$$\begin{aligned}(a) \quad AD &= CD - AC \\ &= BC \cdot \cot 30^\circ - 50 \\ &= 50.(\cot 30^\circ - 1) = 50.(\sqrt{3} - 1) \\ &= 50 \times .7320508 \dots \\ &= 36.6025 \dots \text{ feet.}\end{aligned}$$




FIG. 24.

$$(\beta) \quad AB = AC \cdot \sec 45^\circ = 50 \cdot \sqrt{2} = 50 \times 1.4142 \dots = 70.71 \dots \text{ feet.}$$

$$(\gamma) \quad BD = BC \cdot \operatorname{cosec} 30^\circ = 50 \times 2 = 100 \text{ feet.}$$

- (4) If  $h$  be the measure of the height in feet,

$$\begin{aligned}\frac{h}{140} &= \tan 54^\circ. 27' ; \\ \therefore h &= 140 \times 1.399364 = 195.910960 ; \\ \therefore \text{height is } 196 \text{ feet nearly.}\end{aligned}$$

(5) Let  $PC$  be the hill.




FIG. 25.

Then  $\angle PAC = 32^\circ. 14'$ , and  
 $\angle PBC = 63^\circ. 26'$ .

Then  $PC = BC \tan PBC$ ,  
 and  $PC = AC \cdot \tan PAC$ .

$$\therefore BC \cdot \tan PBC = AC \cdot \tan PAC ;$$

$$\therefore BC \times 1.998 = (500 + BC) \times .63,$$

whence  $BC = 230$  nearly.

$$\text{Hence } PC = 230 \times 1.998 = 459.54 \\ = 460 \text{ yards nearly.}$$

(6) Let  $\theta$  represent the sun's altitude.

$$\text{Then } \tan \theta = \frac{150}{75} = 2 ;$$

$$\therefore L \tan \theta = 10 + \log 2 = 10.3010200.$$

$$\text{Hence } \theta = 63^\circ. 26'. 6''.$$

(7) Let  $BC$  be the breadth of the river.



FIG. 26.

Then  $AC = BC \cdot \tan 60^\circ$ ,

and  $AC = CD \cdot \tan 50^\circ$ .

$$\therefore BC \cdot \tan 60^\circ = (40 + BC) \tan 50^\circ ;$$

$$\therefore BC \times \sqrt{3} = (40 + BC) \times 1.19 ;$$

$$\therefore BC \cdot (1.73 - 1.19) = 40 \times 1.19 ;$$

$$\therefore .54 BC = 47.6,$$

$$\text{and } \therefore BC = 88 \text{ yards nearly.}$$

(8) Let  $\theta$  be the angle of inclination.

$$\text{Then } \sin \theta = \frac{60}{109} = .55045.$$

$$\text{Hence } \theta = 33^\circ. 23'. 55''. 7.$$

(9) Let  $\theta$  be the angle of inclination.

$$\text{Then } \sin \theta = \frac{140}{221} = .6306306 ;$$

$$\therefore \theta = 39^\circ. 5'. 47''. 9.$$

(10) Let  $PC$  be the tower ;  $\angle PAC = 55^\circ$  ;  $\angle PBC = 48^\circ$ .

$$\text{Then } \frac{PA}{AB} = \frac{\sin 48^\circ}{\sin BPA},$$

$$\text{or } \frac{PA}{30} = \frac{\sin 48^\circ}{\sin 7^\circ};$$

$$\therefore PA = 30 \times \frac{\sin 48^\circ}{\sin 7^\circ},$$

$$\text{and } AC = PA \cdot \cos PAC = PA \cdot \sin 35^\circ.$$

Hence if  $b$  be the breadth of the river in feet,

$$b = 30 \times \sin 35^\circ \times \frac{\sin 48^\circ}{\sin 7^\circ}$$

$$\begin{aligned} \therefore \log b &= \log 30 + L \sin 35^\circ + L \sin 48^\circ - L \sin 7^\circ - 10 \\ &= 1.47712 + 9.75859 + 9.87107 - 9.08589 - 10 \\ &= 2.02089; \end{aligned}$$

$$\therefore b = 104.93 \text{ feet.}$$




FIG. 27.

(11) Let  $AB$  be the height of the house,  $BD$  the length,  $C$  the place of observation.

Then  $ABC$  and  $CBD$  are right angles.

Then  $BC = BD \cdot \cot BCD$ ,

and since  $\cos BCD = \frac{1}{\sqrt{5}}$ ,  $\cot BCD = \frac{1}{2}$  ;

$$\therefore BC = 150 \times \frac{1}{2} = 75 \text{ feet.}$$




FIG. 28.

Again,  $AB = BC \cdot \tan ACB$ ,

$$\text{and since } \sin ACB = \frac{3}{\sqrt{34}}, \tan ACB = \frac{3}{5};$$

$$\therefore AB = 75 \times \frac{3}{5} = 45 \text{ feet.}$$

(12) Making the same construction as in Example (11),

$$BC = AB \cdot \cot ACB = 45 \times \frac{5}{3} = 75 \text{ feet,}$$

$$\text{and } BD = BC \cdot \tan BCD = 75 \times 2 = 150 \text{ feet.}$$

EXAMPLES—LII. (p. 174).

$$(1) \cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{169 + 1600 - 1369}{1040} = \frac{5}{13};$$

$$\therefore \sin A = \frac{12}{13} = .9230769.$$

$$\text{Hence } A = 67^\circ. 22'. 48''.5.$$

$$(2) \cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{841 + 14400 - 10201}{6960} = \frac{63}{87};$$

$$\therefore \sin A = \frac{60}{87} = .6896552.$$

$$\text{Hence } A = 43^\circ. 36'. 10''.1.$$

$$(3) s = \frac{1}{2}(37 + 13 + 30) = 40;$$

$$\therefore \sin \frac{A}{2} = \sqrt{\frac{27 \times 10}{13 \times 30}} = \sqrt{\frac{9}{13}};$$

$$\therefore L \sin \frac{A}{2} = 10 + \frac{1}{2} \left\{ .9542425 - 1.1139434 \right\}$$

$$= 10 - .0798504 = 9.9201496.$$

$$\text{Hence } A = 112^\circ. 37'. 11''.5.$$

$$(4) s = \frac{1}{2}(409 + 241 + 600) = 625 ;$$

$$\begin{aligned}\therefore \sin A &= \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)} \\ &= \frac{2}{144600} \sqrt{625 \times 216 \times 384 \times 25} \\ &= \frac{2 \times 36000}{144600} = \frac{360}{723} ;\end{aligned}$$

$$\therefore L \sin A = 10 + 2.5563025 - 2.8591383 = 9.6971642.$$

$$\text{Hence } A = 29^{\circ}. 51'. 46''. 1.$$

$$2. \quad \frac{\tan \frac{C-A}{2}}{\tan \frac{C+A}{2}} = \frac{c-a}{c+a} ;$$

$$\therefore \tan \frac{C-A}{2} = \frac{c-a}{c+a} \cdot \cot \frac{B}{2}.$$

$$\text{Now } c-a = 1859 \text{ and } c+a = 13419 ;$$

$$\therefore L \tan \frac{C-A}{2} = \log(c-a) - \log(c+a) + L \cot \frac{B}{2} ;$$

$$\begin{aligned}\therefore L \tan \frac{C-A}{2} &= 3.26928 - 4.12772 + 10.40312 \\ &= 9.54468.\end{aligned}$$

$$\text{Hence } \frac{C-A}{2} = 19^{\circ}. 18'. 50''.$$

$$\text{Also } \frac{C+A}{2} = 68^{\circ}. 26'. 0'' ;$$

$$\therefore C = 87^{\circ}. 44'. 50'', \text{ and } A = 49^{\circ}. 7'. 10''.$$

$$3. \quad b = a \cdot \frac{\sin B}{\sin A} ;$$

$$\begin{aligned}\therefore \log b &= \log a + L \sin B - L \sin A \\ &= 1.7403627 + 9.9764927 - 9.8188779 \\ &= 1.8979775 ; \\ \therefore b &= 79.063.\end{aligned}$$

$$\begin{aligned}
 4. \quad b &= c \cdot \frac{\sin B}{\sin C}; \\
 \therefore \log b &= \log c + L \sin B - L \sin C \\
 &= 2.1613680 + 9.9982047 - 9.8183919 \\
 &= 2.3411808; \\
 \therefore b &= 219.37.
 \end{aligned}$$

$$\begin{aligned}
 5. \quad \sin A &= \sin B \cdot \frac{a}{b}; \\
 \therefore L \sin A &= L \sin B + \log a - \log b \\
 &= 9.7175280 + 2.7537623 - 2.5465269 \\
 &= 9.9247634.
 \end{aligned}$$

Hence *one* value of  $A$  is  $57^\circ. 14'. 21''$ .

And since  $a$  is greater than  $b$ ,  $A$  is greater than  $B$ , and we may have the same given parts in a triangle where  $A$  is the supplement of  $57^\circ. 14'. 21''$ , or  $122^\circ. 45'. 39''$ .

$$\begin{aligned}
 6. \quad \sin B &= \frac{b}{c} \cdot \sin C = \frac{16}{8} \cdot \sin 30^\circ = \frac{2}{1} \times \frac{1}{2} = 1; \\
 \therefore B &= 90^\circ, \text{ and the triangle is not ambiguous.}
 \end{aligned}$$

$$\begin{aligned}
 7. \quad \text{In the equilateral triangle } a &= b = c; \\
 \therefore \cos A &= \frac{a^2 + a^2 - a^2}{2a^2} = \frac{a^2}{2a^2} = \frac{1}{2}.
 \end{aligned}$$

$$8. \quad \text{Let } A = 60^\circ, \frac{b}{c} = \frac{19}{1}, \text{ and } \therefore \frac{b-c}{b+c} = \frac{18}{20} = \frac{9}{10}.$$

$$\begin{aligned}
 \text{Now } \tan \frac{B-C}{2} &= \frac{b-c}{b+c} \cdot \cot \frac{A}{2} \\
 &= \frac{9}{10} \times \frac{\sqrt{3}}{1} = \frac{3^2 \times 3^{\frac{1}{2}}}{10} = \frac{3^{\frac{5}{2}}}{10};
 \end{aligned}$$

$$\begin{aligned}
 \therefore L \tan \frac{B-C}{2} &= 10 + \frac{5}{2} \log 3 - \log 10 \\
 &= 10 + 1.1928032 - 1 \\
 &= 10.1928032;
 \end{aligned}$$

$$\therefore \frac{B-C}{2} = 57^\circ. 19'. 11'',$$

$$\text{and } \frac{B+C}{2} = 60^\circ. 0'. 0''.$$

$$\therefore B = 117^\circ. 19'. 11'', \text{ and } C = 2^\circ. 40'. 49''.$$



9. Let  $a, b, c$  denote the sides in order of the given values.

$$\text{Then } \cos A = \frac{b^2 + c^2 - a^2}{2bc} = \frac{6 + (1 + \sqrt{3})^2 - 4}{2(1 + \sqrt{3}) \cdot \sqrt{6}} = \frac{6 + 2\sqrt{3}}{2\sqrt{6} + 6\sqrt{2}} = \frac{1}{\sqrt{2}};$$

$$\therefore A = 45^\circ.$$

$$\text{Again, } \sin B = \frac{b}{a} \cdot \sin A = \frac{\sqrt{6}}{2} \cdot \frac{1}{\sqrt{2}} = \frac{\sqrt{3}}{2};$$

$$\therefore B = 60^\circ;$$

$$\text{and } \therefore C = 180^\circ - (60^\circ + 45^\circ) = 180^\circ - 105^\circ = 75^\circ.$$

10. Construct a diagram, as in Art. 213, fig. 2, but with  $A$  and  $B$  interchanged, because  $B$  is here to be the *smaller* angle.




FIG. 29.

Let  $c_1 = A_2B$ , and  $c_2 = A_1B$ .

Then  $c_1 = BD - A_2D = a \cos B - b \cdot \cos CA_2D$ ,

and  $c_2 = BD + A_1D = a \cos B + b \cdot \cos CA_2D$ ;

$$\begin{aligned} \therefore c_1 \cdot c_2 &= a^2 \cdot \cos^2 B - b^2 \cdot \cos^2 CA_2D \\ &= a^2 \cdot \cos^2 B - b^2 \cdot \cos^2 A \\ &= a^2 \cdot (1 - \sin^2 B) - b^2 \cdot (1 - \sin^2 A) \\ &= a^2 - b^2; \\ \therefore c_1 \cdot c_2 + b^2 &= a^2. \end{aligned}$$

11.

Let  $A = 64^\circ. 12'$ , and  $\frac{b}{c} = \frac{9}{7}$ .

$$\text{Then } \frac{b-c}{b+c} = \frac{9-7}{9+7} = \frac{2}{16} = \frac{1}{8}.$$

$$\begin{aligned} \text{And } \tan \frac{B-C}{2} &= \frac{b-c}{b+c} \cdot \cot \frac{A}{2} \\ &= \frac{1}{8} \cdot \cot 32^\circ. 6'. \end{aligned}$$

H

$$\begin{aligned}\therefore L \tan \frac{B-C}{2} &= \log 1 - \log 8 + L \cot 32^\circ.6' \\ &= 0 - 3 \log 2 + L \tan 57^\circ.54' \\ &= -90309 + 10.2025255 \\ &= 9.2994355.\end{aligned}$$

$$\text{Hence } \frac{B-C}{2} = 11^\circ.16'.10'',$$

$$\text{and } \frac{B+C}{2} = 57^\circ.54'.0'';$$

$$\therefore B = 69^\circ.10'.10'', \text{ and } C = 46^\circ.37'.50''.$$

$$12. \quad s = \frac{15}{2}, s-a = \frac{7}{2}, s-b = \frac{5}{2}, s-c = \frac{3}{2}.$$

$$\therefore \cos \frac{B}{2} = \sqrt{\frac{15 \times 5}{2 \cdot 2 \cdot 4 \cdot 6}} = \sqrt{\frac{25}{25}};$$

$$\begin{aligned}\therefore L \cos \frac{B}{2} &= 10 + \frac{1}{2} \left\{ 2 \log 5 - 5 \log 2 \right\} \\ &= 10 + \frac{1}{2} \left\{ 1.3979400 - 1.5051495 \right\} \\ &= 9.9463953.\end{aligned}$$

$$\text{Hence } \frac{B}{2} = 27^\circ.53'.8'', \text{ and } B = 55^\circ.46'.16''.$$

13.

$$\begin{aligned}\tan \frac{A-B}{2} &= \frac{a-b}{a+b} \cdot \cot \frac{C}{2} \\ &= \frac{70-35}{70+35} \cdot \cot \frac{C}{2} \\ &= \frac{1}{3} \cot 18^\circ.26'.6'';\end{aligned}$$

$$\begin{aligned}\therefore L \tan \frac{A-B}{2} &= \log 1 - \log 3 + L \cot 18^\circ.26'.6'' \\ &= 0 - .4771213 + 10.4771213 \\ &= 10;\end{aligned}$$

$$\therefore \frac{A-B}{2} = 45^\circ,$$

$$\text{and } \frac{A+B}{2} = 71^\circ.33'.54'';$$

$$\therefore A = 116^\circ.33'.54'', \text{ and } B = 26^\circ.33'.54''.$$

EXAMPLES—LIII. (p. 176).

- (1)  $c = \sqrt{a^2 + b^2} = \sqrt{16 + 9} = 5,$   
 $\sin A = \frac{4}{5} = \cdot 8.$   
 By the tables  $\sin 53^\circ. 7' = \cdot 7998593,$   
 $\sin 53^\circ. 8' = \cdot 8000338.$   
 Hence  $A = 53^\circ. 7'. 48''. 4,$  and  $B = 36^\circ. 52'. 11''. 6.$

$$a = \sqrt{c^2 - b^2} = 48,$$

$$\sin B = \frac{55}{73} = \cdot 7535068.$$

By the tables  $\sin 48^\circ. 53' = \cdot 7533721,$   
 $\sin 48^\circ. 54' = \cdot 7535634.$   
 Hence  $B = 48^\circ. 53'. 16''. 5,$  and  $A = 41^\circ. 6'. 43''. 6.$

- (3)  $c = \sqrt{a^2 + b^2} = 353,$   
 $\sin A = \frac{272}{353} = \cdot 7705382.$   
 By the tables  $\sin 50^\circ. 24' = \cdot 7705132$   
 $\sin 50^\circ. 25' = \cdot 7706986.$   
 Hence  $A = 50^\circ. 24'. 8''. 1,$  and  $B = 39^\circ. 35'. 51''. 9.$

- (4)  $a = \sqrt{c^2 - b^2} = 40,$   
 $\sin A = \frac{40}{401} = \cdot 0997506.$   
 By the tables  $\sin 5^\circ. 43' = \cdot 0996092,$   
 $\sin 5^\circ. 44' = \cdot 0998986.$   
 Hence  $A = 5^\circ. 43'. 29''. 3,$  and  $B = 84^\circ. 16'. 30''. 7.$

- (5)  $B = 79^\circ. 7'. 9''. 6.$   
 By the tables  $\sin 10^\circ. 52' = \cdot 1885241,$   
 $\sin 10^\circ. 53' = \cdot 1888098.$   
 Hence  $\sin 10^\circ. 52'. 50''. 4 = \cdot 1887639 ;$   
 $\therefore a = c \cdot \sin A = 445 \times \cdot 1887639 = 84,$   
 and  $b = \sqrt{c^2 - a^2} = 437.$

(6)

$$B = 43^{\circ}. 0'. 10''.3.$$

By the tables  $\sin 46^{\circ}. 59' = .7311553$ ,

$$\sin 47^{\circ}. 0' = .7313537.$$

Hence  $\sin 46^{\circ}. 59'. 49''.7 = .7313196$  ;

$$\therefore a = c \cdot \sin A = 629 \times .7313196 = 460,$$

$$\text{and } b = \sqrt{c^2 - a^2} = 429.$$

(7)

$$A = 38^{\circ}. 34'. 48''.3.$$

By the tables  $\sin 51^{\circ}. 25' = .7817019$ ,

$$\sin 51^{\circ}. 26' = .7818833.$$

Hence  $\sin 51^{\circ}. 25'. 11''.7 = .7817372$  ;

$$\therefore b = c \cdot \sin B = 449 \times .7817372 = 351,$$

$$\text{and } a = \sqrt{c^2 - b^2} = 280.$$

(8)

$$A = 31^{\circ}. 2'. 53''.6.$$

By the tables  $\sin 58^{\circ}. 57' = .8567175$ ,

$$\sin 58^{\circ}. 58' = .8568675.$$

Hence  $\sin 58^{\circ}. 57'. 6''.4 = .8567335$  ;

$$\therefore b = c \cdot \sin B = 349 \times .8567335 = 299,$$

$$\text{and } a = \sqrt{c^2 - b^2} = 180.$$

(9)

$$B = 23^{\circ}. 57'. 8''.$$

By the tables  $\tan 23^{\circ}. 57' = .4441834$ ,

$$\tan 23^{\circ}. 58' = .4445318.$$

Hence  $\tan 23^{\circ}. 57'. 8'' = .4442365$  ;

$$\therefore b = a \cdot \tan B = 520 \times .4442365 = 231,$$

$$\text{and } c = \sqrt{a^2 + b^2} = 569.$$

(10)

$$B = 3^{\circ}. 41'. 43''.$$

By the tables  $\tan 86^{\circ}. 18' = 15.463814$ ,

$$\tan 86^{\circ}. 19' = 15.533981.$$

Hence  $\tan 86^{\circ}. 18'. 17'' = 15.483694$  ;

$$\therefore a = b \cdot \tan A = 31 \times 15.483694 = 480,$$

$$\text{and } c = \sqrt{a^2 + b^2} = 481.$$

EXAMPLES—LIV. (p. 177).

(1)  $s=245$ ,  $s-a=48$ ,  $s-b=192$ ,  $s-c=5$ .

$$\text{Then } \tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s \cdot (s-a)}};$$

$$\begin{aligned} \therefore L \tan \frac{A}{2} &= 10 + \frac{1}{2} \left\{ \log 192 + \log 5 - \log 245 - \log 48 \right\} \\ &= 10 + \frac{1}{2} \left\{ 2.2833012 + .6989700 - 2.3891661 - 1.6812412 \right\} \\ &= 10 + \frac{1}{2} \left\{ 2.9822712 - 4.0704073 \right\} \\ &= 9.4559320. \end{aligned}$$

Hence  $\frac{A}{2} = 15^\circ. 56'. 43''.4$ , and  $\therefore A = 31^\circ. 53'. 26''.8$ .

By a similar method we may find  $B = 8^\circ. 10'. 16''.4$ ,  
and  $\therefore C = 139^\circ. 56'. 16''.8$ .

(2.)  $s=605$ ,  $s-a=96$ ,  $s-b=384$ ,  $s-c=125$ .

$$\begin{aligned} L \tan \frac{A}{2} &= 10 + \frac{1}{2} \left\{ \log 384 + \log 125 - \log 605 - \log 96 \right\} \\ &= 10 + \frac{1}{2} \left\{ 2.5843312 + 2.0969100 - 2.7817554 - 1.9822712 \right\} \\ &= 10 + \frac{1}{2} \left\{ 4.6812412 - 4.7640266 \right\} \\ &= 9.9586703. \end{aligned}$$

Hence  $\frac{A}{2} = 42^\circ. 16'. 25''.25$ , and  $\therefore A = 84^\circ. 32'. 50''.5$ .

By a similar method we find  $B = 25^\circ. 36'. 30''.7$ ,  
and  $\therefore C = 69^\circ. 50'. 38''.8$ .

$$(3) \quad s=680, s-a=147, s-b=363, s-c=170.$$

$$\begin{aligned} L \tan \frac{A}{2} &= 10 + \frac{1}{2} \left\{ \log 363 + \log 170 - \log 680 - \log 147 \right\} \\ &= 10 + \frac{1}{2} \left\{ 2.5599066 + 2.2304489 - 2.8325089 - 2.1673173 \right\} \\ &= 10 + \frac{1}{2} \left\{ 4.7903555 - 4.9998262 \right\} \\ &= 9.8952647. \end{aligned}$$

$$\text{Hence } \frac{A}{2} = 28^{\circ}. 9'. 26'', \text{ and } \therefore A = 76^{\circ}. 18'. 52''.$$

By a similar method we find  $B = 35^{\circ}. 18'. 0''. 9$ ,  
and  $\therefore C = 68^{\circ}. 23'. 7''. 1$ .

$$(4) \quad s=808, s-a=243, s-b=363, s-c=202.$$

$$\begin{aligned} L \tan \frac{A}{2} &= 10 + \frac{1}{2} \left\{ \log 363 + \log 202 - \log 808 - \log 243 \right\} \\ &= 10 + \frac{1}{2} \left\{ 2.5599066 + 2.3053514 - 2.9074114 - 2.3856063 \right\} \\ &= 10 + \frac{1}{2} \left\{ 4.8652580 - 5.2930177 \right\} \\ &= 9.7861202. \end{aligned}$$

$$\text{Hence } \frac{A}{2} = 31^{\circ}. 25'. 46''. 45, \text{ and } \therefore A = 62^{\circ}. 51'. 32''. 9.$$

By a similar method we find  $B = 44^{\circ}. 29'. 53''$ ,  
and  $\therefore C = 72^{\circ}. 38'. 34''. 1$ .

$$(5) \quad s=416, s-a=7, s-b=175, s-c=234.$$

$$\begin{aligned} L \tan \frac{A}{2} &= 10 + \frac{1}{2} \left\{ \log 175 + \log 234 - \log 416 - \log 7 \right\} \\ &= 10 + \frac{1}{2} \left\{ 2.2430380 + 2.3692159 - 2.6190933 - .8450980 \right\} \\ &= 10 + \frac{1}{2} \left\{ 4.6122539 - 3.4641913 \right\} \\ &= 10.5740313. \end{aligned}$$

$$\text{Hence } \frac{A}{2} = 75^{\circ}. 4'. 7'', \text{ and } \therefore A = 150^{\circ}. 8'. 14''.$$

By a similar method we can find  $B = 17^{\circ}. 3'. 41''. 5$ ,  
and  $\therefore C = 12^{\circ}. 48'. 4''. 5$ .



$$(6) \quad B = 180^\circ - (A + C) = 11^\circ. 25'. 16''.3,$$

$$a = b \cdot \frac{\sin A}{\sin B} = \frac{29 \times .6896550}{.1980199} = 101,$$

$$c = b \cdot \frac{\sin C}{\sin B} = \frac{29 \times .8193229}{.1980199} = 120.$$

$$(7) \quad B = 180^\circ - (A + C) = 39^\circ. 18'. 27''.5,$$

$$a = b \cdot \frac{\sin A}{\sin B} = \frac{149 \times .9395972}{.6338400} = 221,$$

$$c = b \cdot \frac{\sin C}{\sin B} = \frac{149 \times .9438490}{.6338400} = 222.$$

$$(8) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{72}{130} \cdot \cot 16^\circ. 5'. 26''.9,$$

$$L \tan \frac{A-B}{2} = \log 72 - \log 130 + L \cot 16^\circ. 5'. 26''.9$$

$$= 1.8573325 - 2.1139434 + 10.5399616$$

$$= 10.2833507.$$

$$\text{Hence } \frac{A-B}{2} = 62^\circ. 29'. 16''.8,$$

$$\text{and } \frac{A+B}{2} = 73^\circ. 54'. 33''.1;$$

$$\therefore A = 136^\circ. 23'. 49''.9, \text{ and } B = 11^\circ. 25'. 16''.3.$$

$$\text{Also, } c = \frac{a \cdot \sin C}{\sin A} = \frac{101 \times .5326047}{.6896550} = 78.$$

$$(9) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{360}{442} \cot 48^\circ. 28'. 40''.05,$$

$$L \tan \frac{A-B}{2} = \log 360 - \log 442 + L \cot 48^\circ. 28'. 40''.05$$

$$= 2.5563025 - 2.6454223 + 9.9471473$$

$$= 9.8580275.$$

$$\text{Hence } \frac{A-B}{2} = 35^{\circ}.47'.50''.65,$$

$$\text{and } \frac{A+B}{2} = 41^{\circ}.31'.19''.95;$$

$$\therefore A = 77^{\circ}.19'.10''.6, \text{ and } B = 5^{\circ}.43'.29''.2.$$

$$\text{Also, } c = \frac{a \cdot \sin C}{\sin A} = \frac{401 \times .9926403}{.9756097} = 408.$$

$$(10) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cdot \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{72}{370} \cot 15^{\circ}.20'.17''.5,$$

$$\begin{aligned} L \tan \frac{A-B}{2} &= \log 72 - \log 370 + L \cot 15^{\circ}.20'.17''.5 \\ &= 1.8573325 - 2.5682017 + 10.5617669 \\ &= 9.8508977. \end{aligned}$$

$$\text{Hence } \frac{A-B}{2} = 35^{\circ}.21'.15'',$$

$$\text{and } \frac{A+B}{2} = 74^{\circ}.39'.42''.5;$$

$$\therefore A = 110^{\circ}.0'.57''.5, \text{ and } B = 39^{\circ}.18'.27''.5.$$

$$\text{Also, } c = \frac{a \cdot \sin C}{\sin A} = \frac{221 \times .5101885}{.9395972} = 120.$$

$$(11) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cdot \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{48}{170} \cdot \cot 33^{\circ}.29'.42''.7,$$

$$\begin{aligned} L \tan \frac{A-B}{2} &= \log 48 - \log 170 + L \cot 33^{\circ}.29'.42''.7 \\ &= 1.6812412 - 2.2304489 + 10.1792962 \\ &= 9.6300885. \end{aligned}$$

$$\text{Hence } \frac{A-B}{2} = 23^{\circ}.6'.57''.3,$$

$$\text{and } \frac{A+B}{2} = 56^{\circ}.29'.42''.7;$$

$$\therefore A = 79^{\circ}.36'.40'', \text{ and } B = 33^{\circ}.23'.54''.6.$$

$$\text{Also, } c = \frac{a \cdot \sin C}{\sin A} = \frac{109 \times .9204413}{.9836064} = 102.$$

$$(12) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cdot \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{362}{528} \cdot \cot 43^{\circ}. 57'. 30'',$$

$$\begin{aligned} L \tan \frac{A-B}{2} &= \log 362 - \log 528 + L \cot 43^{\circ}. 57'. 30'' \\ &= 2.5587086 - 2.7226339 + 10.0157949 \\ &= 9.8518696. \end{aligned}$$

$$\text{Hence } \frac{A-B}{2} = 35^{\circ}. 24'. 46'',$$

$$\text{and } \frac{A+B}{2} = 46^{\circ}. 2'. 30'';$$

$$\therefore A = 81^{\circ}. 27'. 16'', \text{ and } B = 10^{\circ}. 37'. 44''.$$

$$\text{Also, } c = \frac{b \cdot \sin C}{\sin B} = \frac{83 \times .999390}{.1844460} = 450.$$

$$(13) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cdot \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{120}{338} \cdot \cot 65^{\circ}. 42'. 22'',$$

$$\begin{aligned} L \tan \frac{A-B}{2} &= \log 120 - \log 338 + L \cot 65^{\circ}. 42'. 22'' \\ &= 2.0791812 - 2.5289167 + 9.6545508 \\ &= 9.2048153. \end{aligned}$$

$$\text{Hence } \frac{A-B}{2} = 9^{\circ}. 6'. 16''6,$$

$$\text{and } \frac{A+B}{2} = 24^{\circ}. 17'. 38'';$$

$$\therefore A = 33^{\circ}. 23'. 54''6, \text{ and } B = 15^{\circ}. 11'. 21''4.$$

$$\text{Also, } c = \frac{b \cdot \sin C}{\sin B} = \frac{109 \times .7499700}{.2620086} = 312.$$

$$(14) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{72}{410} \cdot \cot 52^{\circ}. 1'. 55''.5,$$

$$\begin{aligned} L \tan \frac{A-B}{2} &= \log 72 - \log 410 + L \cot 52^{\circ}. 1'. 55''.5 \\ &= 1.8573325 - 2.6127839 + 9.8923085 \\ &= 9.1368571. \end{aligned}$$

$$\therefore \frac{A-B}{2} = 7^{\circ}. 48'. 12'',$$

$$\text{and } \frac{A+B}{2} = 37^{\circ}. 58'. 4''.5 ;$$

$$\therefore A = 45^{\circ}. 46'. 16''.5, \text{ and } B = 30^{\circ}. 9'. 52''.5.$$

$$\text{Also, } c = \frac{b \sin C}{\sin B} = \frac{169 \times .9900242}{.5024855} = 332.97.$$

$$(15) \quad \tan \frac{A-B}{2} = \frac{a-b}{a+b} \cot \frac{C}{2},$$

$$\tan \frac{A-B}{2} = \frac{72}{410} \cot 7^{\circ}. 41'. 18''.5,$$

$$\begin{aligned} L \tan \frac{A-B}{2} &= \log 72 - \log 410 + L \cot 7^{\circ}. 41'. 18''.5 \\ &= 1.8573325 - 2.6127839 + 10.8696637 \\ &= 10.1142123. \end{aligned}$$

$$\text{Hence } \frac{A-B}{2} = 52^{\circ}. 26'. 54''.1,$$

$$\text{and } \frac{A+B}{2} = 82^{\circ}. 18'. 41''.5 ;$$

$$\therefore A = 134^{\circ}. 45'. 36''.6, \text{ and } B = 29^{\circ}. 51'. 46''.4.$$

$$\text{Also, } c = \frac{b \cdot \sin C}{\sin B} = \frac{169 \times .2651681}{.4982927} = 90.$$

$$(16) \quad \sin B = \frac{b \cdot \sin A}{a} = \frac{37 \times \sin 18^\circ 55' 28''}{13};$$

$$\therefore L \sin B = \log 37 + L \sin 18^\circ 55' 28'' - \log 13$$

$$= 1.5682017 + 9.5109783 - 1.1139434$$

$$= 9.9652366;$$

$$\therefore B = 67^\circ 22' 48'' \cdot 1, \text{ or its supplement } 112^\circ 37' 11'' \cdot 9.$$

$$(17) \quad \sin B = \frac{b \cdot \sin A}{a} = \frac{565 \times \sin 44^\circ 29' 53''}{445};$$

$$\therefore L \sin B = \log 565 + L \sin 44^\circ 29' 53'' - \log 445$$

$$= 2.7520484 + 9.8456468 - 2.6483600$$

$$= 9.9493352;$$

$$\therefore B = 62^\circ 51' 32'' \cdot 9, \text{ or its supplement } 117^\circ 8' 27'' \cdot 1.$$

$$18) \quad \sin B = \frac{b \cdot \sin A}{a} = \frac{836.4 \times \sin 14^\circ 24' 25''}{212.5};$$

$$\therefore L \sin B = \log 836.4 + L \sin 14^\circ 24' 25'' - \log 212.5$$

$$= 2.9224140 + 9.3958630 - 2.3273589$$

$$= 9.9909181;$$

$$\therefore B = 78^\circ 19' 24'', \text{ or its supplement } 101^\circ 40' 36''.$$

$$(19) \quad \sin B = \frac{b \cdot \sin A}{a} = \frac{564.8 \times \sin 40^\circ 32' 16''}{379.5};$$

$$\therefore L \sin B = \log 564.8 + L \sin 40^\circ 32' 16'' - \log 379.5$$

$$= 2.7518947 + 9.8128794 - 2.5792118$$

$$= 9.9855623;$$

$$\therefore B = 75^\circ 18' 28'' \cdot 2, \text{ or its supplement } 104^\circ 41' 31'' \cdot 8.$$

$$(20) \quad \sin B = \frac{b \cdot \sin A}{a} = \frac{8032.29 \times \sin 71^\circ 3' 34''}{9459.31};$$

$$\therefore L \sin B = \log 8032.29 + L \sin 71^\circ 3' 34'' - \log 9459.31$$

$$= 3.9048393 + 9.9758256 - 3.9758594$$

$$= 9.9048055;$$

$$\therefore B = 53^\circ 26' 0'' \cdot 6.$$

## EXAMPLES—LV. (p. 181).

- (1) Let  $QP$  be the hill ;  $\angle QBP = 60^\circ$  ;  $\angle QAP = 45^\circ$ .




FIG. 30.

$$\begin{aligned}\text{Then } QP &= BP \cdot \tan 60^\circ \\ &= (AP - 100) \tan 60^\circ \\ &= (QP - 100) \cdot \sqrt{3} ;\end{aligned}$$

$$\therefore QP = \frac{100\sqrt{3}}{\sqrt{3}-1} = \frac{100\sqrt{3}(\sqrt{3}+1)}{3-1} = 150 + 50\sqrt{3} = 236.602 \dots \text{feet.}$$

- (2) Let  $F$  be the fort ;  $S_1$  and  $S_2$  the ships.

$$\begin{aligned}\text{Then } \angle FS_1S_2 &= 35^\circ. 14', \text{ and } \angle FS_2S_1 = 42^\circ. 12', \\ \text{and } \angle S_1FS_2 &= 180^\circ - 77^\circ. 26'\end{aligned}$$




FIG. 31.

$$\begin{aligned}\text{and } FS_1 &= S_1S_2 \cdot \frac{\sin FS_2S_1}{\sin S_1FS_2} \\ &= 1760 \cdot \frac{\sin 42^\circ. 12'}{\sin 77^\circ. 26'} \\ &= 1760 \times \frac{.671}{.976} = 1210 \text{ yards,}\end{aligned}$$

$$\text{and } FS_2 = 1760 \times \frac{\sin 35^\circ. 14'}{\sin 77^\circ. 26'} = 1760 \times \frac{.577}{.976} = 1040.5 \text{ yards.}$$

- (3) With a construction similar to that in Example (2),

$$FS_1 = 880 \cdot \frac{\sin 85^\circ. 15'}{\sin 11^\circ} = 880 \times \frac{.9965}{.1908} = 4596 \text{ yards nearly,}$$

$$FS_2 = 880 \cdot \frac{\sin 83^\circ. 45'}{\sin 11^\circ} = 880 \times \frac{.9940}{.1908} = 4584.48 \text{ yards.}$$

(4) Let  $AB$  be the flagstaff;  $BP$  the tower;  $Q$  the place of observation.

$$\begin{aligned}\text{Then } \tan BQA &= \tan(AQP - BQP) \\ &= \frac{\tan AQP - \tan BQP}{1 + \tan AQP \cdot \tan BQP} \\ &= \frac{2.05 - 2}{1 + 2.05 \times 2} = \frac{.05}{5.1} = \frac{1}{102}; \\ \therefore L \tan BQA &= 10 + \log 1 - \log 102 \\ &= 10 - 2.0086002 \\ &= 7.9913998; \\ \therefore BQA &= 33'. 42''.\end{aligned}$$




FIG. 32.

(5) Let  $A$  be the top of the steeple;  $D$  the top of the tower.

$$\angle APB = 60^\circ \text{ and } \angle DPB = 45^\circ.$$

$$\text{Then } BA = PB \cdot \tan 60^\circ,$$

$$\text{and } BD = PB \cdot \tan 45^\circ;$$

$$\therefore BA : BD = \tan 60^\circ : \tan 45^\circ$$

$$= \sqrt{3} : 1.$$




FIG. 33.

(6) Let  $PC$  be the river,  $CB$  the column,  $BA$  the statue  
 $CD = 6$  feet; and let  $x$  = breadth of  
 river in feet.

$$\text{Then } \tan APB = \tan DPC = \frac{6}{x},$$

$$\tan APC = \frac{AC}{PC} = \frac{230}{x},$$

$$\tan BPC = \frac{200}{x}.$$




FIG. 34.



Now  $\tan BPC = \tan(APC - APB)$ ;

$$\therefore \frac{200}{x} = \frac{\frac{230}{x} - \frac{6}{x}}{1 + \frac{230}{x} \cdot \frac{6}{x}};$$

$$\therefore \frac{200}{x} = \frac{224x}{x^2 + 1380}, \text{ or } 24x^2 = 276000, \text{ or } x^2 = 11500;$$

$$\therefore x = 107.2 \dots \text{feet.}$$

(7) Let  $A$  be the top of the pole;  $B$  the top of the mound.




FIG. 36.

$$\angle APQ = 60^\circ; \angle BPQ = 30^\circ.$$

$$\text{Then } AQ = PQ \cdot \tan 60^\circ,$$

$$BQ = PQ \cdot \tan 30^\circ;$$

$$\therefore AQ : BQ = \tan 60^\circ : \tan 30^\circ$$

$$= \sqrt{3} : \frac{1}{\sqrt{3}}$$

$$= 3 : 1;$$

$$\therefore AB = 2BQ.$$

(8) Let  $A$  be the top of the flagstaff;  $B$  the top of the tower.




FIG. 36.

Then  $\angle BQP = 90^\circ - \angle AQP$ .

$$\text{Now } AB = AP - BP$$

$$= a(\tan AQP - \tan BQP)$$

$$= a \cdot (\cot a - \tan a) = a \cdot \frac{\cos^2 a - \sin^2 a}{\cos a \cdot \sin a}$$

$$= 2a \cdot \frac{\cos 2a}{\sin 2a} = 2a \cdot \cot 2a.$$

(9) Let  $T$  be the place of the second observation.




FIG. 37.

Then  $a = QP$

$$= PT - TQ$$

$$= BP \cdot \cot \frac{a}{2} - c$$

$$= a \tan a \cdot \cot \frac{a}{2} - c;$$

$$\therefore c = a \left( \frac{2 \tan \frac{a}{2}}{1 - \tan^2 \frac{a}{2}} \cot \frac{a}{2} - 1 \right) = a \left( \frac{2}{1 - \tan^2 \frac{a}{2}} - 1 \right) = a \cdot \frac{1 + \tan^2 \frac{a}{2}}{1 - \tan^2 \frac{a}{2}}$$

$$= a \cdot \frac{\cos^2 \frac{a}{2} + \sin^2 \frac{a}{2}}{\cos^2 \frac{a}{2} - \sin^2 \frac{a}{2}} = \frac{a}{\cos a};$$

$\therefore a = c \cdot \cos a$ , and putting this for  $a$  in the result of Example (8),

$$\text{length of flagstaff} = 2c \cdot \cos a \cdot \cot 2a = 2c \cdot \cos a \cdot \frac{\cos 2a}{\sin 2a}$$

$$= 2c \cdot \frac{\cos 2a}{2 \sin a} = c \cdot \operatorname{cosec} a \cdot \cos 2a.$$

(10) Let  $K$  be the kite ;  $S_1$  and  $S_2$  the places of observation.

Draw  $KA$  perpendicular to  $S_1S_2$ .

Then, since the angles at  $S_1$  and  $S_2$  are equal

$KA$  bisects  $S_1S_2$ .

Then  $KA = AS_1 \cdot \tan KS_1A$

$$= \frac{a}{2} \cdot \tan \beta$$

$$= \frac{a}{2} \cdot \sin \beta \cdot \sec \beta$$

$$= \frac{a}{2} \cdot \sin \alpha \cdot \sec \beta, \text{ because } \alpha = \beta$$




FIG. 38.

(11) Let  $AB$  be the smaller and  $PT$  the greater tower, and  $D$  the point midway between them.




FIG. 39.

Join  $TA, BP, DA, DP$ .

Then  $\angle PDT = \angle DAB$ .

Let  $PT = x$  and  $AB = y$ .

Then  $\frac{x}{60} = \frac{60}{y}$ , or  $x = \frac{3600}{y}$ .

Also,  $\tan PBT = \tan 2ATB$

$$= \frac{2 \tan ATB}{1 - \tan^2 ATB}$$

and  $\tan PBT = \frac{x}{120}$ , and  $\tan ATB = \frac{y}{120}$ ;

$$\therefore \frac{x}{120} = \frac{240y}{14400 - y^2};$$

$$\therefore \frac{3600}{120y} = \frac{240y}{14400 - y^2}.$$

Hence  $y = 40$  feet, and  $\therefore x = 90$  feet.

(12) Since  $\angle ADB = \angle ACB$ , a circle can be described about  $ADCB$ .




FIG. 40.

$$\therefore \angle ABD = \angle ACD = 19^\circ. 15',$$

$$\text{and } \angle DAC = 180^\circ - (40^\circ. 45' + 19^\circ. 15') = 120^\circ.$$

$$\therefore AB = \frac{AD \cdot \sin 30^\circ}{\sin 19^\circ. 15'},$$

$$\text{and } AD = \frac{DC \cdot \sin 19^\circ. 15'}{\sin 120^\circ};$$

$$\therefore \frac{AB}{DC} = \frac{\sin 30^\circ}{\sin 120^\circ} = \frac{\sin 30^\circ}{\sin 60^\circ} = \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}} = \frac{1}{\sqrt{3}}.$$

(13) Let  $x$  be the length of the zigzag road in miles.

$$\text{Then } 5 : 12 = \frac{5}{3} : x;$$

$$\therefore 5x = 20, \text{ or } x = 4 \text{ miles.}$$

(14)  $S_1$  and  $S_2$  are the two positions of the ship,  $A$  and  $B$  the two objects.




FIG. 41.

$$\text{Then } \angle BS_1S_2 = 15^\circ + 45^\circ = 60^\circ$$

$$\angle BS_2S_1 = 90^\circ (\text{since N.W. is at right angles to N.E.})$$

$$\angle S_2AS_1 = 180^\circ - (45^\circ + 60^\circ) = 75^\circ.$$

$$\begin{aligned}\text{Then } BS_1 &= S_1 S_2 \cdot \sec \angle BS_1 S_2 \\ &= 5 \cdot \sec 60^\circ = 10,\end{aligned}$$

$$\text{and } AS_1 = \frac{S_1 S_2 \cdot \sin \angle AS_2 S_1}{\sin S_2 AS_1} = \frac{5 \cdot \sin 45^\circ}{\sin 75^\circ} = \frac{5 \times \frac{1}{\sqrt{2}}}{\frac{\sqrt{3}+1}{2\sqrt{2}}} = \frac{10}{\sqrt{3}+1}.$$

$$\therefore AB = 10 - \frac{10}{\sqrt{3}+1} = \frac{10\sqrt{3}}{\sqrt{3}+1} = \frac{10\sqrt{3}(\sqrt{3}-1)}{3-1} = 5(3-\sqrt{3}).$$

(15) Let  $PQ$  be the tower.

Then  $\angle AQP$  and  $\angle PQB$  are right angles.




FIG. 42.

$$\angle PAQ = 30^\circ, \text{ and } \angle PBQ = 18^\circ.$$

$$\text{Then } AQ = PQ \cdot \cot 30^\circ = PQ \times \sqrt{3},$$

$$BQ = PQ \cdot \cot 18^\circ = PQ \cdot \frac{\sqrt{(10+2\sqrt{5})}}{\sqrt{5}-1}. \quad (\text{See Example xxxvi. 6.})$$

$$\text{Now } BQ^2 - AQ^2 = a^2;$$

$$\therefore PQ^2 \left\{ \frac{10+2\sqrt{5}}{6-2\sqrt{5}} - 3 \right\} = a^2;$$

$$\therefore PQ^2 \left\{ \frac{5+\sqrt{5}}{3-\sqrt{5}} - 3 \right\} = a^2;$$

$$\therefore PQ^2 \cdot \frac{4(\sqrt{5}-1)}{3-\sqrt{5}} = a^2;$$

$$\therefore PQ^2 \cdot \frac{4 \cdot (2+2\sqrt{5})}{4} = a^2;$$

$$\therefore PQ = \frac{a}{\sqrt{(2+2\sqrt{5})}}.$$

(16) Let  $AB$  be the staff,  $C$  the centre of the ring in the vertical line  $ABC$ ,  $D$  the extremity of the shadow; then if  $DE$  be drawn touching the ring in  $E$ ,  $DE$  will be the direction of the sun, and  $CE$  is at right angles to  $DE$ .

Let  $CE=r$ , then  $AB=AD=8r$ , and  $AC=9r$ .

$$\therefore CD^2 = AC^2 + AD^2 = 145r^2,$$

$$\text{and } ED^2 = CD^2 - CE^2 = 144r^2;$$

$$\therefore ED = 12r.$$

$$\text{Hence } \tan ADC = \frac{9}{8}, \text{ and } \tan CDE = \frac{1}{12};$$

$$\therefore \tan ADE = \frac{\frac{9}{8} + \frac{1}{12}}{1 - \frac{9}{96}} = \frac{4}{3}.$$

$$\therefore \text{the sun's altitude} = \tan^{-1} \frac{4}{3}.$$




FIG. 43.

(17) Draw  $DM$ ,  $EN$  perpendicular to  $CB$ , and let  $AB, BC, CA$  be represented by  $c, a, b$ .




FIG. 44.

$$\text{Then } CD^2 = CM^2 + MD^2$$

$$= \frac{a^2}{9} + \frac{4b^2}{9} \quad (\text{EUCLID, VI. 2, Ex. 1.})$$

$$CE^2 = CN^2 + NE^2$$

$$= \frac{4a^2}{9} + \frac{b^2}{9}$$

$$DE^2 = \frac{c^2}{9};$$

$$\therefore CD^2 + CE^2 + DE^2 = \frac{5a^2}{9} + \frac{5b^2}{9} + \frac{a^2 + b^2}{9} = \frac{2}{3}(a^2 + b^2) = \frac{2}{3}c^2.$$

(18) Let  $P$  be the place of observation ;




FIG. 45.

$$BN = x, PB = a.$$

Then  $BA = a$ , because  $\angle APB = 45^\circ$ ,

$$PN = a \cdot \cot 30^\circ = a\sqrt{3},$$

$$\angle PBN = 135^\circ.$$

$$\text{Then } \cos PBN = \frac{PB^2 + BN^2 - PN^2}{2PB \cdot BN},$$

$$\text{or } \cos 135^\circ = \frac{a^2 + x^2 - 3a^2}{2ax},$$

$$\therefore -\frac{1}{\sqrt{2}} = \frac{x^2 - 2a^2}{2ax},$$

$$\therefore \frac{1}{2} = \frac{x^4 - 4a^2x^2 + 4a^4}{4a^2x^2}.$$

$$\text{Hence } x^4 - 6a^2x^2 = -4a^4, \text{ and } x^2 - 3a^2 = \pm\sqrt{5}a^2,$$

$$\text{and } \therefore x = a\sqrt{3 \pm \sqrt{5}}.$$

(19) Let  $BA$  be the first tower;  $AC$  the moat;  $ED$  the other tower.

Draw  $EF$  parallel to  $DA$ . Let  $h$  = height of  $BA$ .

Then since  $\angle BEA = \angle BCA = 45^\circ$ , a circle can be described about  $ABEC$ , and since  $\angle BAC = 90^\circ$ ,  $BC$  is the diameter of the circle, and therefore  $\angle BEC = 90^\circ$ .



FIG. 46.

$$\text{Then } CB^2 = CA^2 + BA^2 = 2h^2,$$

$$\text{and } CB^2 = EC^2 + EB^2$$

$$= a^2 + c^2 + EF^2 + FB^2$$

$$= a^2 + c^2 + (c+h)^2 + (h-a)^2.$$

Hence

$$2h^2 = a^2 + c^2 + c^2 + 2ch + h^2 + h^2 - 2ah + a^2;$$

$$\therefore h = \frac{a^2 + c^2}{a - c}.$$



$$(20) \quad AC = AB \cdot \frac{\sin 15^\circ}{\sin 150^\circ} = 100 \cdot \frac{\sin 15^\circ}{\sin 30^\circ};$$




FIG. 47.

$$\begin{aligned} \therefore AC &= 100 \times \frac{\sqrt{3}-1}{2\sqrt{2}} \div \frac{1}{2} = \frac{100(\sqrt{3}-1)}{\sqrt{2}} \\ &= 50(\sqrt{6}-\sqrt{2}) = 51.76 \dots \text{feet.} \end{aligned}$$

- (21) Since  $BC$  points to N.W. the  $\angle ABC = 45^\circ$ ;  
 $\therefore \angle ACB = 45^\circ$ , and  $AC = AB = 10$  miles.




FIG. 48.

$$\text{Also, } CB = \sqrt{AC^2 + AB^2} = \sqrt{200} = 14.14 \dots \text{miles.}$$

- (22) Let  $CA$  be a line from the end of the shadow in direction of the sun,  $AB$  the wall,  $BC$  the shadow.

$$\text{Then } \tan \angle ACB = \frac{AB}{BC} = \frac{18}{16} = \frac{9}{8}.$$

$$\therefore \tan \angle ACB = 1.125;$$

or,  $\angle ACB = \tan^{-1} 1.125$ , which by the tables  
 we find nearly equal to  $48^\circ. 22'.$




FIG. 49

- (23) Let  $AB$  be the spire ;  $BP$  the tower ;  $Q$  the place of observation.  
Then  $\angle BQP = 30^\circ$ , and  $\angle AQP = 32^\circ$ .




FIG. 50.

Now  $AP = PQ \cdot \tan 32^\circ = 200 \times \cdot 6248694 = 124\cdot97398$   
 $BP = PQ \cdot \tan 30^\circ = 200 \times \cdot 5773503 = 115\cdot47006$ .  
 $\therefore$  height of tower  $= 115\cdot47$  yards nearly,  
 height of spire  $= 9\cdot503$  yards nearly.

$$(24) \quad \cos BAC = \frac{9 + 4 - \frac{324}{100}}{12} = \frac{61}{75} = \cdot 8133333.$$

Hence, by the tables,  $\angle BAC = 35^\circ. 34'. 32''$ ,  
 and  $\therefore \angle BAD = 144^\circ. 25'. 28''$ .




FIG. 51.

$$\begin{aligned}\text{Next, } BD &= \frac{AB \cdot \sin 144^\circ.25'.28''}{\sin 17^\circ.47'.20''} \\ &= \frac{3 \times 5817759}{3055106} = 5.71307 \dots \text{ miles.}\end{aligned}$$

(25)  $\angle BAC = 17^\circ.44'$ ,

$$AB = BC \cdot \frac{\sin 139^\circ.58'}{\sin 17^\circ.44'}.$$

Hence, by the tables,

$$AB = \frac{840.5 \times .6432332}{.3045872} \text{ yards} = 1775 \text{ yards nearly};$$

$\therefore AB$  differs from a mile by about 15 yards.




FIG. 52.

(26)  $\angle BCA = 180^\circ - (50^\circ.20' + 110^\circ.12') = 19^\circ.28'$ ;

$$\therefore BC = AB \cdot \frac{\sin 50^\circ.20'}{\sin 19^\circ.28'}.$$

$$\begin{aligned}\therefore \log BC &= \log 2700 + L \sin 50^\circ.20' - L \sin 19^\circ.28' \\ &= 3.4313638 + 9.8863616 - 9.5227811 \\ &= 3.7949443.\end{aligned}$$

Hence  $BC = 6236.549$  feet.





FIG. 53.

Next, if  $CD$  be the height of the mountain,

$$\begin{aligned}CD &= BC \sin CBD, \\ &= 6236.549 \times \sin 10^\circ.7' \\ &= 6236.549 \times .1756531 \\ &= 1095.47 \dots \text{ feet.}\end{aligned}$$



Let  $EC = x$  feet.

Then  $\tan AEB = \tan(AEC - BEC)$  ;

$$\therefore \tan 10^\circ = \frac{\tan AEC - \tan BEC}{1 + \tan AEC \cdot \tan BEC}.$$

$$\therefore .176327 = \frac{\frac{60}{x} - \frac{40}{x}}{1 + \frac{2400}{x^2}}.$$

$$\therefore .176327 = \frac{20x}{x^2 + 2400} ;$$

and solving this quadratic we get

$$x = 85.28, \text{ or } 28.14.$$

(28)

Let  $CP$  be the height of the hill.




FIG. 55.

Then  $CA = AB \cdot \frac{\sin ABC}{\sin ACB}$

$$= 1760 \times \frac{\sin 2^\circ 45'}{\sin 9^\circ 28'}$$

$$= \frac{1760 \times .0479781}{.1644738} .$$

$$= 513.4 \text{ nearly ;}$$

$$\therefore CP = 513.4 \times \sin CAP$$

$$= 513.4 \times .2116091 = 108.64 \dots \text{ yards.}$$

(29) Let  $AB$  be the tower,  $S$  the ship.




FIG. 56.

$$\begin{aligned}\text{Then } BS &= AB \cdot \cot ASB \\ &= 150 \times 1.3613350 \\ &= 204.2 \dots \text{feet.}\end{aligned}$$

$$\begin{aligned}(30) \quad \tan \frac{A-B}{2} &= \frac{a-b}{a+b} \cdot \cot \frac{C}{2} \\ &= \frac{3225.77}{9541.29} \cot 18^\circ.43'.\end{aligned}$$

$$\begin{aligned}L \tan \frac{A-B}{2} &= 3.5086333 - 3.9795979 + 10.4700495 \\ &= 9.9990849.\end{aligned}$$

$$\text{Hence } \frac{A-B}{2} = 44^\circ.56'.20'',$$

$$\text{and } \frac{A+B}{2} = 71^\circ.17';$$

$$\therefore A = 116^\circ.13'.20'', \text{ and } B = 26^\circ.20'.40''.$$

$$\text{Also } c = \frac{b \cdot \sin C}{\sin B} = 3157.76 \times \frac{.6078379}{.4437665} = 4325.26.$$

$$\begin{aligned}(31) \quad \angle BS_1S_2 &= 55^\circ, \text{ and } \angle BS_2S_1 = 62^\circ.30'; \\ \therefore \angle S_1BS_2 &= 180^\circ - (55^\circ + 62^\circ.30') = 62^\circ.30'; \\ \therefore S_1S_2 &= BS_1 = 1 \text{ mile.}\end{aligned}$$

$$\begin{aligned}\text{Then } S_2B &= \frac{S_1B \cdot \sin BS_1S_2}{\sin BS_2S_1} \\ &= \frac{1 \times \sin 55^\circ}{\sin 62^\circ.30'} = \frac{.8191520}{.8870108} \\ &= .923497 \text{ miles.}\end{aligned}$$




FIG. 57.

(32) From  $E$ , the lower window, draw  $EB$  perpendicular to the tower  $AB$ ; from  $D$ , the upper window, draw  $DC$  perpendicular to the tower.




FIG. 58.

Then  $\angle AEB = 45^\circ$ ,

and  $\angle ADC = 40^\circ$ ,

and  $DC = EB = AB$ .

$$\therefore DC = 20 + AC$$

$$= 20 + DC \cdot \tan 40^\circ.$$

$$\begin{aligned} \therefore DC &= \frac{20}{1 - \tan 40^\circ} = \frac{20}{1 - .8390996} \\ &= \frac{20}{.1609004} = 124.3 \dots \text{feet.} \end{aligned}$$




FIG. 59.

Let  $CD$  be the perpendicular breadth of the river.

$$\text{Now } \angle ACB = 180^\circ - (50^\circ + 65^\circ) = 65^\circ.$$

$$\therefore AC = AB = 400 \text{ yards.}$$

$$\text{Hence } CD = AC \cdot \sin 50^\circ$$

$$= 400 \times .7660444 = 306.4178 \text{ yards.}$$

(34) Let  $AB, AC$  be the lines of the railways,  $D$  the point at which the train travelling 30 miles an hour is in  $2\frac{1}{2}$  hours.

The other train may then be at  $M$  or  $N$ , points on  $AB$  equidistant from  $D$ , and such that  $MD = DN = 50$  miles.




FIG. 60

Also,  $AD = 75$  miles.

$$\text{Then } \sin \angle AND = \frac{75 \cdot \sin 35^\circ \cdot 20'}{50} = \frac{3}{2} \times .5783323 = .8674984.$$

Hence  $\angle AND = 60^\circ \cdot 10'$  nearly,

$\therefore \angle ADN = 84^\circ \cdot 30'$ ,

$$\text{and } AN = \frac{50 \times \sin 84^\circ \cdot 30'}{\sin 35^\circ \cdot 20'} = \frac{50 \times .9953962}{.5783323} = \frac{49.7698100}{.5783323} \text{ miles.}$$

$$\therefore \text{rate of train} = \frac{49.7698100}{.5783323} \div 2\frac{1}{2} = 34.42284 \dots \text{ miles per hour.}$$

Next,  $\angle DMN = \angle AND = 60^\circ \cdot 10'$  nearly;

$\therefore \angle AMD = 119^\circ \cdot 50'$  nearly;

$\therefore \angle ADM = 24^\circ \cdot 50'$  nearly,

$$\text{and } AM = \frac{AD \cdot \sin \angle ADM}{\sin \angle AMD} = \frac{75 \cdot \sin 24^\circ \cdot 50'}{\sin 119^\circ \cdot 50'} = 75 \times \frac{.4199801}{.8674984} \text{ miles;}$$

$$\therefore \text{rate of train} = 75 \times \frac{.4199801}{.8674984} \div 2\frac{1}{2} = 14.524 \dots \text{ miles per hour.}$$

(35) Let  $AB$  be the base of 600 yards;  $C$  the tree;  $CD$  a perpendicular on  $AB$ .

$$\begin{aligned} \text{Then } \angle ACB &= 180^\circ - (52^\circ \cdot 14' + 68^\circ \cdot 32') \\ &= 59^\circ \cdot 14'. \end{aligned}$$




FIG. 61.

Now  $CD = AC \cdot \sin \angle CAD$

$$\begin{aligned} &= \frac{600 \cdot \sin \angle ABC}{\sin \angle ACB} \cdot \sin \angle CAD \\ &= \frac{600 \cdot \sin 68^\circ \cdot 32' \cdot \sin 52^\circ \cdot 14'}{\sin 59^\circ \cdot 14'} \end{aligned}$$

$$= \frac{600 \times .9306306 \times .7905115}{.8592576} = 513.7045 \text{ yards.}$$



(36) Let  $AB$  be the tower ;  $C$  the first place of observation ;  $D$  the second place of observation.




FIG. 62.

Then  $ACD$  and  $ABD$  are right angles.

Now  $AC = AB \cdot \operatorname{cosec} \angle ACB$

$$= 100 \times \operatorname{cosec} 50^\circ = 130.54073.$$

$$AD = \sqrt{(300)^2 + (130.54073)^2}$$

$$= \sqrt{107040.127569}$$

$$= 327.16 \dots$$

$$\sin \angle ADB = \frac{AB}{AD} = \frac{100}{327.16} = .3056608.$$

$$\text{Hence } \angle ADB = 17^\circ. 47'. 50''.$$

(37)




FIG. 63.

Let  $x$  = height of tower in yards ;

$$\tan 48'. 20'' = \tan (\angle ADC - \angle BDC)$$

$$\begin{aligned} & \frac{x+4}{100} - \frac{x}{100} \\ &= \frac{x \cdot (x+4)}{1 + \frac{x \cdot (x+4)}{10000}}; \end{aligned}$$

$$\therefore .0140605 = \frac{400}{10000 + x^2 + 4x}.$$

Solving this quadratic we get  $x = 134$  yards nearly.

(38) Let  $A$  be the object ;  $AB$  a vertical line meeting the horizontal plane through  $C$  in  $B$  ;  $D$  the point 300 yards up the hill.




FIG. 64.

Then  $\angle BCA = 29^\circ. 12'. 40'' = \angle CAD$ .

$$\angle CDA = 16^\circ.$$

$$\text{Then } CA = \frac{CD \cdot \sin 16^\circ}{\sin 29^\circ. 12'. 40''} = \frac{300 \times .2756374}{.4880290} = 169.4392 \text{ yards.}$$

(39) At the end of three hours each engine has passed over 90 miles.  
Let  $AB, AC$  be the distances traversed.

Draw  $AD$  perpendicular to  $BC$ .

Then  $\angle BAD = 25^\circ 10'$ ,

and  $BD = AB \cdot \sin BAD$

$$= 90 \times .4252528.$$

$$\therefore BC = 2 \times 90 \times .4252528 = 76.5455 \dots \text{miles.}$$




FIG. 65.

(40) Diagram as in Example (37); and  $x$  = height of tower in feet.

$$\tan ADB = \frac{6}{150} = \frac{1}{25};$$

$$\therefore \frac{1}{25} = \frac{\frac{x+30}{150} - \frac{x}{150}}{1 + \frac{x(x+30)}{22500}} = \frac{4500}{x^2 + 30x + 22500}.$$

Solving this quadratic  $x = 285$  feet nearly.

#### EXAMPLES—LVI. (p. 199).

$$\begin{aligned} 1. \text{ Area} &= \frac{1}{2} \left\{ 10 \times 12 \times \sin 60^\circ \right\} \text{ square inches} \\ &= \left( 60 \times \frac{\sqrt{3}}{2} \right) \text{ square inches} = 30\sqrt{3} \text{ square inches.} \end{aligned}$$

$$\begin{aligned} 2. \text{ Area} &= \frac{1}{2} \left\{ 40 \times 60 \times \sin 30^\circ \right\} \text{ square feet} \\ &= \left( 1200 \times \frac{1}{2} \right) \text{ square feet} = 600 \text{ square feet} \end{aligned}$$

$$3. \text{ Area} = \frac{1}{2} \left\{ 4 \times 3 \frac{3}{4} \right\} \text{ square feet} = 7 \frac{1}{2} \text{ square feet.}$$

$$\begin{aligned} 4. \text{ Area} &= \sqrt{s \cdot (s-a) \cdot (s-b) \cdot (s-c)} = \sqrt{8 \times 3 \times 2 \times 3} = 4 \times 3 \\ &= 12 \text{ square inches.} \end{aligned}$$

$$\begin{aligned} 5. \text{ Area} &= \sqrt{1017 \times 392 \times 512 \times 113} = \sqrt{9 \times 113 \times 8 \times 49 \times 8 \times 64 \times 113} \\ &= (3 \times 113 \times 8 \times 7 \times 8) = 151872. \end{aligned}$$

$$6. \text{ Area} = \sqrt{544 \times 135 \times 375 \times 34} = \sqrt{17 \times 32 \times 15 \times 9 \times 125 \times 3 \times 17 \times 2} \\ = 17 \times 8 \times 9 \times 25 = 30600.$$

$$7. \text{ Area} = \sqrt{585 \times 8 \times 512 \times 65} = \sqrt{5 \times 13 \times 9 \times 8 \times 64 \times 8 \times 13 \times 5} \\ = 5 \times 13 \times 3 \times 8 \times 8 = 12480.$$

$$8. \quad s \cdot (s-c) = \frac{a+b+c}{2} \cdot \frac{a+b-c}{2} \\ = \frac{(a+b)^2 - c^2}{4} \\ = \frac{(a+b)^2 - (a^2 + b^2)}{4} \\ = \frac{2ab}{4} \\ = \frac{ab}{2} = \text{area of the triangle.}$$

$$9. \text{ Area} = \sqrt{\frac{146 \cdot 27}{2} \times \frac{41 \cdot 21}{2} \times \frac{48 \cdot 75}{2} \times \frac{56 \cdot 31}{2}} \\ \therefore \log \text{ area} = \frac{1}{2} \left\{ \log 146 \cdot 27 + \log 41 \cdot 21 + \log 48 \cdot 75 + \log 56 \cdot 31 - 4 \log 2 \right\} \\ = \frac{1}{2} \left\{ 2 \cdot 1651553 + 1 \cdot 6150026 + 1 \cdot 6879746 \right. \\ \left. + 1 \cdot 7505855 - 1 \cdot 2041200 \right\} \\ = 3 \cdot 0072990 ; \\ \therefore \text{ area} = 1016 \cdot 9487.$$

10. Let  $a, b, c$  be in descending arithmetical progression ;  
then  $a + c = 2b$ .

Thus the perimeter is  $3b$ , and the side of an equilateral triangle of equal perimeter is  $b$ .

$$\text{Then } \sqrt{s \cdot (s-a)(s-b)(s-c)} = \frac{3}{5} \cdot \frac{1}{2} \cdot b^2 \cdot \sin 60^\circ, \\ \text{or } \frac{1}{4} \sqrt{(a+b+c)(b+c-a)(a+c-b)(a+b-c)} = \frac{3\sqrt{3}}{20} b^2,$$

$$\text{or } \sqrt{3b^2(b+c-a)(a+b-c)} = \frac{3\sqrt{3}}{5}b^2$$

$$\sqrt{(b+c-a)(a+b-c)} = \frac{3}{5}b$$

$$\sqrt{\frac{3c-a}{2} \cdot \frac{3a-c}{2}} = \frac{3}{10}(a+c)$$

$$\frac{10ac - 3a^2 - 3c^2}{4} = \frac{9}{100}(a^2 + c^2).$$

Solving this quadratic we get  $\frac{a}{c} = \frac{7}{3}$  or  $\frac{3}{7}$ .

Hence the sides are proportional to 7, 5, 3.

$$\text{Then } \cos A = \frac{b^2 + c^2 - a^2}{2bc} = -\frac{1}{2};$$

and  $\therefore A = 120^\circ$ .

11. Let  $AEB$  be the triangular part turned down.

Then area of  $AEB = \frac{1}{2}AB \cdot AE$ .




FIG. 66.

But  $\frac{AE}{AD} = \frac{AB}{BC}$ , by similar triangles  $AED$ ,  $BAC$ ;

$$\begin{aligned} \therefore \text{area of } AEB &= \frac{1}{2}AB \cdot \frac{AB \cdot AD}{BC} \\ &= \frac{1}{2} \cdot \frac{AB^2}{BC} \cdot (CD - AC) \\ &= \frac{1}{2} \cdot \frac{AB^2}{BC} \cdot \left\{ AB - \sqrt{AB^2 - BC^2} \right\} \end{aligned}$$

$$12. \text{Area} = \frac{bc \cdot \sin A}{2} = \frac{b \sin A \cdot c \sin A}{2 \sin A} = \frac{a \sin B \cdot a \sin C}{2 \sin A} = \frac{a^2 \sin B \cdot \sin C}{2 \sin(B+C)}.$$

$$\begin{aligned} 13. \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} & \left( \frac{a^2}{\sin A} + \frac{b^2}{\sin B} + \frac{c^2}{\sin C} \right) \\ &= \sqrt{\frac{(s-b)(s-c)}{bc}} \cdot \sqrt{\frac{(s-a)(s-c)}{ac}} \cdot \sqrt{\frac{(s-a)(s-b)}{ab}} \cdot \left( \frac{a^2 bc}{2S} + \frac{b^2 ac}{2S} + \frac{c^2 ba}{2S} \right) \\ &= \frac{(s-a)(s-b)(s-c)}{abc} \cdot \frac{abc(a+b+c)}{2S} \\ &= \frac{s \cdot (s-a)(s-b)(s-c)}{S} = S. \end{aligned}$$

$$14. \quad R = \frac{abc}{4S} \text{ and } r = \frac{S}{s};$$

$$\therefore \frac{abc}{4S} = \frac{2S}{s};$$

$$\begin{aligned} \therefore abc &= \frac{8S^2}{s} \\ &= 8(s-a)(s-b)(s-c) \\ &= (b+c-a)(a+c-b)(a+b-c). \end{aligned}$$

Squaring both sides,—

$$\begin{aligned} a^2 b^2 c^2 &= \{a + (b-c)\} \{a - (b-c)\} \times \{b + (a-c)\} \{b - (a-c)\} \\ &\quad \times \{c + (a-b)\} \{c - (a-b)\} \\ &= \{a^2 - (b-c)^2\} \{b^2 - (a-c)^2\} \{c^2 - (a-b)^2\}. \end{aligned}$$

Now this equality can only exist when  $a=b=c$ , for in any other case each factor on the right-hand side is less than the corresponding factor on the left-hand side.

$$\begin{aligned} 15. \quad \frac{b-c}{a} &= \frac{\sin B - \sin C}{\sin A} = \frac{2 \cos \frac{B+C}{2} \cdot \sin \frac{B-C}{2}}{2 \sin \frac{A}{2} \cdot \cos \frac{A}{2}} = \frac{\sin \frac{B-C}{2}}{\cos \frac{A}{2}}; \\ \therefore (b-c) \cos \frac{A}{2} &= a \cdot \sin \frac{B-C}{2}. \end{aligned}$$

16.  $OA$  bisects  $\angle A$ , and  $FE$  at right angles ;

$$\therefore \text{area } FOE = FH \cdot OH$$

$$= r \cos \frac{A}{2} \cdot r \sin \frac{A}{2}$$

$$= r^2 \cdot \frac{1}{2} \sin A$$

$$= \frac{S^2}{s^2} \cdot \frac{S}{bc}$$

$$= \frac{S^3}{s^2 \cdot bc}.$$




FIG. 67.

$\therefore$ , by symmetry,

$$\text{area } FDE = \frac{S^3}{s^2} \left( \frac{1}{bc} + \frac{1}{ca} + \frac{1}{ab} \right)$$

$$= \frac{S^3 \cdot 2s}{s^2 \cdot abc}$$

$$= \frac{2}{abc} \cdot \frac{\{s \cdot (s-a)(s-b)(s-c)\}^{\frac{1}{2}}}{s}$$

$$= \frac{2}{abc} \cdot s^{\frac{1}{2}} \left\{ (s-a)(s-b)(s-c) \right\}^{\frac{1}{2}}.$$

K

17. Area of quadrilateral

$$\begin{aligned}
 &= \frac{1}{2} \{ EA \cdot AD + DA \cdot AC + BA \cdot AC + BA \cdot AE \} \sin A \\
 &= \frac{1}{2} \{ (EA + AC) \cdot AD + (EA + AC) BA \} \sin A \\
 &= \frac{1}{2} \cdot EC \cdot BD \cdot \sin A \\
 &= \frac{1}{2} ab \cdot \sin A.
 \end{aligned}$$




FIG. 68.

$$\begin{aligned}
 18. \quad \frac{a^2 - b^2}{2} \cdot \frac{\sin A \cdot \sin B}{\sin(A - B)} &= \frac{a^2 \sin A \cdot \sin B - b^2 \sin A \cdot \sin B}{2 \sin(A - B)} \\
 &= \frac{ab \sin^2 A - ab \sin^2 B}{2 \sin(A - B)} = \frac{ab \cdot \sin(A + B) \cdot \sin(A - B)}{2 \sin(A - B)} \\
 &= \frac{ab \sin(A + B)}{2} \\
 &= \frac{ab \cdot \sin C}{2} = \text{area of triangle.}
 \end{aligned}$$

19.

$$R = \frac{a}{2 \sin A} = \frac{a}{\sqrt{2}}$$

$$r^a = \frac{S}{s - a} = \frac{\frac{1}{2} ab}{\frac{2a + c}{2} - a} = \frac{ab}{c} = \frac{a^2}{a\sqrt{2}} = \frac{a}{\sqrt{2}};$$

$$\therefore R = r^a.$$



$$\begin{aligned}
 20. \cot(B-A) + \cot 2\left(A + \frac{C}{2}\right) &= \cot(B-A) + \cot(2A+C) \\
 &= \frac{1 + \cot B \cdot \cot A}{\cot B - \cot A} + \frac{1 - \cot 2A \cdot \cot C}{\cot 2A + \cot C} \\
 &= \frac{1+1}{\tan A - \cot A} + \frac{1-0}{\cot 2A + 0} \\
 &= \frac{2}{\tan A - \cot A} + \frac{2 \tan A}{1 - \tan^2 A} \\
 &= \frac{2 \tan A}{\tan^2 A - 1} + \frac{2 \tan A}{1 - \tan^2 A} \\
 &= 0.
 \end{aligned}$$

$$\begin{aligned}
 21. \frac{2abc}{a+b+c} \cdot \cos \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2} \\
 &= \frac{2abc}{a+b+c} \cdot \sqrt{\frac{s \cdot (s-a)}{bc}} \cdot \sqrt{\frac{s \cdot (s-b)}{ac}} \cdot \sqrt{\frac{s \cdot (s-c)}{ab}} \\
 &= \frac{2abc}{a+b+c} \cdot \frac{s}{abc} \cdot \sqrt{s \cdot (s-a)(s-b)(s-c)} \\
 &= \sqrt{s \cdot (s-a)(s-b)(s-c)} \\
 &= \text{area of triangle.}
 \end{aligned}$$

$$\begin{aligned}
 22. \frac{\sin 2A (2a+c)^2}{32 \cdot \cos^4 \frac{A}{2}} &= \frac{\sin 2A \cdot (2s)^2}{32 \cdot \frac{s^2 \cdot (s-a)^2}{b^2 c^2}} \\
 &= \frac{\sin 2A \cdot b^2 c^2}{8 \cdot (s-a)^2} \\
 &= \frac{\sin 2A \cdot b^2 c^2}{8 \left(\frac{c}{2}\right)^2} \\
 &= \frac{b^2 \cdot \sin 2A}{2} = b^2 \cdot \sin A \cdot \cos A = b^2 \cdot \sin A \cdot \frac{c}{2b} \\
 &= \frac{1}{2} bc \cdot \sin A \\
 &= \text{area of triangle.}
 \end{aligned}$$

$$\therefore \text{area} \times 32 \cos^4 \frac{A}{2} = \sin 2A \cdot (2a+c)^2.$$




FIG. 69.

23.  $AD = b \cdot \sin C, \therefore AD \cdot b = b^2 \cdot \sin C.$   
 $AD = c \cdot \sin B, \therefore AD \cdot c = c^2 \cdot \sin B.$




FIG. 70.

$$\therefore AD(b + c) = b^2 \cdot \sin C + c^2 \cdot \sin B;$$

$$\therefore AD = \frac{b^2 \sin C + c^2 \sin B}{b + c}.$$

24. (1)

$$BD = r \cdot \cot \frac{B}{2}$$

$$CD = r \cdot \cot \frac{C}{2};$$




FIG. 71.

$$\therefore r \cdot \left( \cot \frac{B}{2} + \cot \frac{C}{2} \right) = BD + CD = a.$$

$$\therefore r = \frac{a}{\cot \frac{B}{2} + \cot \frac{C}{2}}.$$

(2) From the preceding Example—

$$r = \frac{a \sin \frac{B}{2} \cdot \sin \frac{C}{2}}{\sin \left( \frac{B+C}{2} \right)}$$

$$= \frac{2R \cdot \sin A \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}}{\cos \frac{A}{2}}, \text{ by Art. 221.}$$

(3) Let  $O$  be the centre of the escribed circle touching  $BC$  and the other sides produced, as in diagram to Art. 223.

$$\text{Then } BD = OD \cdot \cot OBD = r_1 \cdot \tan \frac{B}{2},$$

$$\text{and } CD = OD \cdot \cot OCD = r_1 \cdot \tan \frac{C}{2}.$$

$$\therefore BD + CD = r_1 \left( \tan \frac{B}{2} + \tan \frac{C}{2} \right);$$

$$\therefore r_1 = \frac{a}{\tan \frac{B}{2} + \tan \frac{C}{2}}.$$




FIG. 72.

(4) By the preceding Example—

$$r_1 = \frac{a \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2}}{\sin \frac{B+C}{2}}$$

$$= \frac{2R \cdot \sin A \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2}}{\cos \frac{A}{2}}$$

$$= 4R \cdot \sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2}.$$

$$(5) \quad r_1 = 4R \cdot \sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2},$$

$$r_2 = 4R \cdot \cos \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \cos \frac{C}{2},$$

$$r_3 = 4R \cdot \cos \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \sin \frac{C}{2};$$

$$\therefore r_1 + r_2 + r_3 = 4R \cdot \cos \frac{A}{2} \cdot \left( \sin \frac{B}{2} \cdot \cos \frac{C}{2} + \cos \frac{B}{2} \cdot \sin \frac{C}{2} \right) \\ + 4R \cdot \sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2}$$

$$= 4R \cdot \cos \frac{A}{2} \cdot \cos \frac{A}{2} + 4R \cdot \sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2}$$

$$= 2R \cdot (\cos A + 1) + R \cdot (\cos B + \cos C - \cos A + 1)$$

EX. XLVIII. 12.

$$= 3R + R(\cos A + \cos B + \cos C).$$

$$(6) \quad R + r = R + \frac{2R \sin A \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}}{\cos \frac{A}{2}} \text{ by (2)}$$

$$= R + 4R \cdot \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}$$

$$= R \cdot \left( 1 + 4 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} \right)$$

$$= R(\cos A + \cos B + \cos C).$$

EX. XLVIII. 8.

25. Let  $r$  be the radius of the circle.Then area of inscribed polygon of  $2n$  sides  $= nr^2 \cdot \sin \frac{\pi}{n}$ ,area of inscribed polygon of  $n$  sides  $= \frac{nr^2}{2} \cdot \sin \frac{2\pi}{n}$ ;area of circumscribed polygon of  $n$  sides  $= nr^2 \cdot \tan \frac{\pi}{n}$ .

$$\begin{aligned}
 & \text{And} \left( \frac{nr^3}{2} \cdot \sin \frac{2\pi}{n} \right) \times \left( nr^3 \cdot \tan \frac{\pi}{n} \right) \\
 & \quad \frac{n^3 \cdot r^4}{2} \cdot 2 \sin \frac{\pi}{n} \cdot \cos \frac{\pi}{n} \cdot \frac{\sin \frac{\pi}{n}}{\cos \frac{\pi}{n}} \\
 & = n^2 r^4 \cdot \sin^2 \frac{\pi}{n} \\
 & = \left( nr^3 \cdot \sin \frac{\pi}{n} \right)^2
 \end{aligned}$$

26. Let  $O$ ,  $M$  be the centres of the inscribed and escribed circles.




FIG. 78.

Then  $MO = MA - OA$

$$\begin{aligned}
 & = r_1 \operatorname{cosec} \frac{A}{2} - r \cdot \operatorname{cosec} \frac{A}{2} \\
 & = (r_1 - r) \operatorname{cosec} \frac{A}{2} \\
 & = \left\{ 4R \cdot \sin \frac{A}{2} \cdot \cos \frac{B}{2} \cdot \cos \frac{C}{2} - 4R \cdot \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2} \right\} \operatorname{cosec} \frac{A}{2} \\
 & \quad \text{(By Ex. 24.)} \\
 & = 4R \left\{ \cos \frac{B}{2} \cdot \cos \frac{C}{2} - \sin \frac{B}{2} \cdot \sin \frac{C}{2} \right\} \\
 & = 4R \cdot \sin \frac{A}{2},
 \end{aligned}$$

and similarly for the other escribed circles.

(27) Let  $DEF$  be the triangle so formed.

Then since  $\frac{BD}{CD} = \frac{c}{b}$ ,

$$\frac{BD}{BC} = \frac{c}{b+c}, \text{ or, } BD = \frac{ac}{b+c}.$$

So also,  $CD = \frac{ab}{b+c}$ , and similarly for the segments of the other sides.




FIG. 74.

$$\text{Then area } CDE = \frac{1}{2} \cdot \frac{ab}{b+c} \cdot \frac{ab}{a+c} \cdot \sin C = \frac{S \cdot ab}{(a+c)(b+c)}.$$

Similar expressions may be obtained for the areas of  $BFD$ ,  $AFE$ .

$$\therefore \text{area of } DEF = S \left\{ 1 - \frac{ab}{(a+c)(b+c)} - \frac{bc}{(b+a)(c+a)} - \frac{ca}{(c+b)(a+b)} \right\}$$

$$= \frac{2abc \cdot S}{(a+b)(b+c)(c+a)} = 2S \cdot \frac{a}{b+c} \cdot \frac{b}{c+a} \cdot \frac{c}{a+b}.$$

$$\text{Now, } \frac{a}{b+c} = \frac{\sin A}{\sin B + \sin C} = \frac{\sin \frac{A}{2}}{\cos \frac{B-C}{2}}$$

$$\frac{b}{c+a} = \frac{\sin \frac{B}{2}}{\cos \frac{C-A}{2}}, \text{ and } \frac{c}{a+b} = \frac{\sin \frac{C}{2}}{\cos \frac{A-B}{2}}$$

$$\therefore \frac{\text{area } DEF}{\text{area } ABC} = \frac{2 \sin \frac{A}{2} \cdot \sin \frac{B}{2} \cdot \sin \frac{C}{2}}{\cos \frac{B-C}{2} \cdot \cos \frac{C-A}{2} \cdot \cos \frac{A-B}{2}}.$$

$$\begin{aligned} 28. \quad r_1 r_2 + r_2 r_3 + r_3 r_1 &= \frac{S^2}{(s-a)(s-b)} + \frac{S^2}{(s-b)(s-c)} + \frac{S^2}{(s-c)(s-a)} \\ &= s \cdot (s-c) + s \cdot (s-a) + s \cdot (s-b) \\ &= s \cdot \{3s - (a+b+c)\} \\ &= s^2. \end{aligned}$$

$$29. \quad \frac{\sin BAD}{\sin ADB} = \frac{BD}{AB}.$$

$$\frac{\sin ABC}{\sin ACB} = \frac{AC}{AB}.$$

$$\therefore \text{since } \sin ADB = \sin ACB,$$

$$\frac{\sin BAD}{\sin ABC} = \frac{BD}{AC};$$

$$\therefore AC \sin A = BD \cdot \sin B.$$




FIG. 75.

30. Let  $O, P$  be the centres of the inscribed and one of the escribed circles.

Then  $OB$  and  $PB$  bisect the interior and exterior angles at  $B$ ; and  $\therefore OBP$  is a right angle.

Hence  $OBPC$  is a quadrilateral round which a circle may be described.

$$\text{Then } OP = OB \cdot \sec BOP$$

$$= OB \cdot \sec BCP$$

$$= OB \cdot \operatorname{cosec} \frac{C}{2}.$$

$$\text{And } OB = \frac{c \cdot \sin \frac{A}{2}}{\sin AOB} = \frac{c \cdot \sin \frac{A}{2}}{\cos \frac{C}{2}};$$




FIG. 76

$$\therefore OP = \frac{c \cdot \sin \frac{A}{2}}{\sin \frac{C}{2} \cdot \cos \frac{C}{2}} = \frac{2c \cdot \sin \frac{A}{2}}{\sin C} = \frac{2a \cdot \sin \frac{A}{2}}{\sin A} = \frac{a}{\cos \frac{A}{2}}.$$

$$\text{Similarly } OP = \frac{b}{\cos \frac{B}{2}} = \frac{c}{\cos \frac{C}{2}}.$$



$$\begin{aligned}
 31. \quad r \cdot \cos \frac{A}{2} \cdot \operatorname{cosec} \frac{B}{2} \cdot \operatorname{cosec} \frac{C}{2} &= \frac{r \cos \frac{A}{2}}{\sin \frac{B}{2} \cdot \sin \frac{C}{2}}, \\
 &= r \cdot \frac{\sqrt{\frac{s \cdot (s-a)}{bc}}}{\sqrt{\frac{(s-a) \cdot (s-c)}{ac}} \cdot \sqrt{\frac{(s-b)(s-a)}{ab}}} \\
 &= r \cdot \frac{a \sqrt{s}}{\sqrt{(s-a)(s-b)(s-c)}} \\
 &= \frac{S}{s} \cdot \frac{a \sqrt{s}}{\sqrt{(s-a)(s-b)(s-c)}} \\
 &= a.
 \end{aligned}$$

$$\begin{aligned}
 32. \quad \tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2} &= \sqrt{\frac{(s-c)(s-b)}{s \cdot (s-a)}} + \sqrt{\frac{(s-a)(s-c)}{s \cdot (s-b)}} + \sqrt{\frac{(s-b)(s-a)}{s \cdot (s-c)}} \\
 &= \frac{(s-c)(s-b)}{S} + \frac{(s-a)(s-c)}{S} + \frac{(s-b)(s-a)}{S} \\
 &= \frac{1}{4S} \cdot \left\{ (a+b-c) \cdot (a+c-b) + (b+c-a) \cdot (a+b-c) \right. \\
 &\quad \left. + (a+c-b) \cdot (b+c-a) \right\} \\
 &= \frac{1}{4S} \left\{ 2ab + 2ac + 2bc - a^2 - b^2 - c^2 \right\} \\
 &= \frac{1}{S} \cdot \left\{ ab + ac + bc - \frac{a^2 + b^2 + c^2 + 2ab + 2ac + 2bc}{4} \right\} \\
 &= \frac{1}{S} \left\{ ab + ac + bc - s^2 \right\} \\
 &= \frac{ab + ac + bc}{S} - \frac{s^2}{S} \\
 &= \frac{4R}{abc} \cdot (ab + ac + bc) - \frac{s}{r} \\
 &= 4R \cdot \left( \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) - \frac{s}{r}.
 \end{aligned}$$

33.  $PA \cdot BC = BA \cdot PC + AC \cdot BP$

(EUCLID, VI. D.)

$$\text{and } \frac{\sin A}{BC} = \frac{\sin C}{BA} = \frac{\sin B}{AC}.$$




FIG. 77.

$$\therefore PA \cdot \sin A = PC \cdot \sin C + PB \cdot \sin B.$$

34. Each of the angles at  $O = 120^\circ$ .

Let  $OA, OB, OC$  be represented by  $d_1, d_2, d_3$ .

$$c^2 = d_1^2 + d_2^2 - 2d_1d_2 \cdot \cos 120^\circ;$$

$$\therefore c = \sqrt{d_1^2 + d_2^2 + d_1d_2}.$$




FIG. 78.

Similarly for  $a$  and  $b$ .

$$\text{Also, area} = \left( \frac{d_1d_2}{2} + \frac{d_1d_3}{2} + \frac{d_2d_3}{2} \right) \sin 120^\circ$$

$$= \frac{\sqrt{3}}{4} \cdot (d_1d_2 + d_1d_3 + d_2d_3).$$

35. Let  $OA=a$ ,  $OB=b$ ,  $OC=c$ ;  $\angle OBA=\theta$ , and let  $x$  be the side of the square  $ABCD$ .




FIG. 79.

Then  $\angle OBC=90^\circ-\theta$ ,

$$\text{and } a^2=x^2+b^2-2bx\cos\theta;$$

$$c^2=x^2+b^2-2bx\sin\theta;$$

$$\therefore 2bx\cos\theta=x^2+b^2-a^2,$$

$$2bx\sin\theta=x^2+b^2-c^2.$$

Squaring and adding these equations,

$$4b^2x^2=x^4+2(b^2-a^2)x^2+(b^2-a^2)^2+x^4+2(b^2-c^2)x^2+(b^2-c^2)^2;$$

$$\therefore 2x^4-2(a^2+c^2)x^2+(a^2+c^2)^2=2(a^2b^2+a^2c^2+b^2c^2-b^4),$$

$$\text{and } x=\sqrt{\frac{1}{2}\left\{a^2+c^2\pm\sqrt{4(a^2b^2+a^2c^2+b^2c^2-b^4)-(a^2+c^2)^2}\right\}}.$$

(Gaskin's *Solutions of Trigonometrical Examples*.)

36. Let  $ABC$  be any triangle described about a circle.




FIG. 80.

Then area of  $ABC$ =area of  $AOB$ +area of  $BOC$ +area of  $AOC$ .

$$\therefore \text{area of } ABC=\frac{1}{2}\cdot rc+\frac{1}{2}ra+\frac{1}{2}rb.$$

$$=\frac{r}{2}(a+b+c);$$

$\therefore$  since  $r$  is constant,

$$\text{area of } ABC\propto(a+b+c).$$

$$\begin{aligned}
 37. \quad a &= AD = c \cdot \sin B = b \cdot \sin C, \\
 \beta &= BE = c \cdot \sin A, \\
 \gamma &= CF = b \cdot \sin A; \\
 \therefore \frac{a^2}{\beta\gamma} &= \frac{bc \cdot \sin B \cdot \sin C}{bc \cdot \sin A \cdot \sin A} = \frac{bc}{a^2}.
 \end{aligned}$$




FIG. 81.

$$\begin{aligned}
 \text{Similarly } \frac{\beta^2}{\alpha\gamma} &= \frac{ac}{b^2}; \text{ and } \frac{\gamma^2}{\alpha\beta} = \frac{ab}{c^2}; \\
 \therefore \frac{a^2}{\beta\gamma} + \frac{\beta^2}{\alpha\gamma} + \frac{\gamma^2}{\alpha\beta} &= \frac{bc}{a^2} + \frac{ac}{b^2} + \frac{ab}{c^2}.
 \end{aligned}$$

38. Let  $A$  be the observer on the sea-shore,  $O$  the earth's centre,  $BC$  the mountain whose height = 1284·8 yards = ·73 miles.

Then since  $C$  is just visible from  $A$ ,

$AC$  is a tangent at  $A$ .

Join  $OA$  and produce it to  $D$ , making  $AD = 3$  miles; then  $\angle DCA =$  angle of depression of  $C$  from  $D = 2^\circ. 15'$ .




FIG. 82.

Then  $AC = 3 \cdot \cot 2^\circ. 15'$

$$\begin{aligned}
 \log AC &= \log 3 + L \cot 2^\circ. 15' - 10 \\
 &= \cdot 4771213 + 11 \cdot 4057168 - 10 \\
 &= 1 \cdot 8828381; \\
 \therefore AC &= 76 \cdot 3551.
 \end{aligned}$$

Let  $OA$ , the earth's radius,  $=r$ ;

$$\therefore AC^2 = BC(2r + BC) = .73(2r + .73),$$

$$\text{and } \log(2r + .73) = 2 \log AC - \log .73 = 3.9023533;$$

$$\therefore 2r + .73 = 7986.4;$$

$$\therefore r = 3992.835 \text{ miles.}$$

(Gaskin's *Solutions of Trigonometrical Examples*.)

39. Let  $ABC$  be the triangle,  $CO=b$ ,  $BO=a$ ,




FIG. 83.

$\angle BAD = \theta$ , and  $\therefore \angle CAD = 60^\circ - \theta$ .

Let  $AB = x$ .

$$\text{Then } x \cdot \sin \theta = a,$$

$$x \sin(60^\circ - \theta) = b;$$

$$\therefore \frac{\sin(60^\circ - \theta)}{\sin \theta} = \frac{b}{a}.$$

$$\therefore \frac{\sqrt{3}}{2} \cdot \cot \theta - \frac{1}{2} = \frac{b}{a}.$$

$$\therefore a \cdot \cot \theta = \frac{3b + a}{\sqrt{3}}.$$

$$\therefore x = a \cdot \operatorname{cosec} \theta = \sqrt{a^2 + \frac{4b^2 + 4ab + a^2}{3}} = 2\sqrt{\frac{a^2 + ab + b^2}{3}} \quad (\text{Gaskin}).$$

40.




FIG. 84.

$$\frac{AD}{DB} = \frac{AC}{CB} = 2; \therefore AD = 2DB,$$

$$AB = AD + DB = 3DB,$$

$$\frac{AE}{EB} = \frac{AC}{CB} = 2; \therefore AE = 2BE;$$

$$\therefore BE = AB = 3DB;$$

$$\therefore DE = BE + DB = 4DB.$$

Then, by EUCLID, VI. i.

$$\triangle CBD : \triangle ACD : \triangle ABC : \triangle CDE \\ = DB : AD : AB : DE$$

$$= 1 : 2 : 3 : 4. \quad (\text{Gaskin}).$$

$$\begin{aligned}
 41. \quad R \cdot \sin A &= \frac{a}{2}, \text{ by Art. 221 ;} \\
 \therefore Rr \cdot (\sin A + \sin B + \sin C) \\
 &= r \cdot \left( \frac{a+b+c}{2} \right) \\
 &= r \cdot s \\
 &= \text{area of the triangle}
 \end{aligned}$$

$$42. \text{ The circles have the same radius because } R = \frac{b}{2\sin B}.$$

In the example given,  $\sin 50^\circ. 15' = .7688418$  ;

$$\therefore R = \frac{564}{1.5376836} = 366.785.$$

$$43. \text{ Call the angles } x, \frac{x+y}{2}, \frac{x+2y}{2}, \frac{x+3y}{3}.$$

$$\begin{aligned}
 \text{Then } x + \frac{x+y}{2} + \frac{x+2y}{2} + \frac{x+3y}{3} &= 2\pi \\
 \text{and } x + \frac{x+2y}{2} &= \pi
 \end{aligned} \left. \vphantom{\begin{aligned} \text{Then } x + \frac{x+y}{2} + \frac{x+2y}{2} + \frac{x+3y}{3} \\ \text{and } x + \frac{x+2y}{2} \end{aligned}} \right\} ;$$

$$\begin{aligned}
 \therefore 14x + 15y &= 12\pi \\
 3x + 2y &= 2\pi
 \end{aligned} \left. \vphantom{\begin{aligned} \therefore 14x + 15y \\ 3x + 2y \end{aligned}} \right\} .$$

$$\text{Hence } x = \frac{6\pi}{17} \text{ and } y = \frac{8\pi}{17} ;$$

$$\therefore \text{ the angles are } \frac{6\pi}{17}, \frac{7\pi}{17}, \frac{11\pi}{17}, \frac{10\pi}{17}$$

$$\begin{aligned}
 44. \quad \frac{(1 + \cot PCA)^2}{(1 + \cot PCB)^2} &= \frac{\left(1 - \frac{CM}{PM}\right)^2}{\left(1 + \frac{CM}{PM}\right)^2} \\
 &= \frac{(PM - CM)^2}{(PM + CM)^2} \\
 &= \frac{CP^2 - 2CN \cdot PN}{CP^2 + 2CN \cdot PN} \\
 &= \frac{CN \cdot CD - 2CN \cdot PN}{CN \cdot CD + 2CN \cdot PN} \\
 &= \frac{CO - PN}{CO + PN} = \frac{CB - CM}{AC + CM} = \frac{BM}{AM} = \frac{\cot PBA}{\cot PAB}.
 \end{aligned}$$




FIG. 85.

$$\begin{aligned}
 45. \quad r + r_a + r_b - r_c &= \frac{S}{s} + \frac{S}{s-a} + \frac{S}{s-b} - \frac{S}{s-c} \\
 &= \frac{S(2s-a)}{s \cdot (s-a)} + \frac{S \cdot (s-c-s+b)}{(s-b)(s-c)} \\
 &= \frac{S \cdot (b+c)}{s \cdot (s-a)} + \frac{S \cdot (b-c)}{(s-b)(s-c)} \\
 &= S \cdot \left\{ \frac{b \cdot \{(s-b)(s-c) + s \cdot (s-a)\} + c \{(s-b)(s-c) - s \cdot (s-a)\}}{S^2} \right\} \\
 &= \frac{b \{2s^2 - s \cdot (a+b+c) + bc\} + c \{s(b+c-a) + bc\}}{S} \\
 &= \frac{b^2c - \frac{c}{2}(b+c+a)(b+c-a) + bc^2}{S} \\
 &= \frac{c}{2S} \cdot \left\{ 2b^2 - (b+c)^2 + a^2 + 2bc \right\} \\
 &= \frac{c}{2S} (b^2 - c^2 + a^2) = \frac{c}{2S} 2ab \cos C = \frac{abc \cdot \cos C}{S} = 4R \cdot \cos C.
 \end{aligned}$$

46. Let  $C$  be the right angle; then, by Art. 223,




FIG. 86.

$$\begin{aligned}
 S &= \frac{c+b-a}{2} \cdot r, \text{ and} \\
 S &= \frac{c+a-b}{2} \cdot r'; \\
 \therefore S^2 &= \frac{c^2 - (a-b)^2}{4} \cdot rr' \\
 &= \frac{c^2 - a^2 + 2ab - b^2}{4} \cdot rr' = \frac{ab}{2} rr' = S \cdot rr'; \\
 \therefore rr' &= S.
 \end{aligned}$$

47.




FIG. 87.

Using the notation of Art. 228,

$$OB = 4000 \text{ miles,}$$

$$OE = OB \cdot \sec. 1^\circ 58' 10''$$

$$= 4000 \times 1.005910$$

$$= 4002.364.$$

$$\therefore AE = 2.36 \dots \text{miles.}$$



48. Using the notation of Art. 228,

$$\sec EOB = \frac{4001.25}{4000} = 1.0003125,$$

and, by the Tables,  $\sec 1^\circ.26' = 1.0003130$ .

Hence dip of horizon  $= 1^\circ.26'$  nearly.

49. Let  $A$  be the man's eye;  $B$  the lamp;  $C$  the centre of the earth.

Then  $AD + DB = 52800$  feet.

And, if the radius of the earth be  $R$  feet,

$$AD^2 = 6(2R + 6),$$

$$BD^2 = 32(2R + 32).$$




FIG. 88.

Hence, approximately,  $\sqrt{12R} + \sqrt{64R} = 52800$ ,

or,  $\sqrt{R} \cdot (4 + \sqrt{3}) = 26400$ , or,  $\sqrt{R} \times 13 = 26400(4 - \sqrt{3})$ ;

$$\therefore R = \frac{26400 \times 26400 \times (19 - 8\sqrt{3})}{13 \times 13 \times 1760 \times 3} \text{ miles} = 4017.79 \dots \text{ miles.}$$

50. In 72 minutes the ship travels 12 miles.

Then using the notation of Art. 228,

$$BE^2 = CE \cdot EA,$$

$$144 = \left( CA + \frac{90}{5280} \right) \cdot \frac{90}{5280}$$

$$= CA \cdot \frac{90}{5280} \text{ nearly;}$$

$$\therefore CA = \frac{144 \times 528}{9} = 16 \times 528 = 8448.$$

$$\therefore \text{radius} = 4224 \text{ miles.}$$

L

51. Using the notation of Art. 228,

$$\cos EOB = \frac{OB}{OE} = \frac{3956}{3959}.$$

$$\begin{aligned}\therefore L \cos EOB &= 10 + \log 3956 - \log 3959 \\ &= 10 + 3.5972563 - 3.5975855 \\ &= 9.9996708.\end{aligned}$$

Whence, by the tables,

$$EOB = 2^\circ. 13'. 50''.$$

52. Let  $r$  be the radius of a section of the earth, made by a plane through its centre perpendicular to the line joining its centre with the sun's centre. Then if  $\theta$  be the circular measure of the angle subtended by  $r$  at the sun's centre, and  $d$  be the distance between the two centres,

$$\frac{r}{d} = \tan \theta = \theta \text{ nearly, since } \theta \text{ is very small.}$$

$$\therefore \frac{r}{d} = \frac{8.868}{57.29577 \times 60 \times 60}.$$

$$\therefore d = \frac{57.29577 \times 60 \times 60 \times 4000}{8.868}$$

$$= \frac{206264772}{2.217} = 93037786.1 \dots \text{ miles}$$

53. Using the same notation as in Ex. 52,

$$\tan \theta = \frac{4000}{241118};$$

$$\begin{aligned}\therefore L \tan \theta &= 10 + \log 4000 - \log 241118 \\ &= 10 + 3.6020600 - 5.3822296 \\ &= 8.2198304.\end{aligned}$$

Hence, by the tables,

$$\theta = 57'. 1''.5 = \text{nearly.}$$

54. Let  $A, B$  be the two points ; then  $AB$  is a tangent at its middle point  $D$  to the earth's surface.

$$AD = DE \text{ nearly} = 4 \text{ miles,}$$

$$AE = 10 \text{ feet} = \frac{10}{5280} \text{ miles.}$$

Let  $C$  be the earth's centre, and  $CD = r$ .

$$\text{Then } AE(2r + AE) = AD^2.$$

$$\therefore, \text{approximately, } AE \cdot 2r = AD^2 ;$$

$$\therefore r = \frac{16 \times 5280}{10 \times 2} = 4224 \text{ miles.}$$




FIG. 89.

55. The limit of deviation is the angle subtended by the radius of the target at a point 600 feet distant, and if this angle be denoted by  $\theta$

$$\tan \theta = \frac{2}{600} ;$$

$$\therefore \theta = \tan^{-1} 0.00\dot{3}.$$

56. Regard the moon  $M$  as the base of a cone of which  $E$ , the eye of the observer, is the vertex. Then  $S$ , the shilling, will intercept all the rays of light from  $M$  to  $E$ , when it is so near to  $S$  that lines from  $E$  to the circumference of  $S$  do not, when produced, fall within the circumference of  $M$ .



FIG. 90.



3, WATERLOO PLACE, LONDON,  
*January, 1885.*

## EDUCATIONAL WORKS

PUBLISHED BY

MESSRS. RIVINGTON

---

### NEW BOOKS IN PREPARATION AND IN THE PRESS

**Lectures on Greek Prose.** By ARTHUR SIDGWICK, M.A.,  
*Fellow and Tutor of Corpus Christi College, Oxford, and late Assistant  
Master at Rugby School.* [In preparation.]

This book attempts to give help in Greek Prose Composition, not by general rules or principles, but by discussing the best way of turning actual pieces of English, and showing how the difficulties arise in practice, and how they must be met. Illustrations are also given of the errors and pitfalls to which the learner is liable. In his book on Greek Prose Composition the writer gave rules, and hints, and principles. In this book he aims at exhibiting these in their practical application.

**The Jugurtha of Sallust.** By E. P. BROOKE, M.A., *Assistant  
Master at Rugby School.* [In the press.]

**Versiculi.** A Latin Elegiac Verse Book. By the Rev. J. H. RAVEN,  
M.A., *Head Master of the Fauconberge School, Beccles.* Crown  
8vo. 2s. [Now ready.]

**Letters of Cicero.** Selected and Edited, with Introduction and  
Notes. By J. H. MUIRHEAD, B.A., *Assistant to Professor of  
Humanity in the University of Glasgow.* Crown 8vo. 6s.  
[Now ready.]

**Cicero Pro Cluentio.** Edited with Introduction, Notes, &c. By  
W. YORKE FAUSSETT, B.A., *Assistant Master in the Grammar  
School, Manchester.* [In preparation.]

**An Elementary Greek Syntax.** By F. E. THOMPSON, M.A.,  
*Assistant Master at Marlborough College; Author of "A Syntax of  
Attic Greek for the use of Students," &c.* Crown 8vo. 2s.  
[Now ready.]

**Easy Selections from Thucydides.** For the use of Middle  
Forms of Schools. Edited by E. H. MOORE, M.A., *Assistant Master  
in the High School, Plymouth.* Crown 8vo. [Nearly ready.]

---

Waterloo Place, London,

- New Books in Preparation and in the Press—continued.**
- Passages for Latin and Greek Unseen Translation.**  
For the use of Lower and Middle Forms of Schools. By J. ARNOLD TURNER, B.A., *Assistant Master at Hillbrow School, Rugby.* Small 8vo. 2s. 6d. [Now ready.]
- A Collection of Arithmetical Exercises,** progressively arranged. By A. E. DONKIN, M.A., and C. H. HODGES, M.A., *Mathematical Masters at Rugby School.* Small 8vo. 2s. 6d. [Now ready.]
- A History of the Romans.** For the use of Middle Forms of Schools. With Maps and Plans. By R. F. HORTON, M.A., *Fellow and Lecturer of New College, Oxford.* Crown 8vo. 3s. 6d. [Now ready.]
- A History of Hellas, from the Earliest Times to the Death of Alexander.** For the use of Upper Forms of Schools. In two vols. By EVELYN ABBOTT, M.A., LL.D., *Fellow and Tutor of Balliol College, Oxford.* Crown 8vo. [In preparation.]
- History of the Romans to the Establishment of Imperialism.** By J. S. REID, LL.M., *Fellow and Assistant Tutor of Gonville and Caius College, Cambridge; Classical Examiner in the University of London.* [In preparation.]
- A History of England.** For the Use of Middle Forms of Schools. With Maps and Plans. By F. YORK POWELL, M.A., *Senior Student of Christ Church, Oxford, and J. M. MACKAY, M.A., Professor of History at Liverpool College.* In two parts, and also in one volume. Crown 8vo.
- PART I.—From the EARLIEST TIMES to the DEATH of HENRY VII. By F. YORK POWELL, M.A. [Just ready.]
- PART II.—From the DEATH of HENRY VII. to the PRESENT TIME. By J. M. MACKAY, M.A. [In the press.]
- A History of England.** For the use of Middle Forms of Schools. With Contents, Tables, Plans, Maps, Index, &c. By CYRIL RANSOME, M.A., *Professor of Modern Literature and History, Yorkshire College, Leeds.* Small 8vo. [In preparation.]
- Viri illustres Urbis Romae.** An Elementary Latin Reading Book. With Notes and Vocabulary. By G. L. BENNETT, M.A., *Head Master of Sutton Valence School.* Small 8vo. 1s. 6d. [Now ready.]
- Fabulae Faciles.** A First Latin Reader. Containing Detached Sentences and Consecutive Stories. With Notes and a Vocabulary. By F. RITCHIE, M.A., *The Beacon, Sevenoaks, and late Assistant Master in the High School, Plymouth.* Crown 8vo. 2s. 6d. [Now ready.]
- A German Exercise Book.** Containing about 150 Exercises, with the necessary Accidence and Syntax, and a Vocabulary. By W. G. GUILLEMARD, M.A., *Assistant Master at Harrow School.* [In preparation.]

Waterloo Place, London.



**New Books in Preparation and in the Press—continued.**

**Scott's Marmion.** With Introduction, Notes, &c. By F. S. ARNOLD, M.A., *Assistant Master at Bedford Grammar School.* Forming a Volume of "English School Classics." *Small 8vo.*  
[In preparation.]

**Builders' Work and the Building Trades.** By Lieut.-Colonel H. C. SEDDON, R.E., *Examiner in Building Construction, Science and Art Department, South Kensington; Assistant Examiner, H.M. Civil Service Commissioners; Instructor in Construction, School of Military Engineering, Chatham.*  
[In the press.]

This volume contains descriptions of work performed by the different Building Trades; the ordinary Tools and Materials used; methods of executing, measuring, and valuing Builders' work; together with useful data and practical hints connected with the same.

**Highways of History.** A Series of Volumes on portions of English History by various writers. Edited by LOUISE CREIGHTON, *Author of "A First History of England," "Stories from English History," &c.* *Small 8vo.*  
[In the press.]

THE GOVERNMENT OF ENGLAND. *1s. 6d.*

THE CONNECTION BETWEEN ENGLAND AND SCOTLAND. *1s. 6d.*

THE HISTORY OF RELIGION IN ENGLAND. *1s. 6d.*

THE RELATIONS OF ENGLAND WITH FOREIGN POWERS.

ENGLAND AND IRELAND. *1s. 6d.*

THE SOCIAL HISTORY OF ENGLAND.

THE GROWTH OF THE ENGLISH COLONIES. *1s. 6d.*

The aim of this series is to give consecutive treatment to certain important subjects in English History. A strictly chronological arrangement of history distracts the reader's attention from one subject to another. Though England's progress has to be studied by reading her history as a whole, yet clearness is gained on many points by a separate survey of some important line of advance. The treatment of the subjects in this series will be adapted for the use of students who have already mastered the general outline of English History, but wish for more connected information with regard to some special point. Each part will be complete in itself, and will treat of its own subject without reference, except where necessary, to the general course of events.

**Greek Passages adapted for practice in Unseen Translation.** Intended for the use of Middle and Higher Forms of Schools, and for University and other Students. By F. D. MORICE, M.A., *Assistant Master at Rugby School, and Fellow of Queen's College, Oxford.*  
[In preparation.]

**Latin Prose Exercises.** Consisting of about 250 easy Passages for Translation. By T. L. PAPILLON, M.A., *Fellow and Tutor of New College, Oxford.*  
[In preparation.]

**Bacon's Essays.** Complete Edition. By FRANCIS STORR, B.A. Forming a Volume of "English School Classics." *Small 8vo.*  
[Nearly ready.]

Waterloo Place, London,



## KEYS

KEYS are published to the following Educational Works *for the use of Tutors only*. They can *only* be obtained by direct application to the Publishers, who will send a printed Form, to be filled up by the Tutor requiring the KEY. They cannot be supplied through Booksellers.

		Nett Price of the Key.	
		s.	d.
ABBOTT'S	Arnold's Greek Prose Composition . . . . .	3	6
AINGER'S	Clivus. Parts I. and II. . . . .	3	6
ARNOLD'S	Henry's First Latin Book . . . . .	1	0
—	Second Latin Book . . . . .	2	0
—	First Verse Book . . . . .	1	0
—	Latin Prose Composition. Part I. . . . .	1	6
—	First Greek Book . . . . .	1	6
—	Greek Prose Composition. Part I. . . . .	1	6
BENNETT'S	First Latin Writer . . . . .	5	0
—	Second Latin Writer . . . . .	5	0
—	Easy Latin Stories for Beginners . . . . .	5	0
—	Second Latin Reading Book . . . . .	5	0
—	Passages for Latin Unseen Translation . . . . .	3	6
BRADLEY'S	Aids to Writing Latin Prose . . . . .	5	0
—	Arnold's Latin Prose Composition . . . . .	5	0
GEPP'S	Arnold's Henry's First Latin Book . . . . .	5	0
—	Exercises in Latin Elegiac Verse . . . . .	5	0
HEATLEY AND KINGDON'S	Gradatim . . . . .	5	0
—	Excerpta Facilia . . . . .	5	0
HEATLEY'S	Graecula . . . . .	5	0
MORICE'S	Arnold's First Greek Book . . . . .	5	0
RAVEN'S	Latin Grammar Papers . . . . .	5	0
—	Versiculi . . . . .	5	0
RITCHIE AND MOORE'S	Greek Method . . . . .	5	0
RITCHIE'S	First Steps in Latin . . . . .	3	6
SARGENT AND DALLIN'S	Materials and Models for Latin Prose Composition. Latin Version. 116 Selected Pieces . . . . .	5	0
—	Greek Prose Composition. Greek Version. 92 Pieces . . . . .	7	6
SIDGWICK'S	First Greek Writer . . . . .	5	0
—	Greek Prose Composition . . . . .	5	0
SIDGWICK AND MORICE'S	Greek Verse Composition . . . . .	5	0
<i>Keys to the following are sold to the Public without restriction.</i>			
ARNOLD'S	First German Book . . . . .	2	6
—	First French Book . . . . .	2	6
—	First Italian Book . . . . .	1	6
—	First Hebrew Book . . . . .	3	6
SMITH'S (J. HAMBLIN)	Elementary Algebra . . . . .	9	0
—	Arithmetic . . . . .	9	0
—	Geometry . . . . .	8	6
—	Statics and Hydrostatics . . . . .	6	0
—	Trigonometry . . . . .	7	6
—	Latin Prose Composition . . . . .	5	0
SPRATT AND PRETOR'S	Latin and Greek Translation at Sight . . . . .	4	6

Waterloo Place, London.

## ENGLISH

*Select Plays of Shakspeare.* RUGBY EDITION.*With Introduction and Notes to each Play. Small 8vo.*

- As You Like It. 2s.      King Lear. 2s. 6d.  
 Hamlet. 2s. 6d.      Macbeth. 2s.  
 Romeo and Juliet. 2s.      King Henry the Fifth. 2s.  
 A Midsummer Night's Dream. 2s.      King John. 2s.  
*Edited by C. E. MOBERLY, M.A., formerly Scholar of Balliol College, Oxford.*  
 Coriolanus. 2s. 6d. *Edited by ROBERT WHITELAW, M.A., Assistant Master at Rugby School.*  
 The Tempest. 2s. *Edited by J. SURTEES PHILLPOTTS, M.A., Head Master of Bedford Grammar School.*

*Small 8vo. 2s.*

*Shakspeare's Othello.* With Introduction and Notes. *Edited by E. K. PURNELL, M.A., Assistant Master at Wellington College.*

*Highways of History.* A Series of Volumes on portions of English History, by various writers. *Edited by LOUISE CREIGHTON, Author of "A First History of England," &c. Small 8vo.*

- The Government of England. 1s. 6d.  
 The Connection between England and Scotland. 1s. 6d.  
 The History of Religion in England. 1s. 6d.  
 The Relations of England with Foreign Powers. England and Ireland. 1s. 6d.  
 The Social History of England.  
 The Growth of the English Colonies. 1s. 6d.

Waterloo Place, London.

## *English School Classics*

*With Introductions, and Notes at the end of each Book.*

Edited by FRANCIS STORR, B.A.,

CHIEF MASTER OF MODERN SUBJECTS AT MERCHANT TAYLORS' SCHOOL.

*Small 8vo.*

**Thomson's Seasons: Winter.** *With Introduction to the Series, by the REV. J. FRANCK BRIGHT, D.D., Master of University College, Oxford.* 1s.

**Cowper's Task.** By FRANCIS STORR, B.A. 2s.  
Books I. and II., 9d.; Books III. and IV., 9d.; Books V. and VI., 9d.

**Cowper's Simple Poems.** *With Life of the Author.* By FRANCIS STORR, B.A. 1s.

**Scott's Lay of the Last Minstrel.** By J. SURTEES PHILLPOTTS, M.A., *Head Master of Bedford School.* 2s. 6d.  
Canto I., 9d.; Cantos II. and III., 9d.; Cantos IV. and V., 9d.; Canto VI., 9d.

**Scott's Lady of the Lake.** By R. W. TAYLOR, M.A., *Head Master of Kelly College, Tavistock.* 2s.  
Cantos I. and II., 9d.; Cantos III. and IV., 9d.; Cantos V. and VI., 9d.

**Scott's Marmion.** By F. S. ARNOLD, M.A., *Assistant Master at Bedford School.* [In preparation.]

**Notes to Scott's Waverley.** By H. W. EVE, M.A., *Head Master of University College School, London.* 1s.; or with the Text, 2s. 6d.

**Bacon's Essays.** Complete Edition. By FRANCIS STORR, B.A. [In the press.]

**Twenty of Bacon's Essays.** By FRANCIS STORR, B.A. 1s.

**Simple Poems.** Edited by W. E. MULLINS, M.A., *Assistant Master at Marlborough College.* 8d.

**Selections from Wordsworth's Poems.** By H. H. TURNER, B.A., *late Scholar of Trinity College, Cambridge.* 1s.

**Wordsworth's Excursion: The Wanderer.** By H. H. TURNER, B.A. 1s.

**Milton's Paradise Lost.** By FRANCIS STORR, B.A. Book I., 9d. Book II., 9d.

**Milton's L'Allegro, Il Penseroso, and Lycidas.** By EDWARD STORR, M.A., *late Scholar of New College, Oxford.* 1s.

**Selections from the Spectator.** By OSMOND AIRY, M.A., *H. M. Inspector of Schools.* 1s.

Waterloo Place, London.



Browne's Religio Medici. By W. P. SMITH, M.A., *Assistant Master at Winchester College.* 1s.

Goldsmith's Traveller and Deserted Village. By C. SANKEY, M.A., *Head Master of Bury St. Edmund's Grammar School.* 1s.

Extracts from Goldsmith's Vicar of Wakefield. By C. SANKEY, M.A. 1s.

Poems selected from the Works of Robert Burns. By A. M. BELL, M.A., *Balliol College, Oxford.* 2s.

Macaulay's Essays.

MOORE'S LIFE OF BYRON. By FRANCIS STORR, B.A. 9d.  
BOSWELL'S LIFE OF JOHNSON. By FRANCIS STORR, B.A. 9d.

HALLAM'S CONSTITUTIONAL HISTORY. By H. F. BOYD, *late Scholar of Brasenose College, Oxford.* 1s.

Southey's Life of Nelson. By W. E. MULLINS, M.A. 2s. 6d.

Gray's Poems. Selection from Letters, with Life by Johnson. By FRANCIS STORR, B.A. 1s.

*New Edition. Small 8vo. 2s. 6d.*

**The Rudiments of English Grammar and Composition.** By J. HAMBLIN SMITH, M.A., *of Gonville and Caius College, and late Lecturer at St. Peter's College, Cambridge.*

*New and Revised Edition. Small 8vo. 1s. 6d.*

**A Primer of English Parsing and Analysis.** By CYRIL L. C. LOCKE, M.A., *Assistant Master at Clifton College.*

*Small 8vo. 1s. 6d.*

**The Beginner's Drill-book of English Grammar.** Adapted for Middle Class and Elementary Schools. By JAMES BURTON, B.A., *First English Master in the High School of the Liverpool Institute.*

*Small 8vo. 2s. 6d.*

**A Practical English Grammar.** For Schools and Colleges, and for Students preparing for examinations. By the Rev. W. TIDMARSH, B.A., *late Head Master of Putney School.*

*Small 8vo. 2s. 6d.*

**Short Readings in English Poetry.** Arranged, with occasional Notes, for the use of Schools and Classes. Edited by H. A. HERTZ.

Waterloo Place, London.

## HISTORY

*With Maps and Plans. New Edition, Revised. Crown 8vo.*

**A History of England.** *By the Rev. J. FRANCK BRIGHT, D.D., Master of University College, Oxford, and late Master of the Modern School at Marlborough College.*

PERIOD I.—MEDIÆVAL MONARCHY: The departure of the Romans, to Richard III. From A.D. 449 to A.D. 1485. 4s. 6d.

PERIOD II.—PERSONAL MONARCHY: Henry VII. to James II. From A.D. 1485 to A.D. 1688. 5s.

PERIOD III.—CONSTITUTIONAL MONARCHY: William and Mary, to the present time. From A.D. 1689 to A.D. 1837. 7s. 6d.

*Crown 8vo. 6s.*

**The Rise of Constitutional Government in England.**

*By CYRIL RANSOME, M.A., Professor of Modern Literature and History, Yorkshire College, Leeds.*

**A History of England.** *For the Use of Middle Forms of Schools. With Maps and Plans. By F. YORK POWELL, M.A., Senior Student of Christ Church, Oxford; and J. M. MACKAY, M.A., Professor of History at Liverpool College. In two parts, and also in one volume. Crown 8vo.*

PART I.—From the EARLIEST TIMES to the DEATH of HENRY VII. *By F. YORK POWELL, M.A. [Just ready.*

PART II.—From the DEATH of HENRY VII. to the PRESENT TIME. *By J. M. MACKAY, M.A. [In the press.*

*Second Edition. With Forty Illustrations. 16mo. 2s. 6d.*

**A First History of England.** *By LOUISE CREIGHTON, Author of "Life of the Black Prince," "Sir Walter Raleigh," &c.*

*With numerous Illustrations. Royal 16mo. 3s. 6d.*

**Stories from English History.** *By LOUISE CREIGHTON, Author of "A First History of England," "Life of the Black Prince," &c.*

*New Edition. 18mo. 1s. 6d.*

**A History of England for Children.** *By GEORGE DAVYS, D.D., formerly Bishop of Peterborough.*

---

Waterloo Place, London.

***Historical Handbooks.*** Edited by OSCAR BROWNING,  
M.A., *Fellow of King's College, Cambridge.*

*Crown 8vo.*

- English History in the XIVth Century.** By CHARLES  
H. PEARSON, M.A., *late Fellow of Oriel College, Oxford.* 3s. 6d.
- The Reign of Lewis XI.** By P. F. WILLERT, M.A., *Fellow of  
Exeter College, Oxford.* With Map. 3s. 6d.
- The Roman Empire. A.D. 395-800.** By A. M. CURTEIS,  
M.A. With Maps. 3s. 6d.
- History of the English Institutions.** By PHILIP V.  
SMITH, M.A., *Fellow of King's College, Cambridge.* 3s. 6d.
- History of Modern English Law.** By Sir R. K. WILSON,  
Bart., M.A., *late Fellow of King's College, Cambridge.* 3s. 6d.
- History of French Literature.** Adapted from the French of  
M. DEMOGEOT, by C. BRIDGE. 3s. 6d.

*With Maps and Plans. Crown 8vo. 3s. 6d.*

***A History of the Romans.*** For the use of Middle Forms  
of Schools. By R. F. HORTON, M.A., *Fellow and Lecturer of  
New College, Oxford.*

*Crown 8vo. 7s. 6d.*

***Chapters in the History of English Literature.***  
*From 1509 to the close of the Elizabethan Period.*  
By ELLEN CROFTS, *Lecturer at Newnham College, Cambridge.*

*Crown 8vo. 7s. 6d.*

***Ecclesia Anglicana.*** A History of the Church of Christ in  
England from the earliest to the present times. By ARTHUR  
CHARLES JENNINGS, M.A., *Jesus College, Cambridge; Vicar of  
Whittlesford.*

*Second Edition, Revised. Crown 8vo. 7s. 6d.*

***History of the Church under the Roman Empire,***  
A.D. 30-476. By the Rev. A. D. CRAKE, B.A., *Vicar of Haven  
Street, Ryde.*

Waterloo Place, London.



**Historical Biographies.** Edited by the Rev. M. CREIGHTON, M.A., LL.D., late Fellow and Tutor of Merton College, Oxford. With Maps and Plans. Small 8vo.

Simon de Montfort. By M. CREIGHTON, M.A., LL.D. 2s. 6d.

The Black Prince. By LOUISE CREIGHTON. 2s. 6d.

Sir Walter Raleigh. By LOUISE CREIGHTON. 3s.

Oliver Cromwell. By F. W. CORNISH, M.A. 3s. 6d.

The Duke of Marlborough. By LOUISE CREIGHTON. 3s. 6d.

The Duke of Wellington. By ROSAMOND WAITE. 3s. 6d.

Second Edition. Crown 8vo. 6s.

**A Handbook in Outline of the Political History of England to 1882.** Chronologically arranged. By A. H. DYKE ACLAND, M.A., Christ Church, Oxford, and CYRIL RANSOME, M.A., Professor of Modern Literature and History, Yorkshire College, Leeds.

Crown 8vo. Paper cover, 1s. Cloth limp, 1s. 6d.

**A Handbook in Outline of English Politics for the Last Half Century.** Extracted from "A Handbook of English Political History." With Appendices on the Reform Bills, Disfranchised and Enfranchised Boroughs, &c. By A. H. DYKE ACLAND, M.A., and CYRIL RANSOME, M.A.

Small 8vo. 1s. 6d.

**A Skeleton Outline of the History of England,** being an abridgment of a Handbook in Outline of the Political History of England. By A. H. DYKE ACLAND, M.A., and CYRIL RANSOME, M.A.

Small 8vo. 2s. 6d.

**A Skeleton Outline of Greek History.** Chronologically arranged. By EVELYN ABBOTT, M.A., LL.D., Fellow and Tutor of Balliol College, Oxford.

Small 8vo. 2s.

**A Skeleton Outline of Roman History.** Chronologically arranged. By P. E. MATHESON, M.A., Fellow of New College, Oxford.

Waterloo Place, London.



## SCIENCE

*Third and Enlarged Edition, With Illustrations. 8vo. 21s.*

**Physical Geology for Students and General Readers.** By A. H. GREEN, M.A., F.G.S., Professor of Geology in the Yorkshire College of Science, Leeds.

*Crown 8vo. 2s. 6d.*

**Elementary Course of Practical Physics.** By A. M. WORTHINGTON, M.A., F.R.A.S., Assistant Master at Clifton College.

*New Edition, Revised. With Illustrations. Crown 8vo. 2s. 6d.*

**An Easy Introduction to Chemistry.** Edited by the Rev. ARTHUR RIGG, M.A., and WALTER T. GOOLDEN, M.A., Lecturer in Natural Science at Tonbridge School.

*Second Edition. With Illustrations. Crown 8vo. 5s.*

**A Year's Botany.** Adapted to Home and School Use. By FRANCES ANNE KITCHENER, Illustrated by the Author.

*With Illustrations. Medium 8vo.*

**Notes on Building Construction.**

Arranged to meet the requirements of the syllabus of the Science and Art Department of the Committee of Council on Education, South Kensington.

PART I.—FIRST STAGE, OR ELEMENTARY COURSE. Second Edition. *With 325 woodcuts. 10s. 6d.*

PART II.—COMMENCEMENT OF SECOND STAGE, OR ADVANCED COURSE. Second Edition. *With 300 woodcuts. 10s. 6d.*

PART III.—ADVANCED COURSE. *With 188 woodcuts. 21s.*

REPORT ON THE EXAMINATION IN BUILDING CONSTRUCTION, HELD BY THE SCIENCE AND ART DEPARTMENT, SOUTH KENSINGTON, IN MAY, 1875.—"The want of a text-book on this subject, arranged in accordance with the published syllabus, and therefore limiting the students and teachers to the prescribed course, has lately been well met by a work published by Messrs. Rivingtons, entitled 'Notes on Building Construction,' arranged to meet the requirements of the Syllabus of the Science and Art Department of the Committee of Council on Education, South Kensington."

(Signed) H. C. SEDDON, Major R.E.  
[Instructor in Construction and Estimating at the School of Military Engineering, Chatham.]

June 18, 1875.

Waterloo Place, London.

## MATHEMATICS

### *Rivington's Mathematical Series.*

*Small 8vo. 3s. Without Answers, 2s. 6d.*

**Elementary Algebra.** *By J. HAMBLIN SMITH, M.A., of Gonville and Caius College, and late Lecturer in Classics at St. Peter's College, Cambridge.*

Also an Edition especially prepared to cover the ground required by the Regulations for the University Examinations in India. 3s.

A KEY TO ELEMENTARY ALGEBRA. 9s.

*Small 8vo. 2s. 6d.*

**Exercises on Algebra.** *By J. HAMBLIN SMITH, M.A.*  
(Copies may be had without the Answers.)

*Crown 8vo. 8s. 6d.*

**Algebra. PART II.** *By E. J. GROSS, M.A., Fellow of Gonville and Caius College, Cambridge, and Secretary to the Oxford and Cambridge Schools Examination Board.*

*Small 8vo. 3s. 6d.*

**A Treatise on Arithmetic.** *By J. HAMBLIN SMITH, M.A.*  
(Copies may be had without the Answers.)

A KEY TO ARITHMETIC. 9s.

*Small 8vo. 4s. 6d.*

**Elementary Trigonometry.** *By J. HAMBLIN SMITH, M.A.*

A KEY TO ELEMENTARY TRIGONOMETRY. 7s. 6d.

*Crown 8vo. 5s. 6d.*

**Kinematics and Kinetics.** *By E. J. GROSS, M.A.*

*Crown 8vo. 4s. 6d.*

**Geometrical Conic Sections.** *By G. RICHARDSON, M.A., Assistant Master at Winchester College.*

*Small 8vo. 3s.*

**Elementary Statics.** *By J. HAMBLIN SMITH, M.A.*

*Small 8vo. 3s.*

**Elementary Hydrostatics.** *By J. HAMBLIN SMITH, M.A.*

*Crown 8vo. 6s.*

**A Key to Elementary Statics and Hydrostatics.**  
*By J. HAMBLIN SMITH, M.A.*

---

Waterloo Place, London.

*Small 8vo. 3s. 6d.*

**Elements of Geometry.** By J. HAMBLIN SMITH, M.A.

Containing Books 1 to 6, and portions of Books 11 and 12, of EUCLID, with Exercises and Notes, arranged with the Abbreviations admitted in the Cambridge University and Local Examinations. Books 1 and 2, limp cloth, 1s. 6d., may be had separately.

*Crown 8vo. 8s. 6d.*

**A Key to Elements of Geometry.** By J. HAMBLIN SMITH, M.A.

*Small 8vo. 1s.*

**Book of Enunciations for Hamblin Smith's Geometry, Algebra, Trigonometry, Statics, and Hydrostatics.**

*Small 8vo. 3s.*

**An Introduction to the Study of Heat.** By J. HAMBLIN SMITH, M.A.

*Crown 8vo. 5s.*

**Companion to Algebra.** With numerous Examples. By L. MARSHALL, M.A., *Assistant Master at Charterhouse.*

*Crown 8vo. 6s.*

**The Principles of Dynamics.** An Elementary Text-book for Science Students. By R. WORMELL, D.Sc., M.A., *Head Master of the City of London Middle-Class School.*

*Small 8vo.*

**A Collection of Arithmetical Exercises,** progressively arranged. By A. E. DONKIN, M.A., and C. H. HODGES, M.A., *Mathematical Masters at Rugby School.*

*Small 8vo. 3s. 6d.*

**Army and Civil Service Examination Papers** in Arithmetic, including Mensuration and Logarithms. With Arithmetical Rules, Tables, Formulæ and Answers, for the use of Students preparing for Examination. By the Rev. A. DAWSON CLARKE, M.A., *St. John's College, Cambridge.*

*New Edition, Revised. Crown 8vo. 6s. 6d.*

**Arithmetic, Theoretical and Practical.** By W. H. GIRDLESTONE, M.A., *of Christ's College, Cambridge.*

Also a School Edition. *Small 8vo. 3s. 6d.*

Waterloo Place, London,



**Latin Composition Books.**

- BENNETT'S First Latin Writer, *p.* 15.  
 ——— First Latin Exercises, *p.* 15.  
 ——— Second Latin Writer, *p.* 15.  
 RITCHIE'S First Steps in Latin, *p.* 16.  
 DAWE'S Beginner's Latin Exercise Book, *p.* 16.  
 ARNOLD'S Henry's First Latin Book, *p.* 18.  
 GEPP'S Arnold's Henry's First Latin Book, *p.* 18.  
 PROWDE SMITH'S Latin Prose Exercises, *p.* 19.  
 HAMBLIN SMITH'S Latin Exercises, *p.* 19.  
 ARNOLD'S Latin Prose, *p.* 18.  
 BRADLEY'S Arnold's Latin Prose, *p.* 18.  
 ——— Aids to Latin Prose, *p.* 18.  
 SARGENT AND DALLIN'S Materials,

**Greek Composition Books.**

- SIDGWICK'S First Greek Writer, *p.* 23  
 ARNOLD'S First Greek Book, *p.* 23.  
 MORICE'S Arnold's First Greek Book, *p.* 23.  
 RITCHIE AND MOORE'S Greek Method, *p.* 22.  
 ARNOLD'S Greek Prose, *p.* 24.  
 ABBOTT'S Arnold's Greek Prose, *p.* 24.  
 SIDGWICK'S Greek Prose, *p.* 23.  
 SARGENT AND DALLIN'S Materials, *p.* 25.

**Latin Reading Books.**

- BENNETT'S Easy Latin Stories, *p.* 15.  
 ——— Viri Illustres, *p.* 15.  
 ——— Second Latin Reading Book, *p.* 15.  
 ——— Selections from Vergil and Caesar, *p.* 15.  
 RITCHIE'S Fabulae Faciles, *p.* 16.  
 HEATLEY AND KINGDON'S Gradatim, *p.* 16.  
 ——— Excerpta Facilia, *p.* 16.  
 TAYLOR'S Stories from Ovid, *p.* 19.

**Greek Reading Books.**

- HEATLEY'S Græcula, *p.* 22.  
 MORICE'S Stories in Attic Greek, *p.* 26.  
 PHILLPOTT'S Stories from Herodotus, *p.* 25.  
 MOORE'S Selections from Thucydides, *p.* 2.  
 ABBOTT'S Selections from Lucian, *p.* 25.  
 MOBERLY'S Alexander the Great, *p.* 26.  
 SIDGWICK'S Scenes from Greek Plays, *p.* 26.

**Latin and Greek Unseen Translation.**

- BENNETT'S Easy Latin Passages, *p.* 16.  
 TURNER'S Latin and Greek Passages, *p.* 2.  
 SARGENT'S Latin Passages, *p.* 20.  
 SPRATT AND PRETOR'S Latin and Greek Passages, *p.* 20.

---

Waterloo Place, London.

## LATIN

*New Edition, Revised. Crown 8vo. 3s. 6d.*

**First Latin Writer.** Comprising Accidence, the Easier Rules of Syntax illustrated by copious Examples, and progressive Exercises in Elementary Latin Prose, with Vocabularies. *By G. L. BENNETT, M.A., Head Master of Sutton Valence School.*

A KEY for the use of Tutors only. 5s.

*Crown 8vo. 2s. 6d.*

**First Latin Exercises.** Being the Exercises, with Syntax Rules and Vocabularies, from a "First Latin Writer." *By G. L. BENNETT, M.A.*

*Crown 8vo. 1s. 6d.*

**Latin Accidence.** From a "First Latin Writer." *By G. L. BENNETT, M.A.*

*New Edition. Crown 8vo. 3s. 6d.*

**Second Latin Writer.** *By G. L. BENNETT, M.A.*

A KEY for the use of Tutors only. 5s.

*Small 8vo.*

**Viri Illustres Urbis Romae.** An Elementary Latin Reading Book. With Notes and Vocabulary. *By G. L. BENNETT, M.A.*

*New Edition, Revised. Crown 8vo. 2s. 6d.*

**Easy Latin Stories for Beginners.** With Vocabulary and Notes. *By G. L. BENNETT, M.A.*

A KEY for the use of Tutors only. 5s.

*Crown 8vo. 2s. 6d.*

**Second Latin Reading Book.** Forming a continuation of "Easy Latin Stories for Beginners." *By G. L. BENNETT, M.A.*

A KEY for the use of Tutors only. 5s.

*Small 8vo. 2s.*

**Selections from Caesar. The Gallic War.** With Notes, Map, &c. *By G. L. BENNETT, M.A.*

*Small 8vo. 1s. 6d.*

**Selections from the Aeneid of Vergil.** With Notes, &c. *By G. L. BENNETT, M.A.*

---

Waterloo Place, London,

16mo. Paper cover, 1s. Cloth, 1s. 6d.

**Easy Graduated Latin Passages.** For Translation into English, for use in Schools as Unseen Pieces. By G. L. BENNETT, M.A., *Head Master of Sutton Valence School.*

A KEY for the use of Tutors only. 3s. 6d.

Crown 8vo. 2s. 6d.

**Fabulae Faciles.** A First Latin Reader. Containing Detached Sentences and Consecutive Stories. With Notes and a Vocabulary. By F. RITCHIE, M.A., *The Beacon, Sevenoaks, and late Assistant Master in the High School, Plymouth.*

Small 8vo. 1s. 6d.

**Latin Grammar and Junior Scholarship Papers.**

By J. H. RAVEN, M.A., *Head Master of Fauconberge School, Beccles, Suffolk.*

A KEY for the use of Tutors only. 5s.

Small 8vo. 2s.

**Easy Latin and Greek Grammar Papers.** Prepared by H. R. HEATLEY, M.A., *Assistant Master at Hillbrow School, Rugby.*

Second Edition. Crown 8vo. 1s. 6d.

**First Steps in Latin.** By F. RITCHIE, M.A., *The Beacon, Sevenoaks, and late Assistant Master in the High School, Plymouth.*

A KEY for the use of Tutors only. 3s. 6d.

New Edition. Small 8vo. 1s. 6d.

**Gradatim.** An Easy Translation Book for Beginners. With Vocabulary. By H. R. HEATLEY, M.A., *Assistant Master at Hillbrow School, Rugby, and H. N. KINGDON, B.A., Head Master of Dorchester Grammar School.*

A KEY for the use of Tutors only. 5s.

Second Edition. Small 8vo. 2s. 6d.

**Excerpta Facilia.** A Second Latin Translation Book, containing a Collection of Stories from various Latin Authors, with Notes at end, and a Vocabulary. By H. R. HEATLEY, M.A., *Assistant Master at Hillbrow School, Rugby, and H. N. KINGDON, B.A., Head Master of Dorchester Grammar School.*

A KEY for the use of Tutors only. 5s.

Second Edition. Crown 8vo. 1s. 6d.

**The Beginner's Latin Exercise Book.** Affording Practice, oral and written, on Latin Accidence. With Vocabulary. By C. J. SHERWILL DAWE, B.A., *Lecturer and Assistant Chaplain at St. Mark's College, Chelsea.*

Waterloo Place, London,



18mo.

**Latin Texts.** For use in schools, &c.

THE AENEID OF VERGIL. BOOKS I. II. III. IV. V. VII.  
VIII. IX. 2d. each. BOOKS VI. X. XI. XII. 3d. each.

THE GEORGICS OF VERGIL. BOOKS I.-IV. 2d. each.

THE BUCOLICS OF VERGIL. 2d.

**Vergil.** The Bucolics, Georgics, and Aeneid in One Volume. 2s. 6d.

CAESAR DE BELLO GALLICO. BOOKS I. V. VII. VIII. 3d. each.  
BOOKS II. III. IV. VI. 2d. each.

**Caesar De Bello Gallico.** In One Volume. 1s. 6d.

Crown 8vo. On a card, 9d.

**Elementary Rules of Latin Pronunciation.** By  
ARTHUR HOLMES, M.A., late Senior Fellow and Dean of Clare  
College, Cambridge.

Sixth Edition. Crown 8vo. 3s. 6d.

**Progressive Exercises in Latin Elegiac Verse.**  
By C. G. GEPP, M.A., Assistant Master at Bradfield College.

A KEY for the use of Tutors only. 5s.

Twelfth Edition. 12mo. 2s.

**A First Verse Book.** Being an Easy Introduction to the Mechan-  
ism of the Latin Hexameter and Pentameter. By THOMAS KER-  
CHEVER ARNOLD, M.A.

A KEY for the use of Tutors only. 1s.

New Edition. Crown 8vo.

**Clivus.** Elementary Exercises in Latin Elegiac Verse. Compiled  
By A. C. AINGER, M.A., Assistant Master at Eton College.

PART I. 2s. 6d. PART II. 2s. 6d.

A KEY for the use of Tutors only. 3s. 6d.

Crown 8vo. 3s. 6d.

**An Elementary Latin Grammar.** By J. HAMBLIN SMITH,  
M.A., of Gonville and Caius College, and late Lecturer in Classics at  
St. Peter's College, Cambridge.

---

Waterloo Place, London.



*Twenty-sixth Edition. 12mo. 3s.*

**Henry's First Latin Book.** *By* T. K. ARNOLD, M.A.,

A KEY for the use of Tutors only. 1s.

*New Edition, Revised. 12mo. 3s.*

**Arnold's Henry's First Latin Book.** *By* C. G. GEPP, M.A.,  
*Assistant Master at Bradfield College, Author of "Progressive Exercises in Latin Elegiac Verse."*

A KEY for the use of Tutors only. 5s.

*Twenty-first Edition. 8vo. 6s. 6d.*

**A Practical Introduction to Latin Prose Composition.** *By* THOMAS KERCHEVER ARNOLD, M.A.

A KEY for the use of Tutors only. 1s. 6d.

*New Edition, Revised. Crown 8vo. 5s.*

**Arnold's Practical Introduction to Latin Prose Composition.** *By* G. GRANVILLE BRADLEY, D.D., *Dean of Westminster, late Master of University College, Oxford, and formerly Master of Marlborough College.*

A KEY for the use of Tutors only. 5s.

*Crown 8vo. 5s.*

**Aids to Writing Latin Prose.** Containing 144 Exercises, with an Introduction comprising Preliminary Hints, Directions, Explanatory Matter, &c. *By* G. G. BRADLEY, D.D., *Dean of Westminster, and* T. L. PAPILLON, M.A., *Fellow and Tutor of New College, Oxford.*

A KEY for the use of Tutors only. 5s.

*Crown 8vo.*

**The Aeneid of Vergil.** *Edited, with Notes at the end, by* FRANCIS STORR, B.A., *Chief Master of Modern Subjects at Merchant Taylors' School.*

BOOKS I. and II. 2s. 6d. BOOKS XI. and XII. 2s. 6d.

*Small 8vo. 1s. 6d.*

**Virgil, Georgics.** BOOK IV. *Edited, with Life, Notes, Vocabulary, and Index, by* C. G. GEPP, M.A., *Assistant Master at Bradfield College.*

*Third Edition, Revised. Crown 8vo. 3s. 6d.*

**Stories from Ovid in Elegiac Verse.** With Notes and Marginal References to the "Public School Latin Primer." By R. W. TAYLOR, M.A., *Head Master of Kelly College, Tavistock.*

*New Edition, Revised. Crown 8vo. 2s. 6d.*

**Stories from Ovid in Hexameter Verse. Metamorphoses.** With Notes and Marginal References to the "Public School Latin Primer." By R. W. TAYLOR, M.A.

*New Edition, Revised. 12mo. 2s. 6d.*

**Eclogæ Ovidianæ.** From the Elegiac Poems. With English Notes. By THOMAS KERCHEVER ARNOLD, M.A.

*Second Edition. Small 8vo. 2s.*

**Cicero de Amicitia.** Edited, with Introduction and Notes, by ARTHUR SIDGWICK, M.A., *Fellow and Tutor of Corpus Christi College, Oxford.*

*Small 8vo. 3s. 6d.*

**Cæsar. De Bello Gallico.** BOOKS I.-III. Edited by J. H. MERRYWEATHER, M.A., and C. C. TANCOCK, M.A., *Assistant Masters at Charterhouse.*

BOOK I. separately. 2s.

*Crown 8vo. 3s. 6d.*

**Exercises on the Elementary Principles of Latin Prose Composition.** By J. HAMBLIN SMITH, M.A., *of Gonville and Caius College, and late Lecturer in Classics at St. Peter's College, Cambridge.* A KEY. 5s.

*Small 8vo. 3s. 6d.*

**Livy.** BOOK II. Chiefly from the text of MADVIG, with Notes, Translations, and Appendices. Edited by HENRY BELCHER, M.A., *Master of the Matriculation Class, King's College School, London.*

*Fourth Edition. Crown 8vo. 2s. 6d.*

**Latin Prose Exercises.** For Beginners, and Junior Forms of Schools. By R. PROWDE SMITH, B.A., *Assistant Master at Cheltenham College.*

---

Waterloo Place, London.

*Crown 8vo. 2s. 6d.*

**Latin Passages adapted for Practice in Unseen Translation.**

For the use of Middle Forms of Schools. By J. Y. SARGENT, M.A., *Fellow and Tutor of Hertford College, Oxford*; and Editor of "Materials and Models for Latin and Greek Prose Composition."

*New Edition, re-arranged, with fresh Pieces and additional References.*

*Crown 8vo. 6s. 6d.*

**Materials and Models for Latin Prose Composition.**

Selected and arranged by J. Y. SARGENT, M.A., *Fellow and Tutor of Hertford College, Oxford*, and T. F. DALLIN, M.A., *late Tutor and Fellow of Queen's College, Oxford*.

A KEY to SELECTED PIECES (116), for the use of Tutors only. 5s.

*Crown 8vo.*

**Exercises in Translation at Sight.**

A Selection of Passages from Greek and Latin Authors. For the use of Students. Arranged and translated by A. W. SPRATT, M.A., and A. PRETOR, M.A., *Fellows of St. Catharine's College, Cambridge*.

Vol. I.—THE ORIGINAL PASSAGES. 4s. 6d.

Vol. II.—THE ENGLISH VERSIONS. 4s. 6d.

16mo.

**A Latin-English Dictionary for Junior Forms of Schools.**

By C. G. GEPP, M.A., *Assistant Master at Bradfield College*. [In the press.]

This work aims at supplying in a concise form and at a low price all the information required by boys in Middle Class Schools, or in the Junior Forms of Public Schools. Archaisms (with the exception of such as occur in the most commonly read authors), words peculiar to Plautus, and words found only in late or ecclesiastical Latin, have been accordingly excluded. On the other hand, Proper Names have been briefly yet adequately treated in alphabetical order in the body of the work. No effort has been spared to ensure completeness and accuracy, all references having been verified from the latest and most approved editions of modern scholars.

*8vo. On a Card, 1s.*

**Outlines of Latin Sentence Construction.**

By E. D. MANSFIELD, M.A., *Assistant Master at Clifton College*.

*Second Edition. Crown 8vo. 7s. 6d.*

**Classical Examination Papers.**

Edited, with Notes and References, by P. J. F. GANTILLON, M.A., *Classical Master at Cheltenham College*.

Or, interleaved with writing-paper, half-bound, 10s. 6d.

Waterloo Place, London.



*Crown 8vo.*

**Terenti Comædiæ.** Edited by T. L. PAPILLON, M.A., *Fellow and Tutor of New College, Oxford.*

ANDRIA ET EUNUCHUS. With Introduction on Prosody. 4s. 6d.

Or separately, ANDRIA. 3s. 6d. EUNUCHUS. 3s.

*Crown 8vo. 5s.*

**Juvenalis Satiræ.** THIRTEEN SATIRES. Edited by G. A. SIMCOX, M.A., *Fellow of Queen's College, Oxford.*

*Crown 8vo. 3s. 6d.*

**Persii Satiræ.** Edited by A. PRETOR, M.A., *Fellow of St. Catharine's College, Cambridge.*

*Crown 8vo. 7s. 6d.*

**Horati Opera.** By J. M. MARSHALL, M.A., *Head Master of Durham School.*

VOL. I.—THE ODES, CARMEN SECULARE, AND EPODES.

Also separately, THE ODES. BOOKS I. to IV. 1s. 6d. each.

*Crown 8vo.*

**Taciti Historiæ.** Edited by W. H. SIMCOX, M.A., *Fellow of Queen's College, Oxford.*

BOOKS I. and II., 6s. BOOKS III., IV., and V., 6s.

*Crown 8vo. Paper cover, 1s. each.*

**Plays of the Oratory School, Birmingham.**

TERENCE—ANDRIA—PHORMIO—PINCERNA.

With English Notices to assist the representation.

---

Waterloo Place, London.

## GREEK

*New Edition, Revised. Crown 8vo. 3s. 6d.*

**A Primer of Greek Grammar.** With a Preface by JOHN PERCIVAL, M.A., LL.D., *President of Trinity College, Oxford; late Head Master of Clifton College.*

*Or separately, crown 8vo. 2s. 6d.*

**Accidence.** By EVELYN ABBOTT, M.A., LL.D., *Fellow and Tutor of Balliol College, Oxford; and E. D. MANSFIELD, M.A., Assistant Master at Clifton College.*

*Crown 8vo. 1s. 6d.*

**Syntax.** By E. D. MANSFIELD, M.A., *Assistant Master at Clifton College.*

*New Edition. Crown 8vo. 3s. 6d.*

**A Practical Greek Method for Beginners.** Being a Graduated application of Grammar to Translation and Composition. By F. RITCHIE, M.A., *The Beacon, Sevenoaks, and late Assistant Master at the High School, Plymouth; and E. H. MOORE, M.A., Assistant Master at the High School, Plymouth.*

A KEY for the use of Tutors only. 5s.

*Crown 8vo. 2s. 6d.*

**A Manual of Greek Verbs.** With Rules for the Formation of Tenses, and Tables of Verbs for Practice. By F. RITCHIE, M.A., *The Beacon, Sevenoaks, and late Assistant Master at the High School, Plymouth; and E. H. MOORE, M.A., Assistant Master at the High School, Plymouth.*

*Small 8vo. 1s. 6d.*

**Graecula.** A First Book of Greek Translation. With Rules, Short Sentences, Stories for Translation, and a Vocabulary. By H. R. HEATLEY, M.A., *Assistant Master at Hillbrow School, Rugby.*

A KEY for the use of Tutors only. 5s.

*Small 8vo. 2s.*

**Easy Latin and Greek Grammar Papers.** For the use of Public and Private Schools. Prepared by H. R. HEATLEY, M.A., *Assistant Master at Hillbrow School, Rugby.*

---

Waterloo Place, London.

*New Edition, Revised. Crown 8vo. 3s. 6d.*

**A First Greek Writer.** By ARTHUR SIDGWICK, M.A., *Fellow and Tutor of Corpus Christi College, Oxford; late Assistant Master at Rugby School, and Fellow of Trinity College, Cambridge.*

A KEY for the use of Tutors only. 5s.

*Fourth Edition, Revised. Crown 8vo. 5s.*

**An Introduction to Greek Prose Composition.** With Exercises. By ARTHUR SIDGWICK, M.A.

A KEY for the use of Tutors only. 5s.

*Second Edition, Revised. Crown 8vo. 5s.*

**An Introduction to Greek Verse Composition.** With Exercises. By ARTHUR SIDGWICK, M.A., *Fellow and Tutor of Corpus Christi College, Oxford; and F. D. MORICE, M.A., Assistant Master at Rugby School, and Fellow of Queen's College, Oxford.*

A KEY for the use of Tutors only. 5s.

*Sixth Edition. 12mo. 5s.*

**The First Greek Book.** On the plan of *Henry's First Latin Book.* By THOMAS KERCHEVER ARNOLD, M.A.

A KEY for the use of Tutors only. 1s. 6d.

*New Edition, Revised. Crown 8vo. 3s. 6d.*

**Arnold's First Greek Book.** By FRANCIS DAVID MORICE, M.A., *Assistant Master at Rugby School, and Fellow of Queen's College, Oxford.*

A KEY for the use of Tutors only. 5s.

*Cloth limp, 8vo. 6d.*

**Elementary Card on Greek Prepositions.** By Rev E. PRIESTLAND, M.A., *Spondon House School, Derbyshire.*

*Crown 8vo. 9d.*

**A Short Greek Syntax.** Extracted from "*Xenophon's Anabasis, with Notes.*" By R. W. TAYLOR, M.A., *Head Master of Kelly College, Tavistock.*



*Crown 8vo. 8s. 6d.*

***A Syntax of Attic Greek for the use of Students and Schools.*** By F. E. THOMPSON, M.A., Assistant Master at Marlborough College.

*Third Edition. Imperial 16mo. 8s. 6d.*

***Madvig's Syntax of the Greek Language, especially of the Attic Dialect.*** For the use of Schools. Edited by THOMAS KERCHEVER ARNOLD, M.A.

*Tenth Edition. 8vo. 5s. 6d.*

***A Practical Introduction to Greek Accidence.*** By THOMAS KERCHEVER ARNOLD, M.A.

*Fourteenth Edition. 8vo. 5s. 6d.*

***A Practical Introduction to Greek Prose Composition.*** By THOMAS KERCHEVER ARNOLD, M.A.

A KEY for the use of Tutors only. 1s. 6d.

*New Edition, Revised. Crown 8vo. 3s. 6d.*

***Arnold's Practical Introduction to Greek Prose Composition.*** By EVELYN ABBOTT, M.A., LL.D., Fellow and Tutor of Balliol College, Oxford.

A KEY for the use of Tutors only. 3s. 6d.

*Crown 8vo. 4s. 6d.*

***Elements of Greek Accidence.*** By EVELYN ABBOTT, M.A., LL.D., Fellow and Tutor of Balliol College, Oxford.

*Crown 8vo. 4s. 6d.*

***An Elementary Greek Grammar.*** By J. HAMBLIN SMITH, M.A., of Gonville and Caius College, and late Lecturer in Classics at St. Peter's College, Cambridge.

*Cloth limp, 8vo. 1s.*

***A Table of Irregular Greek Verbs, classified according to the arrangement of Crusius's Greek Grammar.*** By FRANCIS STORR, B.A., Chief Master of Modern Subjects at Merchant Taylors' School, and late Assistant Master at Marlborough College.

---

Waterloo Place, London.



*Crown 8vo.*

**Exercises in Translation at Sight.** A Selection of Passages from Greek and Latin Authors. For the use of Students. Arranged and Translated by A. W. SPRATT, M.A., and A. PRETOR, M.A., *Fellows of St. Catharine's College, Cambridge.*

Vol. I.—THE ORIGINAL PASSAGES. 4s. 6d.

Vol. II.—THE ENGLISH VERSIONS. 4s. 6d.

*Crown 8vo. 7s. 6d.*

**Etyma Graeca.** An Etymological Lexicon of Classical Greek. By E. R. WHARTON, M.A., *Lecturer and late Fellow of Jesus College, Oxford.*

*Second Edition. Crown 8vo. 7s. 6d.*

**Classical Examination Papers.** Edited, with Notes and References, by P. J. F. GANTILLON, M.A., *Classical Master at Cheltenham College.*

Or interleaved with writing-paper, half-bound, 10s. 6d.

*Second Edition, containing fresh Pieces and additional References.*

*Crown 8vo. 5s.*

**Materials and Models for Greek Prose Composition.**

*Selected and arranged by J. Y. SARGENT, M.A., Fellow and Tutor of Hertford College, Oxford; and T. F. DALLIN, M.A.*

A KEY to SELECTED PIECES (92), for the use of Tutors only. 7s. 6d.

*Crown 8vo. 2s.*

**Iophon: An Introduction to the Art of Writing Greek Iambic Verses.** By the WRITER of "Nuces" and "Lucretillus."

*Fifth Edition. Crown 8vo. 1s. 6d.*

**Stories from Herodotus.** The Tales of Rhampsinitus and Polycrates, and the Battle of Marathon and the Alcmaeonidae. In Attic Greek. Edited by J. SURTEES PHILLPOTTS, M.A., *Head Master of Bedford Grammar School.*

*New Edition, Revised. Small 8vo. 3s. 6d.*

**Selections from Lucian.** With English Notes. By EVELYN ABBOTT, M.A., LL.D., *Fellow and Tutor of Balliol College, Oxford.*

---

Waterloo Place, London.

B\*

*Small 8vo. 1s. 6d. each.*

## **Scenes from Greek Plays.** RUGBY EDITION.

*Abridged and adapted for the use of Schools, by ARTHUR SIDGWICK, M.A., Fellow of Corpus Christi College, Oxford, late Assistant Master at Rugby School, and Fellow of Trinity College, Cambridge.*

### **Aristophanes.**

THE CLOUDS. THE FROGS. THE KNIGHTS. PLUTUS.

### **Euripides.**

IPHIGENIA IN TAURIS. THE CYCLOPS. ION.  
ELECTRA. ALCESTIS. BACCHÆ. HECUBA.

*Fourth Edition. Crown 8vo. 3s. 6d.*

**Stories in Attic Greek.** Forming a Greek Reading Book for the use of Junior Forms in Schools. With Notes and Vocabulary. By FRANCIS DAVID MORICE, M.A., Assistant Master at Rugby School, and Fellow of Queen's College, Oxford.

*New Edition. Crown 8vo.*

**The Anabasis of Xenophon.** Edited, with Preface, Introduction, Historical Sketch, Itinerary, Syntax Rules, Notes, Indices, Vocabularies, and Maps, by R. W. TAYLOR, M.A., Head Master of Kelly College, Tavistock, and late Fellow of St. John's College, Cambridge.

BOOKS I. and II. 3s. 6d. BOOKS III. and IV. 3s. 6d.

Also separately, BOOK I., 2s. 6d.; BOOK II., 2s.

*Crown 8vo. 2s. 6d.*

**Xenophon's Agesilaus.** Edited, with Syntax Rules, and References, Notes, and Indices, by R. W. TAYLOR, M.A.

*Second Edition. Small 8vo. 2s.*

**Xenophon's Memorabilia.** BOOK I., with a few omissions. Edited, with an Introduction and Notes, by C. E. MOBERLY, M.A., formerly Scholar of Balliol College, Oxford.

*New Edition. Small 8vo. 2s.*

**Alexander the Great in the Punjaub.** Adapted from Arrian, Book V. An Easy Greek Reading Book. Edited, with Notes and a Map, by C. E. MOBERLY, M.A., formerly Scholar of Balliol College, Oxford.

Waterloo Place, London.

*Small 8vo.*

**Homer's Iliad.** Edited, with Notes at the end for the Use of Junior Students, by ARTHUR SIDGWICK, M.A., *Fellow of Corpus Christi College, Oxford; late Assistant Master at Rugby School, and Fellow of Trinity College, Cambridge.*

BOOKS I. and II. 2s. 6d.

BOOK XXI. 1s. 6d.

BOOK XXII. 1s. 6d.

*Small 8vo.* 2s.

**Homer without a Lexicon, for Beginners.** ILIAD, BOOK VI. Edited, with Notes giving the meanings of all the less common words, by J. SURTEES PHILLPOTTS, M.A., *Head Master of Bedford Grammar School.*

*Fifth Edition.* 12mo. 3s. 6d.

**Homer for Beginners.** ILIAD, BOOKS I.-III. With English Notes. By THOMAS KERCHEVER ARNOLD, M.A.

*Fifth Edition.* 12mo. 12s.

**The Iliad of Homer.** With English Notes and Grammatical References. By THOMAS KERCHEVER ARNOLD, M.A.

*Crown 8vo.* 6s.

**The Iliad of Homer.** BOOKS I.-XII. From the Text of Dindorf. With Preface and Notes. By S. H. REYNOLDS, M.A., *late Fellow and Tutor of Brasenose College, Oxford.*

*8vo.* 18s.

**Myths of the Odyssey in Art and Literature.** Illustrated with Outline Drawings. By J. E. HARRISON.

*New Edition.* 12mo. 9s.

**A Complete Greek and English Lexicon for the Poems of Homer and the Homeridæ.** By G. CH. CRUSIUS. Translated from the German. Edited by T. K. ARNOLD, M.A.

*Crown 8vo.* 4s. 6d.

**Isocratis Orationes.** AD DEMONICUM ET PANEGYRICUS. Edited by JOHN EDWIN SANDYS, M.A., *Fellow and Tutor of St. John's College, Cambridge, and Public Orator of the University.*



*Third Edition. Crown 8vo. 2s. 6d.*

**Short Notes on the Greek Text of the Gospel of St. Mark.** By J. HAMBLIN SMITH, M.A., of *Gonville and Caius College, Cambridge.*

*Crown 8vo. 4s. 6d.*

**Notes on the Greek Text of the Acts of the Apostles.** By J. HAMBLIN SMITH, M.A., of *Gonville and Caius College, Cambridge.*

*Crown 8vo. 6s.*

**Notes on the Gospel According to S. Luke.** By the Rev. ARTHUR CARR, M.A., *Assistant Master at Wellington College, late Fellow of Oriel College, Oxford.*

*New Edition. 4 vols. 8vo. 102s.*

**The Greek Testament.** With a Critically Revised Text; a Digest of Various Readings; Marginal References to Verbal and Idiomatic Usage; Prolegomena; and a Critical and Exegetical Commentary. For the use of Theological Students and Ministers. By HENRY ALFORD, D.D., *late Dean of Canterbury.*

The Volumes are sold separately, as follows:—

- Vol. I.—THE FOUR GOSPELS. 28s.
- Vol. II.—ACTS TO 2 CORINTHIANS. 24s.
- Vol. III.—GALATIANS TO PHILEMON. 18s.
- Vol. IV.—HEBREWS TO REVELATION. 32s.

*New Edition. 2 vols. Imperial 8vo. 60s.*

**The Greek Testament.** With Notes, Introductions, and Index. By CHR. WORDSWORTH, D.D., *Bishop of Lincoln.*

The Parts may be had separately, as follows:

- THE GOSPELS. 16s.
- THE ACTS. 8s.
- ST. PAUL'S EPISTLES. 23s.
- GENERAL EPISTLES, REVELATION, AND INDEX. 6s.

---

Waterloo Place, London.

## CATENA CLASSICORUM

*Crown 8vo.***Aristophanis Comœdiae.** *By W. C. GREEN, M.A.*

THE ACHARNIANS AND THE KNIGHTS. 4s.

THE WASPS. 3s. 6d. THE CLOUDS. 3s. 6d.

**Demosthenis Orationes Publicae.** *By G. H. HESLOP, M.A.*

THE OLYNTHIACS. 2s. 6d. } or, in One Volume, 4s. 6d.

THE PHILIPPICS. 3s. }

DE FALSA LEGATIONE. 6s.

**Demosthenis Orationes Privatae.** *By A. HOLMES, M.A.*

DE CORONA. 5s.

**Herodoti Historia.** *By H. G. Woods, M.A.*

BOOK I., 6s. BOOK II., 5s.

**Homeri Ilias.** *By S. H. Keynolls, M.A.*

BOOKS I.-XII. 6s.

**Horati Opera.** *By J. M. Marshall, M.A.*

THE ODES, CARMEN SECLARE, AND EPICURES. 7s. 6d.

THE ODES. BOOKS I. to IV. separately. 1s. 6d. each.

**Isocratis Orationes.** *By JOHN EDWIN SANDYS, M.A.*

AD DEMONICUM ET PANEGYRICUS. 4s. 6d.

**Juvenalis Satirae.** *By G. A. SIMON, M.A.* 5s.**Persii Satirae.** *By A. PATER, M.A.* 3s. 6d.**Sophoclis Tragoediae.** *By R. C. JENN, M.A.*

THE ELECTRA. 3s. 6d. THE AJAX. 3s. 6d.

**Taciti Historiae.** *By W. H. SIMON, M.A.*

BOOKS I. and II., 6s. BOOKS III. IV. and V., 6s.

**Terenti Comœdiae.** *By T. L. PARSONS, M.A.*ANDRIA AND EUNUCHUS. With Introduction in Prose. 4s. 6d. *or separately.*

ANDRIA. With Introduction in Prose. 3s. 6d.

EUNUCHUS. 3s.

**Thucydidis Historia.**BOOKS I. and II. *By CHARLES DODD, M.A.* 6s.BOOKS III. and IV. *By G. A. SIMON, M.A.* 6s.

---

*WILKINSON, PEARCE, LONDON.*

## MURRAY'S DIVINITY

Crown 8vo. 2s. 6d.

*The Gospel according to S. Matthew.* With Explanatory Notes for the Use of Teachers. By HENRY HERBERT WYATT, M.A., Principal of Brighton Training College, and Vicar of Bolney, Sussex. With Commendatory Preface by the Archbishops' Inspector of Training Colleges.

New and Revised Editions. Small 8vo. 3s. 6d. each. Sold separately.

*Manuals of Religious Instruction.* Edited by

JOHN PILKINGTON NORRIS, D.D., Archdeacon of Bristol.

The Old Testament. | The New Testament.  
The Prayer Book.

Cheap Edition. Small 8vo. 1s. 6d. each.

*Keys to Christian Knowledge.* By the Rev. J. H.

BLUNT, D.D., Editor of the "Annotated Book of Common Prayer."

The Holy Bible.	The Church Catechism.
The Book of Common Prayer.	Church History, Ancient.
	Church History, Modern.

By JOHN PILKINGTON NORRIS, D.D., Archdeacon of Bristol.

The Four Gospels. | The Acts of the Apostles.

18mo. 1s. 6d.

*Easy Lessons Addressed to Candidates for Confirmation.* By J. P. NORRIS, D.D., Archdeacon of Bristol.

New Edition. Small 8vo. 1s. 6d.

*A Manual of Confirmation.* By EDWARD MEYRICK GOULBURN, D.D., Dean of Norwich.

Crown 8vo. 7s. 6d.

*Some Helps for School Life.* Sermons preached at Clifton College, 1862-1879. By J. PERCIVAL, M.A., LL.D., President of Trinity College, Oxford, and late Head Master of Clifton College.

Crown 8vo. 1s. Cloth limp, 1s. 6d.

*Study of the Church Catechism.* Adapted for use as a Class Book. By C. J. SHERWILL DAWE, M.A., Lecturer and Assistant Chaplain at St. Mark's College, Chelsea.

Waterloo Place, London.



*New Edition. Small 8vo. 3s. 6d.*

**Household Theology.** A Handbook of Religious Information respecting the Holy Bible, the Prayer Book, the Church, the Ministry, Divine Worship, the Creeds, &c. &c. *By the Rev. JOHN HENRY BLUNT, D.D., F.S.A.*

*Second Edition, Revised. Crown 8vo. 7s. 6d.*

**Rudiments of Theology.** A First Book for Students. *By JOHN PILKINGTON NORRIS, D.D., Archdeacon of Bristol.*

*16mo, 1s. 6d.; Paper Covers, 1s.; or in Three Parts, 6d. each.*

**The Young Churchman's Companion to the Prayer Book.** *Edited by the Rev. J. W. GEDGE, M.A., Winchester, Diocesan Inspector of Schools for West Surrey and the Channel Islands.*

PART I.—MORNING AND EVENING PRAYER, AND LITANY.

PART II.—BAPTISMAL AND CONFIRMATION SERVICES.

PART III.—THE HOLY COMMUNION.

*Second Edition. Large type. 24mo. 1s.*

**Prayers for a Young Schoolboy.** *By the Rev. E. B. PUSEY, D.D. Edited, with a Preface, by H. P. LIDDON, D.D., Canon Residentiary of St. Paul's.*

*Second Edition. 18mo. 1s. 6d.*

**The Way of Life.** A Book of Prayers and Instruction for the Young at School. With a Preparation for Holy Communion. *Compiled by a Priest. Edited by the Rev. T. T. CARTER, M.A.*

*16mo. 2s. 6d.*

**A Plain Exposition of the Thirty-nine Articles of the Church of England.** For the use of Schools. *By the Rev. WILLIAM BAKER, D.D., Head Master of Merchant Taylors' School, and Prebendary of St. Paul's.*

*Crown 16mo. Cloth limp. 1s. 6d.*

**A Manual of Devotion, chiefly for the Use of Schoolboys.** *By WILLIAM BAKER, D.D., Head Master of Merchant Taylors' School. With Preface by J. R. WOODFORD, D.D., Lord Bishop of Ely.*

Waterloo Place, London.

*Small 8vo. 1s. 6d. each.*

## **Scenes from Greek Plays.** RUGBY EDITION.

*Abridged and adapted for the use of Schools, by ARTHUR SIDGWICK, M.A., Fellow of Corpus Christi College, Oxford, late Assistant Master at Rugby School, and Fellow of Trinity College, Cambridge.*

### **Aristophanes.**

THE CLOUDS. THE FROGS. THE KNIGHTS. PLUTUS.

### **Euripides.**

IPHIGENIA IN TAURIS. THE CYCLOPS. ION.  
ELECTRA. ALCESTIS. BACCHÆ. HECUBA.

*Fourth Edition. Crown 8vo. 3s. 6d.*

**Stories in Attic Greek.** Forming a Greek Reading Book for the use of Junior Forms in Schools. With Notes and Vocabulary. By FRANCIS DAVID MORICE, M.A., Assistant Master at Rugby School, and Fellow of Queen's College, Oxford.

*New Edition. Crown 8vo.*

**The Anabasis of Xenophon.** Edited, with Preface, Introduction, Historical Sketch, Itinerary, Syntax Rules, Notes, Indices, Vocabularies, and Maps, by R. W. TAYLOR, M.A., Head Master of Kelly College, Tavistock, and late Fellow of St. John's College, Cambridge.

BOOKS I. and II. 3s. 6d. BOOKS III. and IV. 3s. 6d.

Also separately, BOOK I., 2s. 6d.; BOOK II., 2s.

*Crown 8vo. 2s. 6d.*

**Xenophon's Agesilaus.** Edited, with Syntax Rules, and References, Notes, and Indices, by R. W. TAYLOR, M.A.

*Second Edition. Small 8vo. 2s.*

**Xenophon's Memorabilia.** BOOK I., with a few omissions. Edited, with an Introduction and Notes, by C. E. MOBERLY, M.A., formerly Scholar of Balliol College, Oxford.

*New Edition. Small 8vo. 2s.*

**Alexander the Great in the Punjaub.** Adapted from Arrian, Book V. An Easy Greek Reading Book. Edited, with Notes and a Map, by C. E. MOBERLY, M.A., formerly Scholar of Balliol College, Oxford.

Waterloo Place, London.

*Small 8vo.*

**Homer's Iliad.** Edited, with Notes at the end for the Use of Junior Students, by ARTHUR SIDGWICK, M.A., *Fellow of Corpus Christi College, Oxford; late Assistant Master at Rugby School, and Fellow of Trinity College, Cambridge.*

BOOKS I. and II. 2s. 6d.

BOOK XXI. 1s. 6d.

BOOK XXII. 1s. 6d.

*Small 8vo. 2s.*

**Homer without a Lexicon, for Beginners.** ILIAD, BOOK VI. Edited, with Notes giving the meanings of all the less common words, by J. SURTEES PHILLPOTTS, M.A., *Head Master of Bedford Grammar School.*

*Fifth Edition. 12mo. 3s. 6d.*

**Homer for Beginners.** ILIAD, BOOKS I.-III. With English Notes. By THOMAS KERCHEVER ARNOLD, M.A.

*Fifth Edition. 12mo. 12s.*

**The Iliad of Homer.** With English Notes and Grammatical References. By THOMAS KERCHEVER ARNOLD, M.A.

*Crown 8vo. 6s.*

**The Iliad of Homer.** BOOKS I.-XII. From the Text of Dindorf. With Preface and Notes. By S. H. REYNOLDS, M.A., *late Fellow and Tutor of Brasenose College, Oxford.*

*8vo. 18s.*

**Myths of the Odyssey in Art and Literature.** Illustrated with Outline Drawings. By J. E. HARRISON.

*New Edition. 12mo. 9s.*

**A Complete Greek and English Lexicon for the Poems of Homer and the Homeridæ.** By G. CH. CRUSIUS. Translated from the German. Edited by T. K. ARNOLD, M.A.

*Crown 8vo. 4s. 6d.*

**Isocratis Orationes.** AD DEMONICUM ET PANEGYRICUS. Edited by JOHN EDWIN SANDYS, M.A., *Fellow and Tutor of St. John's College, Cambridge, and Public Orator of the University.*



8vo. 16s.

**Hellenica.** *A Collection of Essays on Greek Poetry, Philosophy, History, and Religion.* Edited by EVELYNABBOTT, M.A., LL.D., *Fellow and Tutor of Balliol College, Oxford.*

CONTENTS.—Aeschylus. E. Myers, M.A.—The Theology and Ethics of Sophocles. E. Abbott, M.A., LL.D.—System of Education in Plato's Republic. R. L. Nettleship, M.A.—Aristotle's Conception of the State. A. C. Bradley, M.A.—Epicurus. W. L. Courtney, M.A.—The Speeches of Thucydides. R. C. Jebb, M.A., LL.D.—Xenophon. H. G. Dakyns, M.A.—Polybius. J. L. S. Davidson, M.A.—Greek Oracles. F. W. H. Myers, M.A.

8vo. 18s.

**The Antiquities of Greece.** THE STATE. Translated from the German of G. F. SCHOEMANN. By E. G. HARDY, M.A., *Head Master of the Grammar School, Grantham*; and J. S. MANN, M.A., *Fellow of Trinity College, Oxford.*

Crown 8vo.

**Herodoti Historia.** Edited by H. G. WOODS, M.A., *Fellow of Trinity College, Oxford.*

BOOK I. 6s. BOOK II. 5s.

12mo.

**Demosthenes.** Edited, with English Note, and Grammatical References, by THOMAS KERCHEVER ARNOLD, M.A.

OLYNTIAC ORATIONS. Third Edition. 3s.

ORATION ON THE CROWN. Second Edition. 4s. 6d.

Crown 8vo. 5s.

**Demosthenis Orationes Privatae.** DE CORONA.Edited by ARTHUR HOLMES, M.A., *late Senior Fellow and Dean of Clare College, Cambridge.*

Crown 8vo.

**Demosthenis Orationes Publicae.** Edited by G. II.HESLOP, M.A., *late Fellow and Assistant Tutor of Queen's College, Oxford*; *Head Master of St. Bees.*

OLYNTIACS, 2s. 6d. } or, in One Volume, 4s. 6d.

PHILIPPICS, 3s.

DE FALSA LEGATIONE, 6s.

Crown 8vo.

**Aristophanis Comædiæ.** Edited by W. C. GREEN, M.A., *late Fellow of King's College, Cambridge*; *Assistant Master at Rugby School.*

THE ACHARNIANS and THE KNIGHTS. 4s.

THE CLOUDS. 3s. 6d.

THE WASPS. 3s. 6d.

Waterloo Place, London.

*Second Edition, Revised and Enlarged. Crown 8vo. 10s. 6d.*

**An Introduction to Aristotle's Ethics.** BOOKS I.-IV.  
(BOOK X., c. vi.-ix. in an Appendix). With a Continuous Analysis  
and Notes. Intended for the use of Beginners and Junior Students.  
*By the Rev. EDWARD MOORE, D.D., Principal of St. Edmund Hall,  
and late Fellow and Tutor of Queen's College, Oxford.*

*Second Edition. Crown 8vo. 3s. 6d.*

**Selections from Aristotle's Organon.** Edited by JOHN  
R. MAGRATH, D.D., Provost of Queen's College, Oxford.

12mo.

**Sophocles.** Edited by T. K. ARNOLD, M.A., ARCHDEACON PAUL,  
and HENRY BROWN, M.A.

AJAX. 3s.

OEDIPUS TYRANNUS. 4s.

*Crown 8vo.*

**Sophoclis Tragicæ.** Edited by R. C. JEBB, M.A., LL.D.,  
Professor of Greek at the University of Glasgow, late Fellow and  
Tutor of Trinity College, Cambridge.

ELECTRA. 3s. 6d.

AJAX. 3s. 6d.

*Crown 8vo. 8s. 6d.*

**Sophocles.** Translated into English Verse. By ROBERT WHITELAW,  
M.A., Assistant Master in Rugby School; late Fellow of Trinity  
College, Cambridge.

*Crown 8vo. 6s.*

**Thucydidis Historia.** BOOKS I. and II. Edited by CHARLES  
BIGG, D.D., late Senior Student and Tutor of Christ Church,  
Oxford; formerly Principal of Brighton College.

*Crown 8vo. 6s.*

**Thucydidis Historia.** BOOKS III. and IV. Edited by G. A.  
SIMCOX, M.A., Fellow of Queen's College, Oxford.

*Sixth Edition. 8vo. 21s.*

**A Copious Phraseological English-Greek Lexicon.**  
Founded on a work prepared by J. W. FRÄDERSDORFF, Ph.D., late  
Professor of Modern Languages, Queen's College, Belfast. Revised,  
Enlarged, and Improved by THOMAS KERCHEVER ARNOLD, M.A.,  
and HENRY BROWNE, M.A.

---

Waterloo Place, London.

*Third Edition. Crown 8vo. 2s. 6d.*

**Short Notes on the Greek Text of the Gospel of St. Mark.** *By J. HAMBLIN SMITH, M.A., of Gonville and Caius College, Cambridge.*

*Crown 8vo. 4s. 6d.*

**Notes on the Greek Text of the Acts of the Apostles.** *By J. HAMBLIN SMITH, M.A., of Gonville and Caius College, Cambridge.*

*Crown 8vo. 6s.*

**Notes on the Gospel According to S. Luke.** *By the Rev. ARTHUR CARR, M.A., Assistant Master at Wellington College, late Fellow of Oriel College, Oxford.*

*New Edition. 4 vols. 8vo. 102s.*

**The Greek Testament.** With a Critically Revised Text; a Digest of Various Readings; Marginal References to Verbal and Idiomatic Usage; Prolegomena; and a Critical and Exegetical Commentary. For the use of Theological Students and Ministers. *By HENRY ALFORD, D.D., late Dean of Canterbury.*

The Volumes are sold separately, as follows :—

Vol. I.—THE FOUR GOSPELS. 28s.

Vol. II.—ACTS TO 2 CORINTHIANS. 24s.

Vol. III.—GALATIANS TO PHILEMON. 18s.

Vol. IV.—HEBREWS TO REVELATION. 32s.

*New Edition. 2 vols. Imperial 8vo. 60s.*

**The Greek Testament.** With Notes, Introductions, and Index. *By CHR. WORDSWORTH, D.D., Bishop of Lincoln.*

The Parts may be had separately, as follows :

THE GOSPELS. 16s.

THE ACTS. 8s.

ST. PAUL'S EPISTLES. 23s.

GENERAL EPISTLES, REVELATION, AND INDEX. 6s.

Waterloo Place, London.



## CATENA CLASSICORUM

*Crown 8vo.***Aristophanis Comœdiæ.** *By* W. C. GREEN, M.A.

THE ACHARNIANS AND THE KNIGHTS. 4s.

THE WASPS. 3s. 6d. THE CLOUDS. 3s. 6d.

**Demosthenis Orationes Publicæ.** *By* G. H. HESLOP, M.A.THE OLYNTHIACS. 2s. 6d. } or, in One Volume, 4s. 6d.  
THE PHILIPPICS. 3s.

DE FALSA LEGATIONE. 6s.

**Demosthenis Orationes Privatæ.** *By* A. HOLMES, M.A.

DE CORONA. 5s.

**Herodoti Historia.** *By* H. G. WOODS, M.A.

BOOK I., 6s. BOOK II., 5s.

**Homeri Ilias.** *By* S. H. REYNOLDS, M.A.

BOOKS I.-XII. 6s.

**Horati Opera.** *By* J. M. MARSHALL, M.A.

THE ODES, CARMEN SECULARE, AND EPODES. 7s. 6d.

THE ODES. BOOKS I. to IV. separately. 1s. 6d. each.

**Isocratis Orationes.** *By* JOHN EDWIN SANDYS, M.A.

AD DEMONICUM ET PANEGYRICUS. 4s. 6d.

**Juvenalis Satiræ.** *By* G. A. SIMCOX, M.A. 5s.**Persii Satiræ.** *By* A. PRETOR, M.A. 3s. 6d.**Sophoclis Tragoediæ.** *By* R. C. JEBB, M.A.

THE ELECTRA. 3s. 6d. THE AJAX. 3s. 6d.

**Taciti Historiæ.** *By* W. H. SIMCOX, M.A.

BOOKS I. and II., 6s. BOOKS III. IV. and V., 6s.

**Terenti Comœdiæ.** *By* T. L. PAPILLON, M.A.ANDRIA AND EUNUCHUS. With Introduction on  
Prosody. 4s. 6d. *Or separately.*

ANDRIA. With Introduction on Prosody. 3s. 6d.

EUNUCHUS. 3s.

**Thucydidis Historia.**BOOKS I. and II. *By* CHARLES BIGG, D.D. 6s.BOOKS III. and IV. *By* G. A. SIMCOX, M.A. 6s.

Waterloo Place, London.

## DIVINITY

*Crown 8vo. 2s. 6d.*

***The Gospel according to S. Matthew.*** With Explanatory Notes for the Use of Teachers. By HENRY HERBERT WYATT, M.A., Principal of Brighton Training College, and Vicar of Bolney, Sussex. With Commendatory Preface by the Archbishops' Inspector of Training Colleges.

*New and Revised Editions. Small 8vo. 3s. 6d. each. Sold separately.*

***Manuals of Religious Instruction.*** Edited by

JOHN PILKINGTON NORRIS, D.D., Archdeacon of Bristol.

The Old Testament. | The New Testament.  
The Prayer Book.

*Cheap Edition. Small 8vo. 1s. 6d. each.*

***Keys to Christian Knowledge.*** By the Rev. J. H.

BLUNT, D.D., Editor of the "Annotated Book of Common Prayer."

The Holy Bible. | The Church Catechism.

The Book of Common Prayer. | Church History, Ancient.  
Church History, Modern.

By JOHN PILKINGTON NORRIS, D.D., Archdeacon of Bristol.

The Four Gospels. | The Acts of the Apostles.

18mo. 1s. 6d.

***Easy Lessons Addressed to Candidates for Confirmation.*** By J. P. NORRIS, D.D., Archdeacon of Bristol.

*New Edition. Small 8vo. 1s. 6d.*

***A Manual of Confirmation.*** By EDWARD MEYRICK GOULBURN, D.D., Dean of Norwich.

*Crown 8vo. 7s. 6d.*

***Some Helps for School Life.*** Sermons preached at Clifton College, 1862-1879. By J. PERCIVAL, M.A., LL.D., President of Trinity College, Oxford, and late Head Master of Clifton College.

*Crown 8vo. 1s. Cloth limp, 1s. 6d.*

***Study of the Church Catechism.*** Adapted for use as a Class Book. By C. J. SHERWILL DAWE, M.A., Lecturer and Assistant Chaplain at St. Mark's College, Chelsea.

Waterloo Place, London.

*New Edition. Small 8vo. 3s. 6d.*

**Household Theology.** A Handbook of Religious Information respecting the Holy Bible, the Prayer Book, the Church, the Ministry, Divine Worship, the Creeds, &c. &c. *By the Rev. JOHN HENRY BLUNT, D.D., F.S.A.*

*Second Edition, Revised. Crown 8vo. 7s. 6d.*

**Rudiments of Theology.** A First Book for Students. *By JOHN PILKINGTON NORRIS, D.D., Archdeacon of Bristol.*

*16mo, 1s. 6d.; Paper Covers, 1s.; or in Three Parts, 6d. each.*

**The Young Churchman's Companion to the Prayer Book.** *Edited by the Rev. J. W. GEDGE, M.A., Winchester, Diocesan Inspector of Schools for West Surrey and the Channel Islands.*

PART I.—MORNING AND EVENING PRAYER, AND LITANY.

PART II.—BAPTISMAL AND CONFIRMATION SERVICES.

PART III.—THE HOLY COMMUNION.

*Second Edition. Large type. 24mo. 1s.*

**Prayers for a Young Schoolboy.** *By the Rev. E. B. PUSEY, D.D. Edited, with a Preface, by H. P. LIDDON, D.D., Canon Residentiary of St. Paul's.*

*Second Edition. 18mo. 1s. 6d.*

**The Way of Life.** A Book of Prayers and Instruction for the Young at School. With a Preparation for Holy Communion. *Compiled by a Priest. Edited by the Rev. T. T. CARTER, M.A.*

*16mo. 2s. 6d.*

**A Plain Exposition of the Thirty-nine Articles of the Church of England.** For the use of Schools. *By the Rev. WILLIAM BAKER, D.D., Head Master of Merchant Taylors' School, and Prebendary of St. Paul's.*

*Crown 16mo. Cloth limp. 1s. 6d.*

**A Manual of Devotion, chiefly for the Use of Schoolboys.** *By WILLIAM BAKER, D.D., Head Master of Merchant Taylors' School. With Preface by J. R. WOODFORD, D.D., Lord Bishop of Ely.*

---

Waterloo Place, London.



## GERMAN

*New Edition, Revised. 4to. 3s. d.*

### ***A German Accidence for the Use of Schools.***

*By J. W. J. VECQUERAY, Assistant Master at Rugby School.*

*Crown 8vo. 2s.*

### ***First German Exercises.*** Adapted to Vecqueray's "German Accidence for the Use of Schools." *By E. F. GRENFELL, M.A., late Assistant Master at Rugby School.*

*Crown 8vo. 2s. 6d.*

### ***German Exercises. Part II.*** With Hints for the Translation of English Prepositions into German. Adapted to Vecqueray's "German Accidence for the Use of Schools." *By E. F. GRENFELL, M.A., late Assistant Master at Rugby School.*

*New Edition. Crown 8vo. 4s. 6d.*

### ***Selections from Hauff's Stories.*** A First German Reading Book. *Edited by W. E. MULLINS, M.A., Assistant Master at Marlborough College, and F. STORR, B.A., Chief Master of Modern Subjects in Merchant Taylors' School.*

*Also, separately, crown 8vo. 2s.*

### ***Kalif Stork and The Phantom Crew.***

*Eighth Edition. 12mo. 5s. 6d.*

### ***The First German Book.*** *By T. K. ARNOLD, M.A., and J. W. FRÄBERSDORFF, Ph.D. KEY, 2s. 6d.*

*Second Edition. Crown 8vo. 2s. 6d.*

### ***Lessing's Fables.*** Arranged in order of difficulty. A First German Reading Book. *By F. STORR, B.A., Chief Master of Modern Subjects in Merchant Taylors' School, and late Assistant Master in Marlborough College.*

*Crown 8vo. 7s. 6d.*

### ***Goethe's Faust.*** PART I. Text, with English Notes, Essays, Verse Translations. *By E. J. TURNER, M.A., and E. D. A. MORSEHEAD, M.A., Assistant Masters at Winchester College.*

---

Waterloo Place, London.

## FRENCH

*Crown 8vo. 1s. 6d.*

**Exercises in French Syntax.** With Rules. By G. SHARP, M.A., Assistant Master at Marlborough College.

*New Edition, Revised. Crown 8vo. 2s. 6d.*

**French Passages for Unseen Translation.** Selected and arranged by C. H. PARRY, M.A., Assistant Master at Charterhouse.

*New Edition. Small 8vo. 2s.*

**A Graduated French Reader.** With an Introduction on the Pronunciation of Consonants and the Connection of Final Letters, a Vocabulary, and Notes, and a Table of Irregular Verbs with the Latin Infinitives. By PAUL BARBIER, Professor at University College, Cardiff.

*Crown 8vo.*

**The Campaigns of Napoleon.** The Text (in French) from M. THIERS' "Histoire de la Révolution Française," and "Histoire du Consulat et de l'Empire." Edited, with English Notes and Maps, for the use of Schools, by EDWARD E. BOWEN, M.A., Master of the Modern Side, Harrow School.

ARCOLA. 4s. 6d.

MARENGO. 4s. 6d.

JENA. 3s. 6d.

WATERLOO. 6s.

*New Editions. Crown 8vo. 3s. 6d. each.*

**Selections from Modern French Authors.** Edited, with English Notes and Introductory Notice, by HENRI VAN LAUN, Translator of TAINÉ's "History of English Literature."

HONORÉ DE BALZAC.

H. A. TAINÉ.

*Small 8vo. 2s.*

**La Fontaine's Fables.** BOOKS I. and II. Edited, with English Notes at the end, by Rev. P. BOWDEN-SMITH, M.A., Assistant Master at Rugby School.

*Sixth Edition. 12mo. 5s. 6d.*

**The First French Book.** By T. K. ARNOLD, M.A.

KEY, 2s. 6d.

---

Waterloo Place, London.

## MISCELLANEOUS

*With Maps. Small 8vo.*

### *A Geography, Physical, Political, and Descriptive.*

For Beginners. *By* L. B. LANG. *Edited by the Rev. M. CREIGHTON, M.A., LL.D., late Fellow and Tutor of Merton College, Oxford.*

VOL. I. THE BRITISH EMPIRE. 2s. 6d.

PART I. THE BRITISH ISLES. 1s. 6d. PART II. THE BRITISH POSSESSIONS. 1s. 6d.

VOL. II. THE CONTINENT OF EUROPE. 3s.

VOL. III. ASIA, AFRICA, AND AMERICA. 2s.

*Small 8vo. 2s. 6d. each part.*

### *Modern Geography for the Use of Schools.*

*By* C. E. MOBERLY, M.A., *formerly Scholar of Balliol College, Oxford.*

PART I. NORTHERN EUROPE.

PART II. THE MEDITERRANEAN & ITS PENINSULAS.

*Crown 8vo. 3s. 6d.*

### *At Home and Abroad; or, First Lessons in Geography.* *By* J. K. LAUGHTON, M.A., F.R.A.S., F.R.G.S., *Mathematical Instructor and Lecturer at the Royal Naval College.*

*Second Edition. Crown 8vo. 2s. 6d.*

### *The Chorister's Guide.* *By* W. A. BARRETT, Mus. Bac. Oxon., *Vicar Choral of St. Paul's Cathedral, Author of "Flowers and Festivals," &c.*

*Crown 8vo. 2s. 6d.*

### *An Introduction to Form and Instrumentation.*

For the use of Beginners in Composition. *By* W. A. BARRETT, Mus. Bac. Oxon., *Vicar Choral of St. Paul's Cathedral.*

*Seventh Edition, Revised. 12mo. 7s. 6d.*

### *The First Hebrew Book.* *By* T. K. ARNOLD, M.A.

KEY, 3s. 6d.

---

Waterloo Place, London.



*By J. Hamblin Smith.*

**Elementary Algebra.** *Small 8vo, 3s. Without Answers, 2s. 6d.*

**Key to Elementary Algebra.** *Crown 8vo, 9s.*

**Exercises on Algebra.** *Small 8vo, 2s. 6d.*

**Arithmetic.** *Small 8vo, 3s. 6d.*

**Key to Arithmetic.** *Crown 8vo, 9s.*

**Elements of Geometry.** *Small 8vo, 3s. 6d.*

Books I. and II., limp cloth, price 1s. 6d., may be had separately.

**Key to Elements of Geometry.** *Crown 8vo, 8s. 6d.*

**Trigonometry.** *Small 8vo, 4s. 6d.*

**Key to Trigonometry.** *Crown 8vo, 7s. 6d.*

**Elementary Statics.** *Small 8vo, 3s.*

**Elementary Hydrostatics.** *Small 8vo, 3s.*

**Key to Elementary Statics and Hydrostatics.** *Crown 8vo, 6s.*

**Book of Enunciations for HAMBLIN SMITH'S Geometry, Algebra, Trigonometry, Statics, and Hydrostatics.** *Small 8vo, 1s.*

**An Introduction to the Study of Heat.** *Small 8vo, 3s.*

**Latin Grammar.** *Crown 8vo, 3s. 6d.*

**Exercises on the Elementary Principles of Latin Prose Composition.** *Crown 8vo, 3s. 6d.*

**Key to Exercises on Latin Prose Composition.** *Crown 8vo, 5s.*

**An Elementary Greek Grammar.** *Crown 8vo, 4s. 6d.*

**The Rudiments of English Grammar and Composition.** *Crown 8vo, 2s. 6d.*

**Notes on the Greek Text of the Acts of the Apostles.** *Crown 8vo, 4s. 6d.*

**Notes on the Greek Text of the Gospel of St. Mark.** *Crown 8vo, 2s. 6d.*

---

Waterloo Place, London,

*By Arthur Sidgwick.*

An Introduction to Greek Prose Composition.

*Crown 8vo, 5s. A KEY, 5s.*

Lectures on Greek Prose.

An Introduction to Greek Verse Composition.

*Crown 8vo, 5s. A KEY, 5s.*

A First Greek Writer. *Crown 8vo, 3s. 6d. A KEY, 5s.*

Cicero de Amicitia. *Small 8vo, 2s.*

Homer's Iliad. *Small 8vo. BOOKS I. and II., 2s. 6d. BOOK XXI., 1s. 6d. BOOK XXII., 1s. 6d.*

Scenes from Greek Plays. *Small 8vo, each 1s. 6d.*

ARISTOPHANES: The Clouds, The Frogs, The Knights, Plutus.

EURIPIDES: Iphigenia in Tauris, The Cyclops, Ion, Electra, Alcestis, Bacchæ, Hecuba.

*By George L. Bennett.*

First Latin Writer. *Crown 8vo, 3s. 6d. A KEY, 5s.*

First Latin Exercises. *Crown 8vo, 2s. 6d.*

First Latin Accidence. *Crown 8vo, 1s. 6d.*

Second Latin Writer. *Crown 8vo, 3s. 6d. A KEY, 5s.*

Easy Latin Stories for Beginners. *Cr. 8vo, 2s. 6d. A KEY, 5s.*

Viri Illustres Urbis Romæ. *Small 8vo.*

Second Latin Reading Book. *Cr. 8vo, 2s. 6d. A KEY, 5s.*

Selections from Caesar. The Gallic War. *Small 8vo, 2s.*

Selections from Vergil. *Small 8vo, 1s. 6d.*

Easy Unseen Latin Passages. *16mo, 1s. A KEY, 3s. 6d.*

*By R. W. Taylor.*

Xenophon's Anabasis. *Crown 8vo. Books I. and II., 3s. 6d. III. and IV., 3s. 6d. Also separately, Book I., 2s. 6d. II., 2s.*

Xenophon's Agesilaus. *Crown 8vo, 2s. 6d.*

A Short Greek Syntax. *Crown 8vo, 9d.*

Stories from Ovid in Elegiac Verse. *Crown 8vo, 3s. 6d.*

Stories from Ovid in Hexameter Verse. METAMORPHOSES. *Crown 8vo, 2s. 6d.*

Scott's Lady of the Lake. Forming a Volume of the "English School Classics." *Small 8vo, 2s.; or in Three Parts, each 9d.*

Waterloo Place, London.


# INDEX

	PAGE		PAGE
ABBOTT (E.), Arnold's Greek Prose	24	Cornish (F. W.), Oliver Cromwell	10
— Elements of Greek Accidence	24	Crake (A. D.), History of the Church	9
— Greek Outline	10	Creighton (L.), First Hist. of Eng.	8
— Hellenica	28	— Highways of History	5
— Lucian, Selections from	25	— Historical Biographies	10
— and Mansfield (E. D.), Primer of Greek Grammar	22	— Stories from English History	8
Acland (A.), Political Hist. of Eng.	10	Crofts (E.), English Literature	9
— Skeleton Outline	10	Crusius (G. C.), Homeric Lexicon	27
Ainger (A. C.), Clivus and Key	17	Curteis (A. M.), The Roman Empire	9
Alford (Dean), Greek Testament	30	DALLIN (T.), Materials and Models	20, 25
Aristophanes	26, 28	Davys (Bishop), History of England	8
Aristotle	29	Dawe (C. J. S.), Latin Exercise Bk.	16
Arnold (T. K.), Crusius' Homeric Lexicon	27	— Church Catechism	32
— Demosthenes	28	Demosthenes	28
— Eclogæ Ovidianæ	19	Donkin (A. E.), Arithmetic	13
— Eng.-Greek Lexicon	29	ENGLISH School Classics	6, 7
— First French Book and Key	35	Euripides, Scenes from	26
— First German Book and Key	34	FRADERSDORFF, Eng.-Greek Lexicon	29
— First Greek Book and Key	23	GANTILLON (P. J. F.), Exam. Papers	20, 25
— — revised by F. D. Morice	23	Gedge (J. W.), Com. to Prayer Book	33
— First Hebrew Book and Key	36	Gepp (C. G.), Arnold's Henry's First Latin Book and Key	18
— First Verse Book and Key	17	— Latin Elegiac Verse and Key	17
— Greek Accidence	24	— Latin-English Dictionary	20
— Greek Prose Comp. and Key	24	— Virgil	18
— — revised by E. Abbott	24	Girdlestone (W. H.), Arithmetic	13
— Henry's First Latin and Key	18	Goethe's Faust	34
— — revised by C. G. Gepp	18	Goolden (W. T.), Intro. to Chemistry	11
— Homer's Iliad	27	Goulburn (Dean), Confirmation	32
— Latin Prose Comp. and Key	18	Greek Plays, Scenes from	26
— — revised by G. G. Bradley	18	Green (A. H.), Geology	11
— Madvig's Greek Syntax	24	— (W. C.), Aristophanes	28
— Sophocles	29	Grenfell (E. F.), German Exercises	34
Arrian	26	Gross (E. J.), Algebra, Part II.	18
BAKER (W.), Manual of Devotion	33	— Kinematics and Kinetics	12
— Thirty-nine Articles	33	HARDY (E. G.), Antiq. of Greece	28
Barbier (P.), French Reader	35	Harrison (J. E.), Myths of the Odyssey	27
Barrett (W. A.), Chorister's Guide	36	Hauff's Stories, Selections from	34
— Form and Instrumentation	36	Heatley (H. R.), Excerpta Facilia and Key	16
Belcher (H.), Livy, Book II.	19	— Græcula and Key	22
Bennett (G. L.), Caesar's Gallic War	15	— Gradatim and Key	16
— Easy Latin Stories and Key	15	— Grammar Papers	16, 22
— Second Latin Reader and Key	15	Hellenica, Essays	28
— First Latin Writer and Key	15	Herodotus, Stories from, Phillpotts	25
— First Latin Exercises	15	— By H. G. Woods	28
— Latin Accidence	15	Hertz (H. A.), English Poetry	7
— Second Latin Writer and Key	15	Heslop (G. H.), Demosthenes	28
— Unseen Latin Passages and Key	16	Highways of History	5
— Vergil, Selections from	15	Historical Biographies	10
— Viri Illustres	15	Historical Handbooks	9
— Works by	38	Hodges (C. H.), Arithmetic	13
Bigg (C.), Thucydides, Books I. II.	29	Holmes (A.), Demosthenes	28
Blunt (J. H.), Household Theology	33	— Rules of Latin Pronunciation	17
— Keys to Christian Knowledge	32	Homer's Iliad	27
Bowen (E. E.), Campaigns of Napoleon	35	Horace, By J. M. Marshall	21
Bradley (G. G.), Aids to Latin Prose	18	Horton (R. F.), Roman History	9
— Arnold's Latin Prose	18	IOPHON	25
Bridge (C.), French Literature	9	Isocrates, By J. E. Sandys	27
Bright (J. F.), History of England	8	JEBB (R. C.), Sophocles	28
Building Construction, Notes on	11	Jennings (A. C.), Ecclesia Anglicana	9
Burton (J.), English Grammar	7	Juvenal, By G. A. Simcox	21
CÆSAR	15, 17, 19	KEYS TO CHRISTIAN KNOWLEDGE	32
Carr (A.), Notes on St. Luke	30	— List of	4
Catena Classicorum	31		
Cicero de Amicitia	19		
Clarke (A. D.), Examination Papers	13		



# INDEX.

	PAGE		PAGE
Kitchener (F. A.), A Year's Botany .	11	SANDYS (J. E.), Isocratis Orationes .	27
Kingdon, <i>Excerpta Facilia</i> and Key .	16	Sargent (J.), Latin Passages . . .	20
— Gradatim and Key . . . . .	16	— Materials and Models . . . . .	20, 25
LATIN TEXT BOOKS . . . . .	17	Schoemann's Antiquities of Greece .	28
La Fontaine's Fables. By P. Smith .	35	Shakspeare's Plays . . . . .	5
Lang (L. B.), Geography for Beginners .	36	Sharp (G.), French Syntax . . . . .	35
Laughton (J.), At Home and Abroad .	36	Sidgwick (A.), Cicero de Amicitia . .	19
Laun Van (H.), French Selections . .	35	— First Greek Writer and Key . . .	23
Lessing's Fables. By F. Storr . . . .	34	— Greek Prose Composition and Key .	23
Livy . . . . .	19	— Greek Verse and Key . . . . .	23
Locke (C. L. C.), English Parsing . .	7	— Homer's Iliad . . . . .	27
Lucian, Selections from . . . . .	25	— Scenes from Greek Plays . . . .	26
MADVIG'S Greek Syntax . . . . .	24	— Works by . . . . .	38
Magrath (J. R.), Aristotle's Organon .	29	Simcox (G. A.), Juvenalis Satire . .	21
Mann (J. S.), Antiquities of Greece .	28	— Thucydides . . . . .	29
Mansfield (E. D.), Latin Sentence . .	30	— (W. H.), Taciti Historia . . . .	21
— Primer of Greek Syntax . . . . .	22	Smith (J. Hamblin), The Acts . . . .	30
Manuals of Religious Instruction . .	32	— Algebra and Key . . . . .	12
Marshall (J. M.), Horati Opera . . .	21	— Algebra, Exercises on . . . . .	12
— (L.), Companion to Algebra . . .	13	— Arithmetic and Key . . . . .	12
Matheson (P. E.), Roman Outline . .	10	— Book of Enunciations . . . . .	13
Merryweather (J. H.), Caesar . . . .	19	— English Grammar . . . . .	7
Moberly (C. E.), Alexander the Great .	26	— Geometry and Key . . . . .	13
— Geography . . . . .	36	— Greek Grammar . . . . .	24
— Shakspeare's Plays . . . . .	5	— Heat, The Study of . . . . .	13
— Xenophon's Memorabilia . . . .	26	— Hydrostatics and Key . . . . .	12
Moore (E.), Aristotle's Ethics . . . .	29	— Latin Grammar . . . . .	17
Moore (E. H.), Greek Method . . . .	22	— Prose Composition and Key . . .	19
— Greek Verbs . . . . .	22	— Statics and Key . . . . .	12
Morice (F. D.), Arnold's First Greek Bk.	23	— St. Mark's Gospel . . . . .	30
— Greek Verse and Key . . . . .	23	— Trigonometry and Key . . . . .	12
— Stories in Attic Greek . . . . .	26	— Works by . . . . .	37
Morshead (E. D.), Goethe's Faust . .	34	— (P. Bowden), La Fontaine's Fables .	35
Mullins (W. E.), Hauff's Stories . . .	34	— (P. V.), English Institutions . .	9
NAPOLEON'S Campaigns . . . . .	35	— (R. Prowde), Latin Prose Ex. . .	19
Norris (J. P.), Confirmation . . . . .	32	Sophocles . . . . .	29
— Keys to Christian Knowledge . . .	32	Spratt (A. W.), Translation at Sight .	20, 25
— Manuals of Religious Instruction .	32	Storr (F.), <i>Æneid</i> of Vergi . . . .	24
— Rudiments of Theology . . . . .	33	— Greek Verbs . . . . .	24
OVIDIANÆ ECLOGÆ. By Arnold . . . .	19	— Hauff's Stories . . . . .	34
Ovid, Stories from. By R. W. Taylor .	19	— Lessing's Fables . . . . .	34
PAPILLON (T. L.), Terenti Comedie .	21	TACITUS. By W. H. Simcox . . . .	21
Parry (C. H.), French Passages . . . .	35	Tancock (C. C.), Caesar . . . . .	19
Pearson (C. H.), English History . . .	9	Taylor (R. W.), Short Greek Syntax .	23
Percival (J.), Helps for School Life .	32	— Stories from Ovid . . . . .	19
Persius. By A. Pretor . . . . .	21	— Xenophon's Agesilaus . . . . .	26
Phillipotts (J. S.), Homer's Iliad . .	27	— Xenophon's Anabasis . . . . .	26
— Shakspeare's Tempest . . . . .	5	— Works by . . . . .	38
— Stories from Herodotus . . . . .	25	Terence . . . . .	21
Plays from Terence . . . . .	21	Tidmarsh (W.), English Grammar . .	7
Powell (F. York), English History . .	8	Thompson (F. E.), Greek Syntax . .	24
Pretor (A.), Persii Satire . . . . .	21	Thucydides . . . . .	29
— Translation at Sight . . . . .	20, 25	Turner (E. J.), Goethe's Faust . . .	34
Priestland (E.), Greek Prepositions . .	23	VECQUERAY (J.), German Accidence .	14
Purnell (E. K.), Shakspeare's Othello .	5	Vergil . . . . .	14, 17, 18
Pusey (E. B.), Prayers for Schoolboy .	33	WAITE (R.), Duke of Wellington . .	10
RANSOME (C.), Political Hist. of Eng.	10	Way of Life . . . . .	33
— Skeleton Outline . . . . .	10	Wharton (E. R.), <i>Etyma Græca</i> . . .	25
— Constitutional Government . . . .	8	Whitelaw (R.), Shakspeare's Coriolanus	5
Raven (J. H.), Latin Gram. Papers . .	16	— Sophocles . . . . .	29
Reynolds (S. H.), Iliad of Homer . .	27	Willert (P. F.), Reign of Lewis XI. .	9
Richardson (G.), Conic Sections . . .	12	Wilson (R. K.), Modern English Law .	9
Rigg (A.), Intro. to Chemistry . . . .	11	Woods (H. G.), Herodoti Historia . .	28
Ritchie (F.), <i>Fabulae Faciles</i> . . . .	16	Wordsworth (Bp.), Greek Testament .	30
— First Steps and Key . . . . .	16	Wormell (R.), Dynamics . . . . .	13
— Greek Method and Key . . . . .	22	Worthington (A. M.), Physics . . . .	11
— Greek Verbs . . . . .	22	Wyatt (H. H.), St. Matthew's Gospel .	32
Rivington's Mathematical Series . . .	12, 13	XENOPHON . . . . .	26



## RIVINGTON'S EDUCATIONAL LIST

### *Arnold's Henry's First*

*Latin Book.* By C. G. GEPF. 3s.  
[The original Edition is still on sale.]

### *Arnold's Latin Prose*

*Composition.* By G. G. BRADLEY. 5s.  
[The original Edition is still on sale.]

### *First Latin Writer.* By

G. L. BENNETT. 3s. 6d.

Or separately—

*First Latin Exercises.* 2s. 6d.

*Latin Accidence.* 1s. 6d.

### *Second Latin Writer.*

By G. L. BENNETT. 3s. 6d.

### *Easy Latin Stories for*

*Beginners.* By G. L. BENNETT.  
2s. 6d.

### *Second Latin Reading*

*Book.* By G. L. BENNETT. 2s. 6d.

### *Selections from Cæsar.*

By G. L. BENNETT. 2s.

### *Selections from Vergil.*

By G. L. BENNETT. 1s. 6d.

### *Cæsar de Bello Gallico.*

Books I.—III. By J. MURPHY-  
WEATHER and C. TANCOCK. 3s. 6d.  
Book I. separately, 2s.

### *Gradatim.* An Easy Latin Trans-

lation Book. By H. R. HEATLEY and  
H. N. KINGDON. 1s. 6d.

### *Excerpta Facilia.* A Second

Latin Translation Book. By H. R.  
HEATLEY and H. N. KINGDON.  
2s. 6d.

### *First Steps in Latin.* By

F. RITCHIE. 1s. 6d.

### *Arnold's Greek Prose*

*Composition.* By E. ARBOTT. 3s. 6d.  
[The original Edition is still on sale.]

### *A Primer of Greek*

*Grammar.* By E. ARBOTT and E.  
D. MANSFIELD. 3s. 6d.

Separately—

*Syntax.* 1s. 6d. *Accidence.* 2s. 6d.

### *A Practical Greek Me-*

*thod for Beginners.* THE SIMPLE  
SENTENCE. By F. RITCHIE and E.  
H. MOORE. 3s. 6d.

### *Stories in Attic Greek.*

By F. D. MORICE. 3s. 6d.

### *A First Greek Writer.*

By A. SIDGWICK. 3s. 6d.

### *An Introduction to Greek*

*Prose Composition.* By A. SIDG-  
WICK. 5s.

### *An Introduction to Greek*

*Verse Composition.* By A. SIDG-  
WICK and F. D. MORICE. 5s.

### *Homer's Iliad.* By A. SIDG-

WICK. Books I. and II., 2s. 6d.—  
Book XXI., 1s. 6d.—Book XXII.,  
1s. 6d.

### *The Anabasis of Xeno-*

phon. By R. W. TAYLOR. Books  
I. and II., 3s. 6d. Separately, Book  
I., 2s. 6d.; Book II., 2s.—Books III.  
and IV., 3s. 6d.

### *Xenophon's Agesilaus.*

By R. W. TAYLOR. 2s. 6d.

### *Stories from Ovid in*

*Elegiac Verse.* By R. W. TAYLOR.  
3s. 6d.

### *Stories from Ovid in*

*Hexameter Verse.* By R. W. TAY-  
LOR. 2s. 6d.

Waterloo Place, Pall Mall, London.



## RIVINGTON'S EDUCATIONAL LIST

### *Catena Classicorum.*

ARISTOPHANIS COMEDIE. THE ACHARSIANS AND THE KNIGHTS. 4s.—THE WASPS. 3s. 6d.—THE CLOUDS. 3s. 6d.  
 DEMOSTHENIS ORATIONES PUBLICÆ. THE OLYNTHIACS. 2s. 6d.—THE PHILIPPICS. 3s. Or, in one volume, 4s. 6d. DE FALSA LEGATIONE. 6s.  
 DEMOSTHENIS ORATIONES PRIVATÆ. DE CORONA. 5s.  
 HERODOTI HISTORIA. Book I., 6s.—Book II., 5s.  
 HOMERI ILIAS. Books I.—XII. 6s.  
 HORATI OPERA. THE ODES, CARMEN SECULARE, and EPODES. 7s. 6d.—THE ODES. Books I.—IV. separately, 1s. 6d. each.  
 ISOCRATIS ORATIONES. AD DEMONICUM ET PANEGYRICUS. 4s. 6d.  
 JUVENALIS SATIRÆ. 5s.  
 PERSII SATIRÆ. 3s. 6d.  
 SOPHOCLES TRAGEDIÆ. THE ELECTRA. 3s. 6d.—THE AJAX. 3s. 6d.  
 TACITI HISTORIÆ. Books I. and II., 6s.—Books III., IV., and V., 6s.  
 TERENCE COMEDIÆ. ANDRIA and EUNUCHUS. 4s. 6d. Or separately, ANDRIA, 3s. 6d.—EUNUCHUS, 2s.  
 THUCYDIDIS HISTORIA. Books I. and II., 6s.—Books III. and IV., 6s.

### *Scenes from Greek Plays.*

By A. SIDGWICK. 1s. 6d. each.  
 ARISTOPHANES. THE CLOUDS. THE FROGS. THE KNIGHTS. PLUTUS.  
 EURIPIDES. IPHIGENIA IN TAURIS. THE CYCLOPS. ION. ELECTRA. ALCESTIS. BACCHÆ. HECUBA.

*Cicero de Amicitia.* By A. SIDGWICK. 2s.

*Exercises in Latin Elegiac Verse.* By C. G. GIFF. 3s. 6d.

*Eclogæ Ovidianæ.* By T. K. ARNOLD. 2s. 6d.

*A First History of England.* By LOUISE CREIGHTON. 2s. 6d.

### *Latin Texts.*

ÆNEID OF VERGIL. Books I., II., III., IV., V., VII., VIII., IX., 2s. each. Books VI., X., XI., XII., 2s. each.  
 GEORGICS OF VERGIL. Books I.—IV. 2s. each.  
 BUCOLICS OF VERGIL. 2s.  
 The above in one volume. 2s. 6d.  
 CÆSAR DE BELLO GALLICO. Books I., V., VII., VIII., 3s. each. Books II., III., IV., VI., 2s. each.  
 The above in one volume, 1s. 6d.

### *Select Plays of Shakspeare.*

RUGBY EDITION.  
 AS YOU LIKE IT. 2s.  
 MACBETH. 2s.  
 HAMLET. 2s. 6d.  
 KING LEAR. 2s. 6d.  
 ROMEO AND JULIET. 2s.  
 KING HENRY THE FIFTH. 2s.  
 A MIDSUMMER NIGHT'S DREAM. 2s.  
 KING JOHN. 2s.  
 CORIOLANUS. 2s. 6d.  
 THE TEMPEST. 2s.

### *A History of England.*

By J. F. BRIGHT.  
 Period I.—MEDIEVAL MONARCHY: A.D. 449—1485. 4s. 6d.  
 Period II.—PERSONAL MONARCHY: A.D. 1485—1688. 5s.  
 Period III.—CONSTITUTIONAL MONARCHY: A.D. 1689—1837. 7s. 6d.

### *Historical Biographies.*

SIMON DE MONTFORT. 2s. 6d.  
 THE BLACK PRINCE. 2s. 6d.  
 SIR WALTER RALEGH. 3s.  
 OLIVER CROMWELL. 3s. 6d.  
 DUKE OF MARLBOROUGH. 3s. 6d.  
 DUKE OF WELLINGTON. 3s. 6d.

### *A Handbook in Outline*

*of English History to 1852.* By ARTHUR H. D. ACLAND and CYRIL RANSOME. 6s.  
 A SKELETON OUTLINE of the above. 1s. 6d.

Waterloo Place, Pall Mall, London.

